

On the semigroup of all partial fence-preserving injections on a finite set

Ilinka Dimitrova

*Faculty of Mathematics
 and Natural Science
 South-West University "Neofit Rilski"
 2700 Blagoevgrad, Bulgaria
 ilinka_dimitrova@swu.bg*

Jörg Kopitz*

*Institute of Mathematics
 University of Potsdam
 14476 Potsdam, Germany
 kopitz@uni-potsdam.de*

Received 20 October 2016
 Accepted 26 November 2016
 Published 12 January 2017

Communicated by L. Bokut

For $n \in \mathbb{N}$, let $X_n = \{a_1, a_2, \dots, a_n\}$ be an n -element set and let $\mathbf{F} = (X_n; <_f)$ be a fence, also called a zigzag poset. As usual, we denote by I_n the symmetric inverse semigroup on X_n . We say that a transformation $\alpha \in I_n$ is *fence-preserving* if $x <_f y$ implies that $x\alpha <_f y\alpha$, for all x, y in the domain of α . In this paper, we study the semigroup PFI_n of all partial fence-preserving injections of X_n and its subsemigroup $IF_n = \{\alpha \in PFI_n : \alpha^{-1} \in PFI_n\}$. Clearly, IF_n is an inverse semigroup and contains all regular elements of PFI_n . We characterize the Green's relations for the semigroup IF_n . Further, we prove that the semigroup IF_n is generated by its elements with $\text{rank} \geq n-2$. Moreover, for $n \in 2\mathbb{N}$, we find the least generating set and calculate the rank of IF_n .

Keywords: Finite transformation semigroup; fence-preserving transformations; inverse semigroup; Green's relations; generators; rank.

Mathematics Subject Classification: 20M20, 20M18

1. Introduction and Preliminaries

For $n \in \mathbb{N}$, let $X_n = \{a_1, a_2, \dots, a_n\}$ be an n -element set. As usual, we denote by I_n the symmetric inverse semigroup on X_n , i.e. the partial one-to-one transformation semigroup on X_n under composition of mappings. The importance of I_n to inverse

*Corresponding author.

semigroup theory may be likened to that of the symmetric group S_n to group theory. Every finite inverse semigroup S is embeddable in I_n , the analogue of Cayley's theorem for finite groups, and to the regular representation of finite semigroups. Thus, just as the study of symmetric, alternating and dihedral groups has made a significant contribution to group theory, so has the study of various subsemigroups of I_n , see for example [1, 3, 5, 6, 13].


Let $\mathbf{F} = (X_n; <_f)$ be a *fence*, also called a *zigzag poset*, i.e. a partially ordered set in which the order relation forms a path with alternating orientations:

$$a_1 <_f a_2 >_f a_3 <_f \cdots a_n$$

or

$$a_1 >_f a_2 <_f a_3 >_f \cdots a_n.$$

Every element of \mathbf{F} is either maximal or minimal. A fence \mathbf{F} is called an *up-fence* (respectively a *down-fence*) if $a_1 <_f a_2$ (respectively $a_1 >_f a_2$). In this paper, without loss of generality, we consider an up-fence.



Several authors have investigated the number of order-preserving maps from fences to themselves, or to fences of other sizes, see for example [2, 4, 10, 11]. Recently, regular semigroups of transformations preserving a fence were characterized in [8, 12].

We begin by recalling some notations and definitions that will be used in the paper. For standard concepts in semigroup and symmetric inverse semigroup theory, see for example [7] and [9]. We denote by $\text{dom } \alpha$ and $\text{im } \alpha$ the domain and the image (range) of $\alpha \in I_n$, respectively. The natural number $\text{rank } \alpha := |\text{im } \alpha|$ is called the rank of α . The inverse element of α is denoted by α^{-1} . For a subset $Y \subseteq X_n$, we denote by $\text{id}|_Y$ the identity mapping on Y . Clearly, if $Y = X_n$ then $\text{id}|_{X_n} =: \text{id}$ is the identity mapping on X_n . For a subset $A \subseteq I_n$, we denote by $\langle A \rangle$ the subsemigroup of I_n generated by A . We say that a transformation $\alpha \in I_n$ is *fence-preserving* if $x <_f y$ implies that $x\alpha <_f y\alpha$, for all $x, y \in \text{dom } \alpha$. We denote by PFI_n the subsemigroup of I_n of all partial fence-preserving injections of X_n . Note that the semigroup PFI_n is not inverse. For example,

$$\alpha = \begin{pmatrix} 1 & 2 & 4 & 5 & 6 \\ 3 & 2 & 6 & 5 & 4 \end{pmatrix} \in PFI_6, \quad \text{but } \alpha^{-1} = \begin{pmatrix} 2 & 3 & 4 & 5 & 6 \\ 2 & 1 & 6 & 5 & 4 \end{pmatrix} \notin PFI_6.$$

Let IF_n be the set of all $\alpha \in PFI_n$ such that $\alpha^{-1} \in PFI_n$. Clearly, IF_n is the set of all $\alpha \in PFI_n$ with $x <_f y$ if and only if $x\alpha <_f y\alpha$, for all $x, y \in \text{dom } \alpha$.

Hence, IF_n is an inverse subsemigroup of PFI_n and contains all regular elements of PFI_n . In Sec. 2, we characterize the Green's relations for the inverse semigroup IF_n . Further, we prove that the semigroup IF_n is generated by its elements with rank $\geq n - 2$. Moreover, for $n \in 2\mathbb{N}$ we find the least generating set and calculate the rank of IF_n .

2. Green's Relations

In this section, we characterize the Green's relations \mathcal{R} , \mathcal{L} , \mathcal{H} and \mathcal{J} on IF_n . Since IF_n is an inverse subsemigroup of I_n , for $\alpha, \beta \in IF_n$, it holds:

- (1) $\alpha\mathcal{L}\beta$ if and only if $\text{im } \alpha = \text{im } \beta$.
- (2) $\alpha\mathcal{R}\beta$ if and only if $\text{dom } \alpha = \text{dom } \beta$.
- (3) $\alpha\mathcal{H}\beta$ if and only if $\text{dom } \alpha = \text{dom } \beta$ and $\text{im } \alpha = \text{im } \beta$.

It remains to describe the relation \mathcal{J} , since this relation is different for the semigroups I_n and IF_n . For example, let

$$\alpha = \begin{pmatrix} 1 & 4 & 5 & 6 \\ 2 & 6 & 5 & 4 \end{pmatrix}, \quad \beta = \begin{pmatrix} 1 & 2 & 5 & 6 \\ 5 & 6 & 1 & 2 \end{pmatrix} \in IF_6.$$

Then $\text{rank } \alpha = \text{rank } \beta$, but α and β are not \mathcal{J} related.

Definition 2.1. For $Y \subseteq X$, let Y_S be the set of all subsets

$$\{a_i, a_{i+1}, \dots, a_{i+r}\} \quad (i, r \in \{1, \dots, n\})$$

of Y such that $a_{i-1} \notin Y$ (or $i = 1$) and $a_{i+r+1} \notin Y$ (or $i + r = n$).

Definition 2.2. Let $\alpha \in IF_n$ and let $k \in \mathbb{N}$. Then we put

$$\alpha(k) := \{A \in (\text{dom } \alpha)_S : |A| = k\},$$

$$\alpha^o(2k+1) := \{\{a_i, \dots, a_{i+2k}\} \in \alpha(2k+1) : i \in 2\mathbb{N} - 1\}.$$

Note that $\alpha^o(2k+1) \subseteq \alpha(2k+1)$.

For a set M of natural numbers, let $\max M$ (let $\min M$) be the greatest (the least) natural number in M with respect to the natural order in \mathbb{N} .

Proposition 2.3. Let $\alpha, \beta \in IF_n$. Then the following statements are equivalent:

- (i) $\alpha\mathcal{J}\beta$.
- (ii) $|\alpha(k)| = |\beta(k)|$ and $|\alpha^o(2k+1)| = |\beta^o(2k+1)|$ for all $k \in \mathbb{N}$.

Proof. Suppose that $\alpha\mathcal{J}\beta$. Then there are $\gamma, \delta, \gamma_1, \delta_1 \in IF_n$ such that $\beta = \gamma\alpha\delta$ and $\alpha = \gamma_1\beta\delta_1$. We have $\text{rank } \alpha = \text{rank } \beta$ since $IF_n \leq I_n$. Then from $\alpha = \gamma_1\beta\delta_1$ and $\beta = \gamma\alpha\delta$, we obtain $|(\text{dom } \alpha)_S| = |(\text{dom } \beta)_S|$, and in particular, $|\alpha(k)| = |\beta(k)|$ for all $k \in \mathbb{N}$. Moreover, if $k \in \mathbb{N}$ and $B \in \beta(k)$, then we observe $B\gamma \in (\text{dom } \alpha)_S$ and thus $B\gamma \in \alpha(k)$.

Let $k \in 2\mathbb{N} + 1$ and $B := \{a_i, \dots, a_{i+k-1}\} \in \beta^o(k)$ (for some $i \in 2\mathbb{N} - 1$). We have $B\gamma \in \alpha(k)$ and we will show that $B\gamma \in \alpha^o(k)$. Since i is odd, we have $a_i <_f a_{i+1} >_f \dots <_f a_{i+k-2} >_f a_{i+k-1}$. This implies $a_i\gamma <_f a_{i+1}\gamma >_f \dots <_f a_{i+k-2}\gamma >_f a_{i+k-1}\gamma$ and there is $l \in \{1, \dots, n\}$ with either $a_i\gamma = a_l$ and $a_{i+k-1}\gamma = a_{l+k-1}$ or $a_i\gamma = a_{l+k-1}$ and $a_{i+k-1}\gamma = a_l$. This gives $a_l <_f a_{l+1}$ and $l \in 2\mathbb{N} - 1$, and consequently, $B\gamma \in \alpha^o(k)$. This shows $|\beta^o(k)| \leq |\alpha^o(k)|$. Dually, we can verify the converse inequation. Thus, $|\alpha^o(k)| = |\beta^o(k)|$.

Conversely, let $|\alpha(k)| = |\beta(k)|$ and $|\alpha^o(2k+1)| = |\beta^o(2k+1)|$ for all $k \in \mathbb{N}$. Then for all $k \in \mathbb{N}$, there is a bijection $f_k : \beta(k) \rightarrow \alpha(k)$ such that $f_{2k+1}(B) \in \alpha^o(2k+1)$ for all $B \in \beta^o(2k+1)$. We define now a mapping $\gamma : \text{dom } \beta \rightarrow \text{dom } \alpha$. For $k \in \mathbb{N}$, $B = \{a_i, \dots, a_{i+k-1}\} \in \beta(k)$ and $f_k(B) = \{a_l, \dots, a_{l+k-1}\}$ (with $i, l \in \{1, \dots, n\}$) let

$$a_{i+r}\gamma := \begin{cases} a_{l+r}, & \text{if } k = 1 \text{ or } i \text{ and } l \text{ have the same parity,} \\ a_{l+k-(r+1)}, & \text{otherwise} \end{cases}$$

for $0 \leq r \leq k-1$. The mapping γ is well defined since $\text{dom } \beta = \bigcup_{j=1}^p \beta(j)$, where $p := \max\{k \in \mathbb{N} : \beta(k) \neq \emptyset\}$.

We have to show that $\gamma \in IF_n$. For this let again $B = \{a_i, \dots, a_{i+k-1}\} \in \beta(k)$ and $\{a_l, \dots, a_{l+k-1}\} = f_k(B)$ for some $i, l, k \in \{1, \dots, n\}$.

We consider here the case $i \in 2\mathbb{N} - 1$, the case $i \in 2\mathbb{N}$ can be handled in the same matter. Suppose that $k \in 2\mathbb{N} + 1$. Then $B \in \beta^o(k)$ and $f_k(B) \in \alpha^o(k)$, i.e. $l \in 2\mathbb{N} - 1$. Since i and l are odd, we have $a_i <_f a_{i+1} >_f \dots <_f a_{i+k-2} >_f a_{i+k-1}$ and $a_l <_f a_{l+1} >_f \dots <_f a_{l+k-2} >_f a_{l+k-1}$, i.e. $a_i\gamma <_f a_{i+1}\gamma >_f \dots <_f a_{i+k-2}\gamma >_f a_{i+k-1}\gamma$.

Now, suppose that $k \in 2\mathbb{N}$. Since i is odd, we have $a_i <_f a_{i+1} >_f \dots >_f a_{i+k-2} <_f a_{i+k-1}$. If l is odd, then $a_l <_f a_{l+1} >_f \dots >_f a_{l+k-2} <_f a_{l+k-1}$, i.e. $a_i\gamma <_f a_{i+1}\gamma >_f \dots >_f a_{i+k-2}\gamma <_f a_{i+k-1}\gamma$. If l is even then $a_l >_f a_{l+1} <_f \dots <_f a_{l+k-2} >_f a_{l+k-1}$, i.e. $a_{i+k-1}\gamma >_f a_{i+k-2}\gamma <_f \dots <_f a_{i+1}\gamma >_f a_i\gamma$.

This shows that $\gamma \in PFI_n$. Let $r \in \{1, \dots, n-1\}$ with $a_r, a_{r+1} \in A$ for some $A \in (\text{im } \gamma)_S$. We observe that $\{B\gamma : B \in (\text{dom } \beta)_S\} = (\text{dom } \alpha)_S$. Thus, there is $B \in (\text{dom } \beta)_S$ such that $B\gamma = A$ and there is $s \in \{1, \dots, n\}$ with $a_s = a_r\gamma^{-1}$. Then $a_{r+1}\gamma^{-1} \in \{a_{s+1}, a_{s-1}\}$. If r is odd then $a_r <_f a_{r+1}$. Assume that $a_r\gamma^{-1} >_f a_{r+1}\gamma^{-1}$. Then s is even, i.e. $a_s\gamma >_f a_{s+1}\gamma$ (if $a_{r+1}\gamma^{-1} = a_{s+1}$) and $a_{s-1}\gamma <_f a_s\gamma$ (if $a_{r+1}\gamma^{-1} = a_{s-1}$). This gives $a_r >_f a_{r+1}$, a contradiction. If r is even then $a_r >_f a_{r+1}$ and we obtain $a_r\gamma^{-1} >_f a_{r+1}\gamma^{-1}$ by the same arguments. This provides $\gamma^{-1} \in PFI_n$, i.e. $\gamma \in IF_n$.

Finally, we define $\delta : \text{im } \alpha \rightarrow \text{im } \beta$ by

$$\delta := \alpha^{-1}\gamma^{-1}\beta.$$

Since $\alpha, \beta, \gamma \in IF_n$, we have $\delta = \alpha^{-1}\gamma^{-1}\beta \in IF_n$.

There holds $\beta = \gamma\alpha\delta$. In fact, for $a \in \text{dom } \beta$, we obtain $a\gamma\alpha\delta = a\gamma\alpha\alpha^{-1}\gamma^{-1}\beta = a\beta$ since $\text{dom } \alpha = \text{im } \gamma$ and $\text{dom } \gamma = \text{dom } \beta$. \square

3. Generating Sets

For convenience, we arrange such that X_n is the set of the first positive integers n for some $n \in \mathbb{N}$, i.e. $X_n = \{1, \dots, n\}$ with

$$1 <_f 2 >_f 3 <_f \dots n.$$

Clearly, the minimal elements of the fence $\mathbf{F} = (X_n, <_f)$ are odd and maximal elements are even. For $a, b \in X_n$, we will write $a \equiv b \pmod{2}$ or shorter $a \equiv_2 b$ if a and b have the same parity. Further, we denote by ε_i the identity mapping on $X_n \setminus \{i\}$ for $i = 1, \dots, n$, i.e. $\varepsilon_i := id|_{X_n \setminus \{i\}}$.

Notation 3.1. Let us put

$$J := \{\alpha \in IF_n : \text{rank } \alpha \geq n - 2\}.$$

The aim of this section is to show that J is a generating set for the semigroup IF_n . Note, $\varepsilon_i^{-1} = \varepsilon_i \in J$ for $1 \leq i \leq n$.

Lemma 3.2. Let $m, p \in \mathbb{N}$ with $m + p \leq n$ and $m \equiv_2 m + p$. Then

$$\alpha = \begin{pmatrix} 1 & \dots & m-2 & m & \dots & m+p & m+p+2 & \dots & n \\ 1 & \dots & m-2 & m+p & \dots & m & m+p+2 & \dots & n \end{pmatrix} \in J$$

and $\alpha^{-1} \in J$.

Proof. By simple calculations, one can see that $\alpha \in IF_n$. Since $\text{rank } \alpha = n - 2$ and $\alpha^{-1} = \alpha$, we obtain $\alpha, \alpha^{-1} \in J$. \square

Lemma 3.3. Let $m, p \in \mathbb{N}$ such that $m + p + 2 \leq n$. Then

$$\alpha = \begin{pmatrix} 1 & \dots & m-2 & m & \dots & m+p & m+p+4 & \dots & n \\ 1 & \dots & m-2 & m+2 & \dots & m+p+2 & m+p+4 & \dots & n \end{pmatrix} \in \langle J \rangle$$

and $\alpha^{-1} \in \langle J \rangle$.

Proof. We have to consider two cases.

(1) Suppose that p is even. Then $m \equiv_2 m + p$ and we consider the following transformations with $\text{rank } \geq n - 2$:

$$\beta_1 = \begin{pmatrix} 1 & \dots & m-2 & m & \dots & m+p+2 & m+p+4 & \dots & n \\ 1 & \dots & m-2 & m+p+2 & \dots & m & m+p+4 & \dots & n \end{pmatrix}$$

and

$$\beta_2 = \begin{pmatrix} 1 & \dots & m & m+2 & \dots & m+p+2 & m+p+4 & \dots & n \\ 1 & \dots & m & m+p+2 & \dots & m+2 & m+p+4 & \dots & n \end{pmatrix}.$$

Clearly, $\beta_1, \beta_2 \in J$ by Lemma 3.2 and it is easy to verify that

$$\alpha = \beta_1 \beta_2 \varepsilon_m \text{ and } \alpha^{-1} = \varepsilon_m \beta_2 \beta_1,$$

where $\varepsilon_m \in J$. Thus, we obtain $\alpha, \alpha^{-1} \in \langle J \rangle$.

(2) Now suppose that p is odd. Then $m \not\equiv_2 m+p$ and we consider the following transformations with $\text{rank} \geq n-2$:

$$\beta_3 = \begin{pmatrix} 1 & \cdots & m-2 & m & \cdots & m+p+1 & m+p+3 & \cdots & n \\ 1 & \cdots & m-2 & m+p+1 & \cdots & m & m+p+3 & \cdots & n \end{pmatrix}$$

and

$$\beta_4 = \begin{pmatrix} 1 & \cdots & m-1 & m+1 & \cdots & m+p+2 & m+p+4 & \cdots & n \\ 1 & \cdots & m-1 & m+p+2 & \cdots & m+1 & m+p+4 & \cdots & n \end{pmatrix}.$$

Clearly, $\beta_3, \beta_4 \in J$ by Lemma 3.2 and it is easy to verify that

$$\alpha = \beta_3\beta_4 \text{ and } \alpha^{-1} = \beta_4\beta_3.$$

Thus, $\alpha, \alpha^{-1} \in \langle J \rangle$. □

Corollary 3.4. *Let $m, p, k \in \mathbb{N}$ such that $m+p+2k \leq n$. Then*

$$\alpha = \begin{pmatrix} 1 & \cdots & m-2 & m & \cdots & m+p & m+p+2k+2 & \cdots & n \\ 1 & \cdots & m-2 & m+2k & \cdots & m+p+2k & m+p+2k+2 & \cdots & n \end{pmatrix} \in \langle J \rangle$$

and $\alpha^{-1} \in \langle J \rangle$.

Proof. For $0 \leq i < k$ we define the transformations

$$\beta_i = \begin{pmatrix} 1 & \cdots & m+2i-2 & m+2i & \cdots & m+p+2i & m+p+2i+4 & \cdots & n \\ 1 & \cdots & m+2i-2 & m+2i+2 & \cdots & m+p+2i+2 & m+p+2i+4 & \cdots & n \end{pmatrix}.$$

Note that $\beta_i, \beta_i^{-1} \in \langle J \rangle$ ($0 \leq i < k$) by Lemma 3.3. It is easy to verify that $\alpha = \beta_0 \cdots \beta_{k-1} \in \langle J \rangle$ and $\alpha^{-1} = \beta_{k-1}^{-1} \cdots \beta_0^{-1} \in \langle J \rangle$. □

Lemma 3.5. *Let $m, p \in \mathbb{N}$ such that p is odd and $m+p+1 \leq n$. Then*

$$\alpha = \begin{pmatrix} 1 & \cdots & m-2 & m & \cdots & m+p & m+p+3 & \cdots & n \\ 1 & \cdots & m-2 & m+p+1 & \cdots & m+1 & m+p+3 & \cdots & n \end{pmatrix} \in \langle J \rangle$$

and $\alpha^{-1} \in \langle J \rangle$.

Proof. We define a transformation

$$\beta_1 = \begin{pmatrix} 1 & \cdots & m-2 & m & \cdots & m+p+1 & m+p+3 & \cdots & n \\ 1 & \cdots & m-2 & m+p+1 & \cdots & m & m+p+3 & \cdots & n \end{pmatrix}.$$

Clearly, $\beta_1 \in J$ by Lemma 3.2. Then we can verify that $\alpha = \beta_1 \varepsilon_m \in \langle J \rangle$ and $\alpha^{-1} = \varepsilon_m \beta_1 \in \langle J \rangle$. □

Corollary 3.6. *Let $m, p, k \in \mathbb{N}$ such that p is odd and $m+p+2k-1 \leq n$. Then*

$$\alpha = \begin{pmatrix} 1 & \cdots & m-2 & m & \cdots & m+p & m+p+2k+1 & \cdots & n \\ 1 & \cdots & m-2 & m+p+2k-1 & \cdots & m+2k-1 & m+p+2k+1 & \cdots & n \end{pmatrix} \in \langle J \rangle$$

and $\alpha^{-1} \in \langle J \rangle$.

Proof. Let

$$\beta_1 = \begin{pmatrix} 1 & \cdots & m-2 & m & \cdots & m+p & m+p+2k & \cdots & n \\ 1 & \cdots & m-2 & m+2k-2 & \cdots & m+p+2k-2 & m+p+2k & \cdots & n \end{pmatrix}$$

and

$$\beta_2 = \begin{pmatrix} 1 & \cdots & m+2k-4 & m+2k-2 & \cdots & m+p+2k-2 & m+p+2k+1 & \cdots & n \\ 1 & \cdots & m+2k-4 & m+p+2k-1 & \cdots & m+2k-1 & m+p+2k+1 & \cdots & n \end{pmatrix}.$$

Note that $\beta_1 \in \langle J \rangle$ (by Corollary 3.4) and $\beta_2 \in \langle J \rangle$ (by Lemma 3.5). It is easy to verify that $\alpha = \beta_1\beta_2$ and $\alpha^{-1} = \beta_2^{-1}\beta_1^{-1}$, and thus $\alpha, \alpha^{-1} \in \langle J \rangle$. \square

Lemma 3.7. *Let $m, p, k \in \mathbb{N}$ such that p is even and $m + p + 2k \leq n$. Then*

$$\alpha = \begin{pmatrix} 1 & \cdots & m-2 & m & \cdots & m+p & m+p+2k+2 & \cdots & n \\ 1 & \cdots & m-2 & m+p+2k & \cdots & m+2k & m+p+2k+2 & \cdots & n \end{pmatrix} \in \langle J \rangle$$

and $\alpha^{-1} \in \langle J \rangle$.

Proof. Let

$$\beta_1 = \begin{pmatrix} 1 & \cdots & m-2 & m & \cdots & m+p & m+p+2k+2 & \cdots & n \\ 1 & \cdots & m-2 & m+2k & \cdots & m+p+2k & m+p+2k+2 & \cdots & n \end{pmatrix}$$

and

$$\beta_2 = \begin{pmatrix} 1 & \cdots & m+2k-2 & m+2k & \cdots & m+p+2k & m+p+2k+2 & \cdots & n \\ 1 & \cdots & m+2k-2 & m+p+2k & \cdots & m+2k & m+p+2k+2 & \cdots & n \end{pmatrix}.$$

Note that $\beta_1 \in \langle J \rangle$ (by Corollary 3.4) and $\beta_2 \in J$ (by Lemma 3.2). It is easy to verify that $\alpha = \beta_1\beta_2$ and $\alpha^{-1} = \beta_2\beta_1^{-1}$, and thus $\alpha, \alpha^{-1} \in \langle J \rangle$. \square

Lemma 3.8. *Let $Y \subseteq X_n$. Then $id|_{X_n \setminus Y} \in \langle J \rangle$.*

Proof. If $Y = \emptyset$, i.e. $X_n \setminus Y = X_n$, then $id|_{X_n} = id \in J$. Let $\emptyset \neq Y := \{i_1, \dots, i_k\} \subseteq X_n$ with $k \in \{1, \dots, n\}$. Then it is easy to verify that $id|_{X_n \setminus Y} = \varepsilon_{i_1} \cdots \varepsilon_{i_k} \in \langle J \rangle$. \square

Proposition 3.9. *Let $\alpha \in IF_n$. Then there are transformations $\eta_1, \dots, \eta_k, \eta_{k+1}, \dots, \eta_l \in J$ ($k < l \in \mathbb{N}$) such that $\eta_1^{-1}, \dots, \eta_k^{-1}, \eta_{k+1}^{-1}, \dots, \eta_l^{-1} \in J$, $\text{dom } \alpha \subseteq \text{im } (\eta_1 \cdots \eta_k)$, $\text{im } \alpha \subseteq \text{dom } (\eta_{k+1} \cdots \eta_l)$ and $x(\eta_1 \cdots \eta_k \alpha \eta_{k+1} \cdots \eta_l) \equiv_2 x$ for all $x \in \text{dom } (\eta_1 \cdots \eta_k \alpha \eta_{k+1} \cdots \eta_l)$.*

Proof. If $a \equiv_2 a\alpha$ for all $a \in \text{dom } \alpha$ then $id|_{\text{dom } \alpha} \alpha id|_{\text{im } \alpha} = \alpha$. This shows the assertion, since $id|_{\text{dom } \alpha}, id|_{\text{im } \alpha} \in \langle J \rangle$ by Lemma 3.8.

Let $a \in \text{dom } \alpha$ such that $a \not\equiv_2 a\alpha$. Then it is clear that $a-1, a+1 \notin \text{dom } \alpha$.

If a is even then we put

$$\eta = \begin{pmatrix} 1 & 3 & \cdots & a & a+2 & \cdots & n \\ a & 1 & \cdots & a-2 & a+2 & \cdots & n \end{pmatrix} \in J.$$

We observe that $\eta^{-1} \in \langle J \rangle$. Moreover, it is easy to see that $\text{im } \alpha = \text{im } (\eta\alpha)$, $x\alpha^{-1} \equiv_2 x(\eta\alpha)^{-1}$ for all $x \in \text{im } \alpha \setminus \{a\alpha\}$ and $a\alpha\alpha^{-1} = a \not\equiv_2 1 = a\eta^{-1} = a\alpha(\eta\alpha)^{-1}$. This shows that

$$|\{x \in \text{im } \alpha : x \not\equiv_2 x(\eta\alpha)^{-1}\}| = |\{x \in \text{im } \alpha : x \not\equiv_2 x\alpha^{-1}\}| - 1.$$

If a is odd, then $a\alpha$ is even and we put

$$\eta = \begin{pmatrix} 1 & \cdots & a\alpha - 2 & a\alpha & a\alpha + 2 & \cdots & n \\ 3 & \cdots & a\alpha & 1 & a\alpha + 2 & \cdots & n \end{pmatrix} \in J$$

with $\eta^{-1} \in \langle J \rangle$. By dual arguments, we obtain

$$|\{x \in \text{dom } \alpha : x \not\equiv_2 x(\alpha\eta)\}| = |\{x \in \text{dom } \alpha : x \not\equiv_2 x\alpha\}| - 1.$$

Continuing in this way, starting with the even cases, we obtain transformations $\eta_1, \dots, \eta_k, \eta_{k+1}, \dots, \eta_l \in J$ ($k < l \in \mathbb{N}$) such that $\eta_1^{-1}, \dots, \eta_k^{-1}, \eta_{k+1}^{-1}, \dots, \eta_l^{-1} \in J$ and $x(\eta_1 \dots \eta_k \alpha \eta_{k+1} \dots \eta_l) \equiv_2 x$ for all $x \in \text{dom } (\eta_1 \dots \eta_k \alpha \eta_{k+1} \dots \eta_l)$. \square

Notation 3.10. Let $\alpha \in PFI_n$ and let $A, B \in (\text{dom } \alpha)_S$ (or $A, B \in (\text{im } \alpha)_S$). Then we write $A < B$ if all elements in A are less than any element in B with respect to the natural order of \mathbb{N} . Further, we write

$$A \prec B$$

if $A < B$ and for each $C \in (\text{dom } \alpha)_S$ (for each $C \in (\text{im } \alpha)_S$, respectively) the following implication holds: $A \leq C \leq B \Rightarrow A = C$ or $B = C$.

Any transformation $\alpha \in IF_n$ with $a \equiv_2 a\alpha$ for all $a \in \text{dom } \alpha$ can be written in the following form:

Notation 3.11. Let

$$\alpha = \begin{pmatrix} A_1 & \prec \cdots \prec & A_{i-1} & \prec & A_i & \prec \cdots \prec & A_p \\ A_1 & \prec \cdots \prec & A_{i-1} & < & B_i & \cdots & B_p \end{pmatrix} \in IF_n$$

with $i \leq p \in \{1, \dots, n\}$, and $a \equiv_2 a\alpha$ for all $a \in \text{dom } \alpha$ such that $i = 1$ or

- (i) $a\alpha = a$ for all $a \in A_1 \cup \dots \cup A_{i-1}$ and
- (ii) $A_{i-1} < B_l$ for all $l \in \{i, \dots, p\}$.

Further, let

$$r_j := \min A_j, \quad s_j := \max A_j, \quad t_j := \min B_j, \quad u_j := \max B_j$$

for $1 \leq j \leq p$.

Proposition 3.12. *Let α be as in Notation 3.11. Then there exist $\omega_1, \omega_2 \in \langle J \rangle$ with $\omega_1^{-1}, \omega_2^{-1} \in \langle J \rangle$, $\text{dom } \alpha \subseteq \text{im } \omega_1$, $\text{im } \alpha \subseteq \text{dom } \omega_2$ such that $\omega_1 \alpha \omega_2$ has the form*

$$\omega_1 \alpha \omega_2 = \begin{pmatrix} A_1 & \prec \cdots \prec & A_{i-1} & \prec & A'_i & \prec & A'_{i+1} & \prec \cdots \prec & A'_p \\ A_1 & \prec \cdots \prec & A_{i-1} & \prec & B'_i & & B'_{i+1} & \cdots & B'_p \end{pmatrix} \in IF_n$$

with $a \equiv_2 a(\omega_1 \alpha \omega_2)$ for all $a \in \text{dom } (\omega_1 \alpha \omega_2)$, and $B'_i < B'_l$ for all $l \in \{i+1, \dots, p\}$ such that $i = 1$ or $a(\omega_1 \alpha \omega_2) = a$ for all $a \in A_1 \cup \dots \cup A_{i-1}$.

Proof. We will define the transformations ω_1 and ω_2 with $\text{dom } \alpha \subseteq \text{im } \omega_1$ and $\text{im } \alpha \subseteq \text{dom } \omega_2$ such that $\omega_1 \alpha \omega_2$ is the required mapping of our assertion. The concrete calculations we leave to the reader.

Let $k \in \{i, \dots, p\}$ such that $A_{i-1} \prec B_k$ if $i > 1$, and $B_k < B_l$ for all $l \in \{1, \dots, p\} \setminus \{k\}$ if $i = 1$, respectively. Note that if $k = i$ then $\omega_1 = \omega_2 = id \in J$. Thus, let $k > i$. Then we consider the following seven cases. Note that the cases are not mutually exclusive (i.e. the transformation α can satisfy more than one case), but cover all the possibilities.

(1) If $r_i \equiv_2 s_k$, then we put $\omega_2 = id$ and

$$\omega_1 = \begin{pmatrix} 1 & \cdots & r_i - 2 & r_i & \cdots & s_k & s_k + 2 & \cdots & n \\ 1 & \cdots & r_i - 2 & s_k & \cdots & r_i & s_k + 2 & \cdots & n \end{pmatrix}.$$

(2) If $r_i \not\equiv_2 s_k$ and $r_i - 2 \notin \text{dom } \alpha$ (or $r_i - 1 = 1$), then we put $\omega_2 = id$ and

$$\omega_1 = \begin{pmatrix} 1 & \cdots & r_i - 3 & r_i - 1 & \cdots & s_k & s_k + 2 & \cdots & n \\ 1 & \cdots & r_i - 3 & s_k & \cdots & r_i - 1 & s_k + 2 & \cdots & n \end{pmatrix}.$$

(3) If $r_i \not\equiv_2 s_k$ and $s_k + 2 \notin \text{dom } \alpha$ (or $s_k + 1 = n$), then we put $\omega_2 = id$ and

$$\omega_1 = \begin{pmatrix} 1 & \cdots & r_i - 2 & r_i & \cdots & s_k + 1 & s_k + 3 & \cdots & n \\ 1 & \cdots & r_i - 2 & s_k + 1 & \cdots & r_i & s_k + 3 & \cdots & n \end{pmatrix}.$$

(4) If $u_i \equiv_2 t_k$, then we put $\omega_1 = id$ and

$$\omega_2 = \begin{pmatrix} 1 & \cdots & t_k - 2 & t_k & \cdots & u_i & u_i + 2 & \cdots & n \\ 1 & \cdots & t_k - 2 & u_i & \cdots & t_k & u_i + 2 & \cdots & n \end{pmatrix}.$$

(5) If $u_i \not\equiv_2 t_k$ and $t_k - 2 \notin \text{im } \alpha$ (or $t_k - 1 = 1$), then we put $\omega_1 = id$ and

$$\omega_2 = \begin{pmatrix} 1 & \cdots & t_k - 3 & t_k - 1 & \cdots & u_i & u_i + 2 & \cdots & n \\ 1 & \cdots & t_k - 3 & u_i & \cdots & t_k - 1 & u_i + 2 & \cdots & n \end{pmatrix}.$$

(6) If $u_i \not\equiv_2 t_k$ and $u_i + 2 \notin \text{im } \alpha$ (or $u_i + 1 = n$), then we put $\omega_1 = id$ and

$$\omega_2 = \begin{pmatrix} 1 & \cdots & t_k - 2 & t_k & \cdots & u_i + 1 & u_i + 3 & \cdots & n \\ 1 & \cdots & t_k - 2 & u_i + 1 & \cdots & t_k & u_i + 3 & \cdots & n \end{pmatrix}.$$

Clearly, $\text{dom } \alpha \subseteq \text{im } \omega_1$, $\text{im } \alpha \subseteq \text{dom } \omega_2$, and $\omega_1, \omega_2 \in J$ (by Lemma 3.2) for all cases 1–6.

(7) It remains the case $r_i \not\equiv_2 s_k$ and $u_i \not\equiv_2 t_k$ and $r_i - 2, s_k + 2 \in \text{dom } \alpha$ and $t_k - 2, u_i + 2 \in \text{im } \alpha$, where $1 = r_1 \in \text{dom } \alpha$ and $1 = t_k \in \text{im } \alpha$ in the case $i = 1$.

(7.1) Let $k = i + 1$. First, we will show that $r_i = t_{i+1}$. In the case $i = 1$, it is clear. For the case $i > 1$, we have that $t_{i+1} = u_{i-1} + 2$ (since $A_{i-1} \prec B_{i+1}$ and $t_{i+1} - 2 \in \text{im } \alpha$), $r_i = s_{i-1} + 2$ (since $A_{i-1} < A_i$ and $r_i - 2 \in \text{dom } \alpha$) and $u_{i-1} + 2 = s_{i-1} + 2$ (since $a\alpha = a$ for all $a \in A_1 \cup \dots \cup A_{i-1}$). Altogether, we obtain $r_i = t_{i+1}$. Since $r_i \not\equiv_2 s_{i+1}$, we have $r_i = t_{i+1} = r_{i+1}\alpha \equiv_2 r_{i+1}$. Thus, we get $r_{i+1} \not\equiv_2 s_{i+1}$ and we put $\omega_1 = \eta_1\eta_2$ and $\omega_2 = \text{id}$, where

$$\eta_1 = \begin{pmatrix} 1 & \cdots & r_i - 2 & r_i & \cdots & s_{i+1} - 1 & s_{i+1} + 1 & \cdots & n \\ 1 & \cdots & r_i - 2 & s_{i+1} - 1 & \cdots & r_i & s_{i+1} + 1 & \cdots & n \end{pmatrix}$$

$$\eta_2 = \begin{pmatrix} 1 & \cdots & r_{i+1} - 3 & r_{i+1} - 1 & \cdots & s_{i+1} - 1 & s_{i+1} + 2 & \cdots & n \\ 1 & \cdots & r_{i+1} - 3 & s_{i+1} & \cdots & r_{i+1} & s_{i+1} + 2 & \cdots & n \end{pmatrix}.$$

Clearly, $\eta_1 \in J$ by Lemma 3.2 and $\eta_2 \in \langle J \rangle$ by Lemma 3.5. Note that $r_{i+1} - 2 \notin \text{dom } \alpha$, since otherwise $s_i = r_{i+1} - 2 \equiv_2 r_{i+1} \equiv_2 r_i$ implies $u_i \equiv_2 r_i = t_{i+1}$ which is a contradiction. Thus, it is easy to verify that $\text{dom } \alpha \subseteq \text{im } \omega_1$.

(7.2) Let $k > i + 1$. We define a transformation τ as following:

(a) If $r_{i+1} \equiv_2 s_k$, then we put

$$\tau = \begin{pmatrix} 1 & \cdots & r_{i+1} - 2 & r_{i+1} & \cdots & s_k & s_k + 2 & \cdots & n \\ 1 & \cdots & r_{i+1} - 2 & s_k & \cdots & r_{i+1} & s_k + 2 & \cdots & n \end{pmatrix}.$$

(b) If $r_{i+1} \not\equiv_2 s_k$, i.e. $r_i \equiv_2 r_{i+1}$, then we put

$$\tau = \begin{pmatrix} 1 & \cdots & r_{i+1} - 3 & r_{i+1} - 1 & \cdots & s_k & s_k + 2 & \cdots & n \\ 1 & \cdots & r_{i+1} - 3 & s_k & \cdots & r_{i+1} - 1 & s_k + 2 & \cdots & n \end{pmatrix}.$$

By Lemma 3.2, we have $\tau, \tau^{-1} \in J$. We have to verify that $r_{i+1} - 2 \notin \text{dom } \alpha$. Assume the opposite that $r_{i+1} - 2 \in \text{dom } \alpha$. Then $s_i = r_{i+1} - 2$ and thus $s_i = r_{i+1} - 2 \equiv_2 r_{i+1} \equiv_2 r_i$. Therefore, we have $r_i \equiv_2 s_i \equiv_2 t_i \equiv_2 u_i$. Moreover, we have $r_i = t_k = 1$ in the case $i = 1$. If $i > 1$ then $u_{i-1} = s_{i-1} = r_i - 2 \equiv_2 r_i$ and $u_{i-1} \equiv_2 u_{i-1} + 2 = t_k$ implies $r_i \equiv_2 t_k$. Thus, we obtain $u_i \equiv_2 t_k$, a contradiction. Hence, $\text{dom } \alpha \subseteq \text{im } \tau$.

Now, we consider the transformation

$$\tau\alpha = \begin{pmatrix} A_1 & \cdots & A_{i-1} & A_i & A_{i+1}^* & \cdots & A_k^* & A_{k+1} & \cdots & A_p \\ A_1 & \cdots & A_{i-1} & B_i & B_{i+1}^* & \cdots & B_k^* & B_{k+1} & \cdots & B_p \end{pmatrix} \in IF_n$$

with $A_{i-1} \prec B_{i+1}^*$. For this transformation, we have the case 7.1. with corresponding transformations $\eta_1, \eta_2 \in \langle J \rangle$. Then we put $\omega_1 = \eta_1\eta_2\tau$ and $\omega_2 = \text{id}$ with $\omega_1^{-1}, \omega_2^{-1} \in \langle J \rangle$, $\text{dom } \alpha \subseteq \text{im } \omega_1$ and $\text{im } \alpha \subseteq \text{dom } \omega_2$. \square

Proposition 3.13. *Let α be as in Notation 3.11 with $A_{i-1} \prec B_i$. Then there exist $\omega_1, \omega_2 \in \langle J \rangle$ such that $\omega_1^{-1}, \omega_2^{-1} \in \langle J \rangle$, $\text{dom } \alpha \subseteq \text{im } \omega_1$, $\text{im } \alpha \subseteq \text{dom } \omega_2$, and*

$$\omega_1 \alpha \omega_2 = \begin{pmatrix} A_1 & \prec \cdots \prec & A_{i-1} & \prec & A'_i & \prec & A'_{i+1} & \prec \cdots \prec & A'_p \\ A_1 & \prec \cdots \prec & A_{i-1} & \prec & A'_i & & B'_{i+1} & \cdots & B'_p \end{pmatrix} \in IF_n$$

with $a(\omega_1 \alpha \omega_2) = a$ for all $a \in (A_1 \cup \cdots \cup A_{i-1} \cup A'_i)$.

Proof. If $a\alpha = a$ for all $a \in A_i$, then $\omega_1 = \omega_2 = \text{id}$. Let $a\alpha \neq a$ for some $a \in A_i$. Then we put

$$\eta_1 = \begin{pmatrix} 1 & \cdots & t_i - 2 & r_i \alpha & \cdots & s_i \alpha & s_i + 2 & \cdots & n \\ 1 & \cdots & t_i - 2 & r_i & \cdots & s_i & s_i + 2 & \cdots & n \end{pmatrix}, \quad \text{if } r_i \geq t_i$$

and

$$\eta_2 = \begin{pmatrix} 1 & \cdots & r_i - 2 & r_i \alpha & \cdots & s_i \alpha & u_i + 2 & \cdots & n \\ 1 & \cdots & r_i - 2 & r_i & \cdots & s_i & u_i + 2 & \cdots & n \end{pmatrix}, \quad \text{if } r_i \leq t_i.$$

Clearly, $\eta_1, \eta_2, \eta_1^{-1}, \eta_2^{-1} \in \langle J \rangle$ by Corollary 3.4 (if $r_i \alpha = t_i$) or Corollary 3.6 (if $r_i \alpha = u_i$ and $r_i \not\equiv_2 s_i$) or Lemma 3.7 (if $r_i \alpha = u_i$ and $r_i \equiv_2 s_i$). If $r_i \geq t_i$ then $\text{dom } \alpha \subseteq \text{im } \eta_1$ and we put $\omega_1 = \eta_1$ and $\omega_2 = \text{id}$. If $r_i \leq t_i$ then $\text{im } \alpha \subseteq \text{dom } \eta_2$ and we put $\omega_1 = \text{id}$ and $\omega_2 = \eta_2$. \square

From Propositions 3.9, 3.12 and 3.13 (frequently used), we obtain

Corollary 3.14. *Let $\alpha \in IF_n$. Then there exist $\omega_1, \omega_2 \in \langle J \rangle$ such that $\omega_1^{-1}, \omega_2^{-1} \in \langle J \rangle$, $\text{dom } \alpha \subseteq \text{im } \omega_1$, $\text{im } \alpha \subseteq \text{dom } \omega_2$, and $a(\omega_1 \alpha \omega_2) = a$ for all $a \in \text{dom } (\omega_1 \alpha \omega_2)$.*

Theorem 3.15. $IF_n = \langle J \rangle$.

Proof. Let $\alpha \in IF_n$. Then by Corollary 3.14, there exist $\omega_1, \omega_2 \in \langle J \rangle$ such that $\omega_1^{-1}, \omega_2^{-1} \in \langle J \rangle$, $\text{dom } \alpha \subseteq \text{im } \omega_1$, $\text{im } \alpha \subseteq \text{dom } \omega_2$, and $a(\omega_1 \alpha \omega_2) = a$ for all $a \in \text{dom } (\omega_1 \alpha \omega_2)$. Therefore, we have

$$\omega_1 \alpha \omega_2 = \varepsilon_{i_1} \cdots \varepsilon_{i_k} \in \langle J \rangle$$

(by Lemma 3.8), where $\{i_1, \dots, i_k\} = X_n \setminus \text{dom } (\omega_1 \alpha \omega_2)$, $k \in \{1, \dots, n\}$.

Finally, we obtain $\alpha \in \langle J \rangle$, since $\alpha = \omega_1^{-1} \omega_1 \alpha \omega_2 \omega_2^{-1}$. \square

4. Rank of the Semigroup IF_n for Even n

Let $n \in 2\mathbb{N}+1$. Using the GAP software, we have observed that IF_n is not generated by the set $\{\alpha \in IF_n : \text{rank } \alpha \geq n-1\}$. Moreover, there is no least generating set for IF_n . But, in the case n is even the situation is different. There is a least generating set and all its elements have rank $\geq n-1$.

Throughout this section, let $n \in 2\mathbb{N}$ and let X_n be again the up-fence $1 <_f 2 >_f \dots <_f n$. We describe the least generating set and calculate the rank of the semigroup IF_n .

Notation 4.1. We put

$$\sigma_1 := \begin{pmatrix} 1 & 3 & \cdots & n \\ n & 1 & \cdots & n-2 \end{pmatrix};$$

$$\sigma_2 := \sigma_1^{-1} = \begin{pmatrix} 1 & \cdots & n-2 & n \\ 3 & \cdots & n & 1 \end{pmatrix};$$

$$\gamma_i := \begin{pmatrix} 1 & \cdots & i-1 & i+1 & \cdots & n \\ i-1 & \cdots & 1 & i+1 & \cdots & n \end{pmatrix} \quad \text{for } i \in 2\mathbb{N}, 4 \leq i \leq n;$$

$$\delta_i := \begin{pmatrix} 1 & \cdots & i-1 & i+1 & \cdots & n \\ 1 & \cdots & i-1 & n & \cdots & i+1 \end{pmatrix} \quad \text{for } i \in 2\mathbb{N}-1, 1 \leq i \leq n-3;$$

$$G := \{id\} \cup \{\sigma_1, \sigma_2\} \cup \{\gamma_i : i \in 2\mathbb{N}, 4 \leq i \leq n\} \cup \{\delta_i : i \in 2\mathbb{N}-1, 1 \leq i \leq n-3\}.$$

Note that $\sigma_2^{-1} = \sigma_1$, $\gamma_i^{-1} = \gamma_i$, and $\delta_i^{-1} = \delta_i$.

Theorem 4.2. $IF_n = \langle G \rangle$.

Proof. From Theorem 3.15, we have $IF_n = \langle J \rangle$. It remains to show that $J \subseteq \langle G \rangle$. For this, we have to show that all transformations ε_i for $i \in \{1, \dots, n\}$ as well as all transformations which are used in Propositions 3.9, 3.12 and 3.13 belong to $\langle G \rangle$.

We observe that $\varepsilon_i = \gamma_i \gamma_i$ for $i \in 2\mathbb{N}$, $4 \leq i \leq n$ and $\varepsilon_i = \delta_i \delta_i$ for $i \in 2\mathbb{N}-1$, $1 \leq i \leq n-3$ as well as $\varepsilon_2 = \sigma_1 \sigma_2$ and $\varepsilon_{n-1} = \sigma_2 \sigma_1$.

For the transformations in Proposition 3.9, we have

$$\eta = \begin{pmatrix} 1 & 3 & \cdots & i & i+2 & \cdots & n \\ i & 1 & \cdots & i-2 & i+2 & \cdots & n \end{pmatrix} = \delta_{i+1} \sigma_1 \delta_{i-1} \in \langle G \rangle$$

if $i = a$ is even and $i < n$. If $i = n$ then $\eta = \sigma_1$. Further, we have

$$\eta = \begin{pmatrix} 1 & \cdots & i-2 & i & i+2 & \cdots & n \\ 3 & \cdots & i & 1 & i+2 & \cdots & n \end{pmatrix} = \delta_{i-1} \sigma_2 \delta_{i+1} \in \langle G \rangle$$

if $i = a\alpha$ is even.

For the transformations in Proposition 3.12, we put

$$\beta_{i,j} := \begin{pmatrix} 1 & \cdots & i-1 & i+1 & \cdots & j-1 & j+1 & \cdots & n \\ 1 & \cdots & i-1 & j-1 & \cdots & i+1 & j+1 & \cdots & n \end{pmatrix} = \beta_{i,j}^{-1}$$

for $1 \leq i < j \leq n$ and $i \equiv_2 j$. Clearly, $\beta_{i,j} \in \langle G \rangle$ since

$$\beta_{i,j} = \begin{cases} \delta_i \delta_{n-j+i+1} \delta_i, & \text{if } i \text{ and } j \text{ are odd;} \\ \gamma_j \gamma_{j-i} \gamma_j, & \text{if } i \text{ and } j \text{ are even.} \end{cases}$$

It is easy to verify that $\omega_1, \omega_2, \eta_1$ and τ are all of the form $\beta_{i,j}$ for suitable i and j .

Further, we have

$$\eta_2 = \begin{pmatrix} 1 & \cdots & i-1 & i+1 & \cdots & j-2 & j+1 & \cdots & n \\ 1 & \cdots & i-1 & j-1 & \cdots & i+2 & j+1 & \cdots & n \end{pmatrix}$$

for suitable i and j , and $i+1 \not\equiv_2 j-2$. Clearly, $\eta_2 \in \langle G \rangle$ since $\eta_2 = \beta_{i,j} \varepsilon_{i+1}$.

For the transformations in Proposition 3.13, we have $\eta_1, \eta_2 \in \langle J \rangle$ by Corollary 3.4 (if $r_i \alpha = t_i$) or Corollary 3.6 (if $r_i \alpha = u_i$ and $r_i \not\equiv_2 s_i$) or Lemma 3.7 (if $r_i \alpha = u_i$ and $r_i \equiv_2 s_i$).

For the transformation α in Corollary 3.4, we have

$$\alpha = \begin{pmatrix} 1 & \cdots & i-1 & i+1 & \cdots & j-1-2k & j+1 & \cdots & n \\ 1 & \cdots & i-1 & i+1+2k & \cdots & j-1 & j+1 & \cdots & n \end{pmatrix}$$

with $m = i+1$ and $m+p = j-1-2k$. Hence, we obtain $\alpha \in \langle G \rangle$ since

$$\alpha = \begin{cases} \beta_{i,j} \beta_{i+2k,j} \varepsilon_{i+1} \cdots \varepsilon_{i+2k-1}, & \text{if } i \equiv_2 j; \\ \beta_{i,j-1} \beta_{i+2k-1,j} \varepsilon_{i+1} \cdots \varepsilon_{i+2k-2}, & \text{if } i \not\equiv_2 j. \end{cases}$$

For the transformation α in Corollary 3.6, we have

$$\alpha = \begin{pmatrix} 1 & \cdots & i-1 & i+1 & \cdots & j-2k & j+1 & \cdots & n \\ 1 & \cdots & i-1 & j-1 & \cdots & i+2k & j+1 & \cdots & n \end{pmatrix}$$

with $m = i+1$ and $m+p = j-2k$. Then we can verify that $\alpha \in \langle G \rangle$ since $\alpha = \beta_{i,j} \varepsilon_{i+1} \cdots \varepsilon_{i+2k-1}$.

For the transformation α in Lemma 3.7, we have

$$\alpha = \begin{pmatrix} 1 & \cdots & i-1 & i+1 & \cdots & j-2k-1 & j+1 & \cdots & n \\ 1 & \cdots & i-1 & j-1 & \cdots & i+2k+1 & j+1 & \cdots & n \end{pmatrix}$$

with $m = i+1$ and $m+p = j-2k-1$. We have $\alpha \in \langle G \rangle$ since $\alpha = \beta_{i,j} \varepsilon_{i+1} \cdots \varepsilon_{i+2k}$. \square

Proposition 4.3. *The set G is the least generating set for IF_n .*

Proof. Theorem 4.2 shows that G is a generating set for IF_n . Let $\alpha, \beta \in G$ with $\alpha \neq \beta$ and $\{\alpha, \beta\} \neq \{\sigma_1, \sigma_2\}$. It is easy to verify that $\text{rank } \alpha\beta = n-2$. Moreover, we observe $\text{rank } \sigma_1^2 = \text{rank } \sigma_2^2 = n-2$. Let $\alpha = \alpha_1 \cdots \alpha_m$ with $\alpha_1, \dots, \alpha_m \in IF_n$, $2 \leq m \in \mathbb{N}$, such that $\text{rank } \alpha = n-1$. Without loss of generality, we can assume that $\alpha_i \neq id$ for $1 \leq i \leq m$. Then $\alpha_1, \dots, \alpha_m \in \{\beta \in IF_n : \text{rank } \beta = n-1\}$. Since G is a generating set for $\{\beta \in IF_n : \text{rank } \beta = n-1\}$, there is $\rho \in G$

such that $\alpha_1, \dots, \alpha_m \in \{\rho^j : j \in \mathbb{N}\}$ or $\alpha_1, \dots, \alpha_m \in \{\sigma_1, \sigma_2\}$ with $\alpha_i \neq \alpha_{i+1}$ for $1 \leq i \leq m-1$. This shows that any $\alpha \in G$ cannot be generated by a set without this α . Thus, each generating set of IF_n has to contain G and the assertion is shown. \square

Since $|G| = n + 1$ from Theorem 4.2 and Proposition 4.3, we obtain

Theorem 4.4. *Let $n \in 2\mathbb{N}$. Then $\text{rank } IF_n = n + 1$.*

Acknowledgments

The authors wish to thank the Alexander von Humboldt Foundation for the financial support and Assoc. Prof. Kalcho Todorov for his essential suggestions concerning this research.

The project is supported by the Alexander von Humboldt Foundation.

References

- [1] L. Bracci and L. E. Picasso, Representations of semigroups of partial isometries, *Bull. Lond. Math. Soc.* **39** (2007) 792–802.
- [2] J. D. Currie and T. I. Visentin, The number of order-preserving maps of fences and crowns, *Order* **8**(2) (1991) 133–142.
- [3] I. Dimitrova and J. Koppitz, The maximal subsemigroups of the ideals of some semigroups of partial injections, *Discuss. Math. Gen. Algebra Appl.* **29** (2009) 153–167.
- [4] J. D. Farley, The number of order-preserving maps between fences and crowns, *Order* **12**(1) (1995) 5–44.
- [5] V. H. Fernandes, The monoid of all injective orientation-preserving partial transformations on a finite chain, *Comm. Algebra* **32** (2000) 3401–3426.
- [6] O. Ganyushkin and V. Mazorchuk, On the Structure of IO_n , *Semigroup Forum* **66** (2003) 455–483.
- [7] J. M. Howie, *Fundamentals of Semigroup Theory* (Clarendon Press, Oxford, 1995).
- [8] K. Jendana and R. Srithus, Coregularity of order-preserving self-mapping semigroups of fences, *Commun. Korean Math. Soc.* **30** (2015) 249–361.
- [9] S. Lipscomb, *Symmetric Inverse Semigroups* Vol. 46 (AMS Mathematical Surveys and Monographs, 1996).
- [10] A. Rutkowski, The number of strictly increasing mappings of fences, *Order* **9**(1) (1992a) 31–42.
- [11] A. Rutkowski, The formula for the number of order-preserving self-mappings of a fence, *Order* **9**(2) (1992b) 127–137.
- [12] R. Tanyawong, R. Srithus and R. Chinram, Regular subsemigroups of the semigroups of transformations preserving a fence, *Asian-Eur. J. Math.* **9**(1) (2016) 1650003.
- [13] A. Umar, On the semigroups of partial one-to-one order-decreasing finite transformations, *Proc. Roy. Soc. Edinburgh A* **123** (1993) 355–363.