

Provided for non-commercial research and educational use.
Not for reproduction, distribution or commercial use.

Mathematica Balkanica

Mathematical Society of South-Eastern Europe
A quarterly published by
the Bulgarian Academy of Sciences – National Committee for Mathematics

The attached copy is furnished for non-commercial research and education use only. Authors are permitted to post this version of the article to their personal websites or institutional repositories and to share with other researchers in the form of electronic reprints.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to third party websites are prohibited.

For further information on Mathematica Balkanica visit the website of the journal
<http://www.mathbalkanica.info>

or contact:

Mathematica Balkanica - Editorial Office;
Acad. G. Bonchev str., Bl. 25A, 1113 Sofia, Bulgaria
Phone: +359-2-979-6311, Fax: +359-2-870-7273,
E-mail: balmat@bas.bg

Exact Constants in Estimations of the Error of the Quadrature Formulae of Simpson with the Averaged Moduli of Smoothness

Irena A. Parvanova and Parvan E. Parvanov

Presented by Bl. Sendov

In this paper the exact constants in estimations of the error of the quadrature formulae of Simpson are obtained. The estimations involve 4—th order in the non-periodic case and high order in the periodic case L_1 -averaged Sendov-Popov moduli of smoothness of a bounded and measurable function f .

AMS Subj. Classification: 26A15, 41A55, 65D32

Key Words: quadrature formulas of Simpson, averaged moduli of smoothness, exact constants in estimation

1. Introduction

Let $M[0, 1]$ be the set of all bounded and measurable on $[0, 1]$ functions and Π_k be the set of all algebraic polynomials of degree at most k . The k —th order local modulus of smoothness of $f \in M[0, 1]$ at the point $x \in [0, 1]$ with a step $\delta \in [0, \frac{1}{k}]$ is a function (see Definition 1.4 of [2])

$$(1.1) \quad \omega_k(f, x; \delta) \stackrel{\text{def}}{=} \sup \left\{ |\Delta_{v, [0, 1]}^k f(t)| \ ; \ t, t + kv \in \left[x - \frac{k\delta}{2}, x + \frac{k\delta}{2} \right] \right\},$$

where

$$\Delta_{t, [0, 1]}^k f(x) \stackrel{\text{def}}{=} \begin{cases} \sum_{i=0}^k (-1)^{k-i} \binom{k}{i} f(x + it) & \text{if } x, x + kt \in [0, 1]; \\ 0 & \text{otherwise.} \end{cases}$$

The k -th order L_1 -averaged Sendov-Popov modulus of smoothness of a function f bounded and measurable on $[0,1]$ is (see Definition 1.5 of [2])

$$(1.2) \quad \tau_k(f, \delta)_{L_1[0,1]} \stackrel{\text{def}}{=} \|\omega_k(f, \cdot; \delta)\|_{L_1[0,1]}.$$

We consider the well-known quadrature formulae of Simpson

$$(1.3) \quad \int_0^1 f(x)dx \approx Q_1^2(f) = \frac{1}{6} \left(f(0) + 4f\left(\frac{1}{2}\right) + f(1) \right)$$

and the n -composite quadrature formulae of Simpson

$$(1.4) \quad \int_0^1 f(x)dx \approx Q_n^2(f) = \frac{h}{6} \left(f(0) + 2 \sum_{i=1}^{n-1} f(x_i) + 4 \sum_{i=1}^n f\left(\frac{x_{i-1} + x_i}{2}\right) + f(1) \right),$$

where $x_i = ih$ and $h = \frac{1}{n}$.

We estimate the error

$$(1.5) \quad R_n^2(f) \stackrel{\text{def}}{=} \left| Q_n^2(f) - \int_0^1 f(x)dx \right|.$$

Our aim is to find for some fixed k and $\alpha > 0$, constants $c_{k,\alpha}$ and $\tilde{c}_{k,\alpha}$ such that

$$(1.6) \quad c(k, \alpha) = \max_{f \in M[0,1], n \in \mathbb{N}} \left\{ \frac{R_n^2(f)}{\tau_k(f, \alpha h)_{L_1[0,1]}} \right\}$$

and

$$(1.7) \quad \tilde{c}(k, \alpha) = \sup_{f \in C^\infty[0,1], n \in \mathbb{N}} \left\{ \frac{R_n^2(f)}{\tau_k(f, \alpha h)_{L_1[0,1]}} \right\}.$$

In Section 2 we find the exact constants $c(4, \frac{1}{4})$ and $\tilde{c}(4, \frac{1}{4})$ for non-periodic functions. In Sections 3 and 4 we consider periodic functions (1-periodic for simplicity). We find for these functions $c(k, \alpha)$ and $\tilde{c}(k, \alpha)$, where $k \in \mathbb{N}$ and $\alpha \geq \frac{1}{2}$ and $c(2, \frac{1}{4})$ and $\tilde{c}(2, \frac{1}{4})$. The methods in Sections 3 and 4 are applicable for other n -composite quadrature formulae Q_n^s of Newton-Cotes type, constructed with respect to s knots.

2. Estimations of the error of the quadrature formulae for non-periodic functions

In this section we consider non-periodic functions and a step $\alpha = \frac{1}{4}$. The Quadrature Formulae of Simpson is exact in Π_3 . Then (see [1]), we can consider $k = 4$. We first prove the following lemma.

Lemma 2.1. *Let $f \in M[0, 1]$ and $n \in \mathbb{N}$. Then*

$$R_n^2(f) \leq \frac{4}{9} \tau_4 \left(f, \frac{1}{4n} \right)_{L_1[0,1]}.$$

Proof. Using (1.5) and (1.4) for $[a, b] = [0, 1]$ we have

$$\begin{aligned} (2.1) \quad R_n^2(f) &= \left| \sum_{i=0}^{n-1} \int_{x_i}^{x_{i+1}} \left(f(x) - \frac{1}{6n} \left(f(x_i) + 4 \left(\frac{x_{i+1} + x_i}{2} \right) + f(x_{i+1}) \right) \right) dx \right| \\ &= \left| \sum_{i=0}^n \frac{4}{6} \int_0^{\frac{h}{4}} \Delta_t^4 f(x_i) dt + \frac{4}{6} \int_0^{\frac{h}{4}} \Delta_{-t}^4 f(x_{i+1}) dt + \frac{4}{9} \int_0^{\frac{h}{4}} \Delta_t^4 f \left(\frac{x_i + x_{i+1}}{2} - 2t \right) dt \right| \\ &\leq \sum_{i=0}^n \frac{4}{6} \left| \int_0^{\frac{h}{4}} \Delta_t^4 f(x_i) dt \right| + \frac{4}{6} \left| \int_0^{\frac{h}{4}} \Delta_{-t}^4 f(x_{i+1}) dt \right| \\ &\quad + \frac{4}{9} \left| \int_0^{\frac{h}{4}} \Delta_t^4 f \left(\frac{x_i + x_{i+1}}{2} - 2t \right) dt \right|. \end{aligned}$$

From (1.1) we have the next four inequalities are true for $t \in [0, \frac{h}{4}]$:

$$(2.2) \quad |\Delta_t^4 f(x_i)| \leq \omega_4 \left(f, x_i + 2t; \frac{h}{4} \right);$$

$$(2.3) \quad |\Delta_{-t}^4 f(x_{i+1})| \leq \omega_4 \left(f, x_{i+1} - 2t; \frac{h}{4} \right);$$

$$(2.4) \quad \left| \Delta_t^4 f \left(\frac{x_i + x_{i+1}}{2} - 2t \right) \right| \leq \omega_4 \left(f, \frac{x_i + x_{i+1}}{2} + \frac{h}{2} - 2t; \frac{h}{4} \right);$$

$$(2.5) \quad \left| \Delta_t^4 f \left(\frac{x_i + x_{i+1}}{2} - 2t \right) \right| \leq \omega_4 \left(f, \frac{x_i + x_{i+1}}{2} - \frac{h}{2} + 2t; \frac{h}{4} \right).$$

Applying the estimatoins of (2.2)-(2.5) in (2.1) we obtain

$$\begin{aligned} R_n^2(f) &\leq \frac{4}{6} \sum_{i=0}^n \int_0^{\frac{h}{4}} \omega_4 \left(f, x_i + 2t; \frac{h}{4} \right) dt + \frac{4}{6} \sum_{i=0}^n \int_0^{\frac{h}{4}} \omega_4 \left(f, x_{i+1} - 2t; \frac{h}{4} \right) dt \\ &\quad + \frac{2}{9} \sum_{i=0}^n \int_0^{\frac{h}{4}} \omega_4 \left(f, \frac{x_i + x_{i+1}}{2} - \frac{h}{2} + 2t; \frac{h}{4} \right) + \omega_4 \left(f, \frac{x_i + x_{i+1}}{2} + \frac{h}{2} - 2t; \frac{h}{4} \right) dt \end{aligned}$$

$$\begin{aligned}
&= \sum_{i=0}^n \frac{1}{3} \int_{x_i}^{x_i + \frac{h}{2}} \omega_4 \left(f, x; \frac{h}{4} \right) dx + \frac{1}{3} \int_{x_i + \frac{h}{2}}^{x_{i+1}} \omega_4 \left(f, x; \frac{h}{4} \right) dx \\
&\quad + \frac{1}{9} \int_{x_i}^{x_{i+1}} \omega_4 \left(f, x; \frac{h}{4} \right) dx = \frac{4}{9} \sum_{i=0}^n \int_{x_i}^{x_{i+1}} \omega_4 \left(f, x; \frac{h}{4} \right) dx \\
&= \frac{4}{9} \tau_4 \left(f; \frac{h}{4} \right)_{L_1[0,1]}.
\end{aligned}$$

Lemma 2.1 is proved. ■

Let $n = 1$. We prove that the constant $\frac{4}{9}$ in Lemma 2.1 is exact. More then, this constant is the solution of another problem. This is (see (1.6)) the constant $c(4, \frac{1}{4})$ for non-periodic functions.

Lemma 2.2. *There is a function $f \in M[0, 1]$ such that*

$$R_1^2(f) = \frac{4}{9} \tau_4 \left(f, \frac{1}{4} \right)_{L_1[0,1]}.$$

Proof. Let $\mathbb{B}$ be the set of the binary rational numbers. For every natural number m we define

$$B_m \stackrel{\text{def}}{=} \left\{ x \in \mathbb{B} ; x = \frac{2s-1}{2^m}, s = 0, \dots, 2^{m-1} \right\}.$$

Let $c = \frac{15}{16}$. We consider the function

$$f(x) \stackrel{\text{def}}{=} \begin{cases} 1 & \text{if } x = 0 \text{ or } 1; \\ \frac{1}{6} & \text{if } x = \frac{1}{2}; \\ \frac{1}{48} c^{m-3} & \text{if } x \in B_m, m \geq 3; \\ 0 & \text{if } x = \frac{1}{4}, \frac{3}{4}, \text{ or } x \notin \mathbb{B}. \end{cases}$$

Obviously, $f \in M[0, 1]$. We prove that

$$(2.6) \quad \sup \left\{ \Delta_t^4 f(x) ; t \leq \frac{1}{4} \right\} \leq 1.$$

Let $x \in [0, \frac{1}{2})$. There are 4 cases of finite differences:

I) $\Delta_{\frac{x}{4}}^4 f(0)$; II) $\Delta_{\frac{x}{2}}^4 f(0)$; III) $\Delta_{\frac{1}{4} - \frac{x}{4}}^4 f(x)$; IV) $\Delta_t^4 f(x)$ for which 0, 1 and $\frac{1}{2}$ are not knots.

In IV) we have $\sup \left\{ \Delta_t^4 f(x) ; t \leq \frac{1}{4} \right\} \leq 8 \max \left\{ f(x) ; x \neq 0, 1, \frac{1}{2} \right\} = \frac{8}{48} < 1$.

In I), II) and III), if $x \notin \mathbb{B}$, we have $\Delta_t^4 f(x) = 1$. Else, (2.6) is equal to:

$$\text{I) } 1 - 8c^2 f(B_m) + 6cf(B_m) + f(B_m) \leq 1;$$

$$\text{II) } 1 - 8cf(B_m) + 6f(B_m) + \frac{1}{c}f(B_m) \leq 1;$$

$$\text{III) } 2f(B_m) - 8cf(B_m) + 1 \leq 1.$$

The last three inequalities are true because of the choice of c . The situation for $x \in (\frac{1}{2}, 1]$ is the same. In this way we prove (2.6).

$$\text{Hence, } \omega_4 \left(f, x; \frac{1}{4} \right) = \begin{cases} 1 & \text{if } x \neq \frac{1}{2}; \\ 3 & \text{if } x = \frac{1}{2}; \end{cases} \quad \text{and } \tau_4 \left(f; \frac{1}{4} \right)_{L_1[0,1]} = 1.$$

Applying (1.5) for this function, we have $R_1^2(f) = \frac{4}{9}$. Then Lemma 2.2 is proved. ■

Lemma 2.3. *There is a functional sequence $\{f_m\}_{m=1}^\infty$, $f_m \in C^\infty[0, 1]$ such that*

$$\lim_{m \rightarrow \infty} \frac{R_1^2(f_m)}{\tau_4 \left(f_m, \frac{1}{4} \right)_{L_1[0,1]}} = \frac{4}{9}.$$

Proof. Let f be the function of Lemma 2.2 and for $m = 1, \dots, \infty$,

$$g_m(x) \stackrel{\text{def}}{=} \begin{cases} e^{\frac{(mx)^2}{(mx)^2 - 1}} & \text{if } |mx| \leq 1; \\ 0 & \text{if } |mx| > 1. \end{cases}$$

$$\text{We set } f_m(x) \stackrel{\text{def}}{=} \sum_{s=0}^{2m} f\left(\frac{s}{2m}\right) g_{2^{2m+1}}\left(x - \frac{s}{2m}\right).$$

For this functional sequence we have

$$f_m \in C^\infty[0, 1], f_m(x) = f(x) \forall x \in \bigcup_{i=1}^m B_i \text{ and } \lim_{m \rightarrow \infty} f_m(x) = f(x) \forall x \in [0, 1].$$

Also for the error of the quadrature formulae and for the local modulus of f_m , we have respectively $R_1^2(f_m) \geq \frac{4}{9} - \frac{1}{2^{2m-1}}$ and

$$\omega_1\left(f_m, x; \frac{1}{4}\right) = 1 + \left(6 \cdot \frac{15}{16} + 1\right) \frac{3^{m-4} 5^{m-3}}{2^{4m-8}} = 1 + \text{const.} \left(\frac{15}{16}\right)^m, \quad x \neq \frac{1}{2}.$$

Since

$$\tau_4\left(f_m; \frac{1}{4}\right)_{L_1[0,1]} = 1 + \text{const.} \left(\frac{15}{16}\right)^m,$$

we finally obtain

$$\lim_{m \rightarrow \infty} \frac{R_1^2(f_m)}{\tau_4\left(f_m, \frac{1}{4}\right)_{L_1[0,1]}} = \frac{4}{9}.$$

Lemma 2.3 is proved. ■

Summarizing the statements from Lemma 2.1, Lemma 2.2 and Lemma 2.3, we obtain the following theorem.

Theorem 2.1. *For non-periodic functions,*

$$c\left(4, \frac{1}{4}\right) = \tilde{c}\left(4, \frac{1}{4}\right) = \frac{4}{9}.$$

3. Estimations of the error of the quadrature formulae for 1-periodic functions and a step $\geq \frac{h}{2}$

In this section we consider 1-periodic functions and a step $\alpha \geq \frac{1}{2}$. There are not periodic algebraic polynomials out of Π_0 . Then (see [1]), we can make estimations with τ_k , for each $k \geq 1$. We first prove the following lemma.

Lemma 3.1. *Let $f \in M[0, 1]$ be 1-periodic, $k, n \in \mathbb{N}$ and $\alpha \geq \frac{1}{2}$. Then,*

$$R_n^2(f) \leq \frac{1}{\binom{k}{[\frac{k}{2}]}} \tau_k\left(f, \frac{\alpha}{n}\right)_{L_1[0,1]}.$$

Proof. We prove the statment for $\alpha = \frac{1}{2}$, because the modulus (1.2) is an increasing function of α .

1) Let $k = 2m$. Using (1.5) and (1.4) we have

$$R_n^2(f) = \left| \sum_{l=0}^{n-1} \frac{1}{3n} f(x_l) - \frac{1}{3} \int_{x_l - \frac{h}{2}}^{x_l + \frac{h}{2}} f(x) dx + \frac{2}{3n} f(x_l + \frac{h}{2}) - \frac{2}{3} \int_{x_l}^{x_{l+1}} f(x) dx \right|.$$

Applying in the last equality the trivial equality

$$\frac{2}{\binom{2m}{m}} \sum_{j=1}^m (-1)^j \binom{2m}{m-j} = -1,$$

we obtain

$$\begin{aligned} (3.1) \quad R_n^2(f) &= \left| \sum_{l=0}^{n-1} \frac{1}{3n} f(x_l) + \frac{2}{3 \binom{2m}{m}} \sum_{j=1}^m (-1)^j \binom{2m}{m-j} \int_{x_l - \frac{h}{2}}^{x_l + \frac{h}{2}} f(x) dx \right. \\ &\quad \left. + \frac{2}{3n} f(x_l + \frac{h}{2}) + \frac{4}{3 \binom{2m}{m}} \sum_{j=1}^m (-1)^j \binom{2m}{m-j} \int_{x_l}^{x_{l+1}} f(x) dx \right| \\ &= \left| \sum_{l=0}^{n-1} \frac{1}{3n} f(x_l) + \frac{2}{3 \binom{2m}{m}} \sum_{j=1}^m \frac{(-1)^j \binom{2m}{m-j}}{j} j \sum_{l=0}^{n-1} \int_{x_l - \frac{h}{2}}^{x_l + \frac{h}{2}} f(x) dx \right. \\ &\quad \left. + \sum_{l=0}^{n-1} \frac{2}{3n} f(x_l + \frac{h}{2}) + \frac{4}{3 \binom{2m}{m}} \sum_{j=1}^m \frac{(-1)^j \binom{2m}{m-j}}{j} j \sum_{l=0}^{n-1} \int_{x_l}^{x_{l+1}} f(x) dx \right| \\ &= \left| \sum_{l=0}^{n-1} \frac{1}{3n} f(x_l) + \frac{2}{3 \binom{2m}{m}} \sum_{j=1}^m \frac{(-1)^j \binom{2m}{m-j}}{j} \sum_{i=0}^{n-1} \sum_{s=-j}^{j-1} \int_{x_i + s \frac{h}{2}}^{x_i + (s+1) \frac{h}{2}} f(x) dx \right. \\ &\quad \left. + \sum_{l=0}^{n-1} \frac{2}{3n} f(x_l + \frac{h}{2}) + \frac{4}{3 \binom{2m}{m}} \sum_{j=1}^m \frac{(-1)^j \binom{2m}{m-j}}{j} \sum_{i=0}^{n-1} \sum_{s=-j}^{j-1} \int_{x_i + (s+1) \frac{h}{2}}^{x_i + (s+2) \frac{h}{2}} f(x) dx \right| \\ &= \left| \sum_{i=0}^{n-1} \frac{1}{3n} f(x_i) + \sum_{i=0}^{n-1} \frac{2}{3 \binom{2m}{m}} \sum_{j=1}^m \frac{(-1)^j \binom{2m}{m-j}}{j} \int_{x_i - j \frac{h}{2}}^{x_i + j \frac{h}{2}} f(x) dx \right. \\ &\quad \left. + \sum_{i=0}^{n-1} \frac{2}{3n} f(x_i + \frac{h}{2}) + \sum_{i=0}^{n-1} \frac{4}{3 \binom{2m}{m}} \sum_{j=1}^m \frac{(-1)^j \binom{2m}{m-j}}{j} \int_{x_i + \frac{h}{2} - j \frac{h}{2}}^{x_i + \frac{h}{2} + j \frac{h}{2}} f(x) dx \right| \end{aligned}$$

$$\begin{aligned}
&= \left| \sum_{i=0}^{n-1} \frac{2}{3 \binom{2m}{m}} \int_0^{\frac{h}{2}} \Delta_t^{2m} f(x_i - mt) dt \right. \\
&\quad \left. + \sum_{i=0}^{n-1} \frac{4}{3 \binom{2m}{m}} \int_0^{\frac{h}{2}} \Delta_t^{2m} f(x_i + \frac{h}{2} - mt) dt \right| \\
&\leq \sum_{i=0}^{n-1} \frac{2}{3 \binom{2m}{m}} \int_0^{\frac{h}{2}} |\Delta_t^{2m} f(x_i - mt)| dt \\
&\quad + \sum_{i=0}^{n-1} \frac{4}{3 \binom{2m}{m}} \int_0^{\frac{h}{2}} |\Delta_t^{2m} f(x_i + \frac{h}{2} - mt)| dt.
\end{aligned}$$

From (1.1) we have the next four inequalities are true for $t \in [0, \frac{h}{2}]$:

$$(3.2) \quad |\Delta_t^{2m} f(x_i - mt)| \leq \omega_{2m} \left(f, x_i + mt - m \frac{h}{2}; \frac{h}{2} \right);$$

$$(3.3) \quad |\Delta_t^{2m} f(x_i - mt)| \leq \omega_{2m} \left(f, x_i - mt + m \frac{h}{2}; \frac{h}{2} \right);$$

$$(3.4) \quad \left| \Delta_t^{2m} f \left(x_i + \frac{h}{2} - mt \right) \right| \leq \omega_{2m} \left(f, x_i + \frac{h}{2} + mt - m \frac{h}{2}; \frac{h}{2} \right);$$

$$(3.5) \quad \left| \Delta_t^{2m} f \left(x_i + \frac{h}{2} - mt \right) \right| \leq \omega_{2m} \left(f, x_i + \frac{h}{2} - mt + m \frac{h}{2}; \frac{h}{2} \right).$$

Applying the estimatoins of (3.2)-(3.5) in (3.1) we obtain

$$\begin{aligned}
(3.6) \quad R_n^2(f) &\leq \sum_{l=0}^{n-1} \frac{1}{3 \binom{2m}{m}} \int_0^{\frac{h}{2}} \left(\omega_{2m} \left(f, x_i + mt - m \frac{h}{2}; \frac{h}{2} \right) \right. \\
&\quad \left. + \omega_{2m} \left(f, x_i - mt + m \frac{h}{2}; \frac{h}{2} \right) \right) dt \\
&+ \sum_{i=0}^{n-1} \frac{2}{3 \binom{2m}{m}} \int_0^{\frac{h}{2}} \left(\omega_{2m} \left(f, x_i + \frac{h}{2} + mt - m \frac{h}{2}; \frac{h}{2} \right) \right. \\
&\quad \left. + \omega_{2m} \left(f, x_i + \frac{h}{2} - mt + m \frac{h}{2}; \frac{h}{2} \right) \right) dt \\
&= \sum_{l=0}^{n-1} \frac{1}{3m \binom{2m}{m}} \left(\int_{x_i - m \frac{h}{2}}^{x_i + m \frac{h}{2}} \omega_{2m} \left(f, x; \frac{h}{2} \right) dx + 2 \int_{x_i + \frac{h}{2} - m \frac{h}{2}}^{x_i + \frac{h}{2} + m \frac{h}{2}} \omega_{2m} \left(f, x; \frac{h}{2} \right) dx \right) \\
&= \sum_{l=0}^{n-1} \frac{1}{\binom{2m}{m}} \int_{x_i}^{x_{i+1}} \omega_{2m} \left(f, x; \frac{h}{2} \right) dx \\
&= \frac{1}{\binom{2m}{m}} \tau_{2m} \left(f; \frac{h}{2} \right)_{L_1[0,1]}.
\end{aligned}$$

2) Let $k = 2m + 1$. Using the same arguments as in 1), we have

$$\begin{aligned}
 (3.7) \quad R_n^2(f) &= \left| \sum_{i=0}^{n-1} \frac{2}{3 \binom{2m+1}{m}} \left(- \int_0^{\frac{h}{2}} \Delta_t^{2m+1} f(x_i - mt) dt \right. \right. \\
 &\quad \left. \left. + \int_0^{\frac{h}{2}} \Delta_t^{2m+1} f(x_i - (m+1)t) dt \right) \right. \\
 &\quad \left. + \sum_{i=0}^{n-1} \frac{4}{3 \binom{2m+1}{m}} \left(- \int_0^{\frac{h}{2}} \Delta_t^{2m+1} f(x_i + \frac{h}{2} - mt) dt \right. \right. \\
 &\quad \left. \left. + \int_0^{\frac{h}{2}} \Delta_t^{2m+1} f(x_i + \frac{h}{2} - (m+1)t) dt \right) \right| \\
 &= \sum_{i=0}^{n-1} \frac{2}{3 \binom{2m+1}{m}} \left(\int_0^{\frac{h}{2}} |\Delta_t^{2m+1} f(x_i - mt)| dt \right. \\
 &\quad \left. + \int_0^{\frac{h}{2}} |\Delta_t^{2m+1} f(x_i - (m+1)t)| dt \right) \\
 &\quad + \sum_{i=0}^{n-1} \frac{4}{3 \binom{2m+1}{m}} \left(\int_0^{\frac{h}{2}} \left| \Delta_t^{2m+1} f(x_i + \frac{h}{2} - mt) \right| dt \right. \\
 &\quad \left. + \int_0^{\frac{h}{2}} \left| \Delta_t^{2m+1} f(x_i + \frac{h}{2} - (m+1)t) \right| dt \right).
 \end{aligned}$$

Also as in 1), from (1.1) we have the next four inequalities are true for $t \in [0, \frac{h}{2}]$:

$$(3.8) \quad |\Delta_t^{2m+1} f(x_i - mt)| \leq \omega_{2m+1} \left(f, x_i + mt - m\frac{h}{2}; \frac{h}{2} \right);$$

$$(3.9) \quad |\Delta_t^{2m+1} f(x_i - (m+1)t)| \leq \omega_{2m+1} \left(f, x_i - mt + m\frac{h}{2}; \frac{h}{2} \right);$$

$$(3.10) \quad \left| \Delta_t^{2m+1} f \left(x_i + \frac{h}{2} - mt \right) \right| \leq \omega_{2m+1} \left(f, x_i + \frac{h}{2} + mt - m\frac{h}{2}; \frac{h}{2} \right);$$

$$(3.11) \quad \left| \Delta_t^{2m+1} f \left(x_i + \frac{h}{2} - (m+1)t \right) \right| \leq \omega_{2m+1} \left(f, x_i + \frac{h}{2} - mt + m\frac{h}{2}; \frac{h}{2} \right).$$

Applying the estimations of (3.8)-(3.11) in (3.7), we obtain

$$(3.12) \quad R_n^2(f) \leq \sum_{l=0}^{n-1} \frac{1}{3 \binom{2m+1}{m}} \int_0^{\frac{h}{2}} \omega_{2m+1} \left(f, x_i + mt; \frac{h}{2} \right) dt$$

$$\begin{aligned}
& + \omega_{2m+1} \left(f, x_i - mt; \frac{h}{2} \right) dt \\
& + \sum_{i=0}^{n-1} \frac{2}{3 \binom{2m+1}{m}} \int_0^{\frac{h}{2}} \omega_{2m+1} \left(f, x_i + \frac{h}{2} + mt; \frac{h}{2} \right) + \omega_{2m+1} \left(f, x_i + \frac{h}{2} - mt; \frac{h}{2} \right) dt \\
& = \sum_{l=0}^{n-1} \frac{1}{3m \binom{2m+1}{m}} \left(\int_{x_i - m \frac{h}{2}}^{x_i + m \frac{h}{2}} \omega_{2m+1} \left(f, x; \frac{h}{2} \right) dx \right. \\
& \quad \left. + 2 \int_{x_i + \frac{h}{2} - m \frac{h}{2}}^{x_i + \frac{h}{2} + m \frac{h}{2}} \omega_{2m+1} \left(f, x; \frac{h}{2} \right) dx \right) \\
& = \sum_{l=0}^{n-1} \frac{1}{\binom{2m+1}{m}} \int_{x_i}^{x_{i+1}} \omega_{2m+1} \left(f, x; \frac{h}{2} \right) dx = \frac{1}{\binom{2m+1}{m}} \tau_{2m+1} \left(f; \frac{h}{2} \right)_{L_1[0,1]}.
\end{aligned}$$

The inequalities (3.6) and (3.12) complete the proof of Lemma 3.1. $\blacksquare$

Lemma 3.2. *Let $k, n \in \mathbb{N}$ and $\alpha \geq \frac{1}{2}$. There is a 1-periodic function $f \in M[0, 1]$ such that*

$$R_n^2(f) = \frac{1}{\binom{k}{\lfloor \frac{k}{2} \rfloor}} \tau_k \left(f, \frac{\alpha}{n} \right)_{L_1[0,1]}.$$

Proof. There are many trivial bounded and measurable 1-periodic functions which prove this lemma but we construct such a function which we use in the next lemma. We use the notations $\mathbb{B}$ and B_m from Lemma 2.1. We set $q_k \stackrel{\text{def}}{=} \frac{2^{k-1} - \binom{k}{\lfloor \frac{k}{2} \rfloor}}{\binom{k}{\lfloor \frac{k}{2} \rfloor}} < 1$, $z_k \stackrel{\text{def}}{=} \left\lceil \frac{k}{4} \right\rceil + 1$ and $\diamond_{k,l} \stackrel{\text{def}}{=} \cup_{i=(l-1)z_k+1}^{lz_k} B_i$, $l \geq 1$. Let $n = 1$ (for $n > 1$ the idea is the same if we assume that $[x_i, x_{i+1}] = [0, 1]$). We define the function

$$f(x) \stackrel{\text{def}}{=} \begin{cases} (q_k)^{l-1} & \text{if } x \in \diamond_{k,l}; \\ 0 & \text{if } x \notin \mathbb{B}. \end{cases}$$

Obviously, $f \in M[0, 1]$. For the finite difference $\Delta_v^k f(x)$ with $v \leq \frac{1}{2n}$ and $x \in [0, 1]$ we have 3 cases:

- I) The knots of the finite difference are binary rational;
- II) The knots of the finite difference are not binary rational;
- III) The knots of the finite difference accept one are not binary rational.

In I) the absolute value of the finite difference is $< \binom{k}{[\frac{k}{2}]}$. In II) we have that the value of this difference is 0. In III) the absolute value of the finite difference is $\leq \binom{k}{[\frac{k}{2}]}$ ("=" iff $f(x) = 1$). Then $\omega_k(f, x; \frac{\alpha}{n}) = \binom{k}{[\frac{k}{2}]}$ and $\tau_k\left(f, \frac{\alpha}{n}\right)_{L_1[0,1]} = \binom{k}{[\frac{k}{2}]}$. On the other hand, applying (1.5) for this function, we have $R_n^2(f) = 1$.

Then Lemma 3.2 is proved. $\blacksquare$

Lemma 3.3. *Let $k, n \in \mathbb{N}$ and $\alpha \geq \frac{1}{2}$. There is a sequence of 1-periodic functions $\{f_m\}_{m=1}^\infty$, $f_m \in C^\infty[0, 1]$ such that*

$$\lim_{m \rightarrow \infty} \frac{R_n^2(f_m)}{\tau_k\left(f_m, \frac{\alpha}{n}\right)_{L_1[0,1]}} = \frac{1}{\binom{k}{[\frac{k}{2}]}}.$$

Proof. Let f be the function of Lemma 3.2 and for $m = 1, \dots, \infty$, $g_m(x)$ be the functions of Lemma 2.2. We set

$$f_m(x) \stackrel{\text{def}}{=} \sum_{s=0}^{2^{mz_k}} f\left(\frac{s}{2^{mz_k}}\right) g_{2^{2mz_k+1}}\left(x - \frac{s}{2^{mz_k}}\right).$$

For this functional sequence we have $f_m \in C^\infty[0, 1]$, $f_m(x) = f(x)$ for every $x \in \cup_{i=1}^{mz_k} B_i$ and $\lim_{m \rightarrow \infty} f_m(x) = f(x)$ for every $x \in [0, 1]$.

Also for the error of the quadrature formulae and for the k -th averaged modulus of f_m we have respectively,

$$\lim_{m \rightarrow \infty} R_n^2(f_m) = 1 \quad \text{and} \quad \lim_{m \rightarrow \infty} \tau_k\left(f_m, \frac{\alpha}{n}\right)_{L_1[0,1]} = \binom{k}{[\frac{k}{2}]}$$

Lemma 3.3 is proved. $\blacksquare$

Summarizing the statements from Lemma 3.1, Lemma 3.2 and Lemma 3.3, we obtain the following theorem.

Theorem 3.1. *Let $k \in \mathbb{N}$ and $\alpha \geq \frac{1}{2}$. For 1-periodic functions*

$$c(k, \alpha) = \tilde{c}(k, \alpha) = \frac{1}{\binom{k}{[\frac{k}{2}]}}.$$

Remark. Obviously the methods in Sections 3 and 4 are applicable for other n -composite quadrature formulae Q_n^s of Newton-Cotes type, constructed with respect to s knots. Let $k, n, s \in \mathbb{N}$ and $\alpha \geq \frac{1}{s}$. For 1-periodic functions we can prove the following estimation:

$$R_n^s(f) \leq \frac{1}{\binom{k}{[\frac{k}{2}]}} \tau_k \left(f, \frac{\alpha}{n} \right)_{L_1[0,1]},$$

where the constant is exact in $M[0, 1]$ and $C^\infty[0, 1]$.

4. Estimations of the error of the quadrature formulae for 1-periodic functions and a step $< \frac{h}{2}$

In this section we consider 1-periodic functions and a step $\alpha < \frac{1}{2}$. There are some specific problems. Here we use not only finite differencies centered in the knots of the quadrature formulae. We demonstrate this for $k = 2$ and $\alpha = \frac{1}{4}$. We first prove the upper estimation.

Lemma 4.1. *Let $f \in M[0, 1]$ be 1-periodic and $n \in \mathbb{N}$. Then,*

$$R_n^2(f) \leq \frac{2}{3} \tau_2 \left(f, \frac{1}{4n} \right)_{L_1[0,1]}.$$

Proof. Using again (1.5) and (1.4), we have

$$\begin{aligned} (4.1) \quad R_n^2(f) &= \left| \sum_{i=0}^{n-1} \frac{1}{3} \int_0^{\frac{h}{4}} \Delta_t^2 f(x_i - t) dt + \sum_{i=0}^{n-1} \frac{2}{3} \int_0^{\frac{h}{4}} \Delta_t^2 f \left(x_i + \frac{h}{2} - t \right) dt \right. \\ &\quad \left. - \sum_{i=0}^{n-1} \frac{1}{3} \int_0^{\frac{h}{2}} \Delta_{\frac{h}{4}}^2 f \left(x_i - \frac{h}{4} + t \right) dt \right| \\ &\leq \sum_{i=0}^{n-1} \frac{1}{3} \int_0^{\frac{h}{4}} |\Delta_t^2 f(x_i - t)| dt + \sum_{i=0}^{n-1} \frac{2}{3} \int_0^{\frac{h}{4}} \left| \Delta_t^2 f \left(x_i + \frac{h}{2} - t \right) \right| dt \\ &\quad + \sum_{i=0}^{n-1} \frac{1}{3} \int_0^{\frac{h}{2}} \left| \Delta_{\frac{h}{4}}^2 f \left(x_i - \frac{h}{4} + t \right) \right| dt. \end{aligned}$$

Again as in the previous sections, from (1.1) we have that the next three inequalities are true:

$$(4.2) \quad \int_0^{\frac{h}{4}} |\Delta_t^2 f(x_i - t)| dt \leq \frac{1}{2} \int_{x_i - \frac{h}{4}}^{x_i + \frac{h}{4}} \omega_2 \left(f, x; \frac{h}{4} \right) dx;$$

$$(4.3) \quad \int_0^{\frac{h}{4}} \left| \Delta_t^2 f \left(x_i + \frac{h}{2} - t \right) \right| dt \leq \frac{1}{2} \int_{x_i + \frac{h}{4}}^{x_i + 3\frac{h}{4}} \omega_2 \left(f, x; \frac{h}{4} \right) dx;$$

$$(4.4) \quad \int_0^{\frac{h}{2}} \left| \Delta_{\frac{h}{4}}^2 f \left(x_i - \frac{h}{4} + t \right) \right| dt \leq \int_{x_i - \frac{h}{4}}^{x_i + \frac{h}{4}} \omega_2 \left(f, x; \frac{h}{4} \right) dx.$$

Applying the estimations of (4.2)-(4.4) in (4.1), we obtain

$$R_n^2(f) \leq \sum_{i=0}^{n-1} \frac{2}{3} \int_{x_i}^{x_{i+1}} \omega_2 \left(f, x; \frac{h}{4} \right) dx = \frac{2}{3} \tau_2 \left(f, \frac{1}{4n} \right)_{L_1[0,1]}.$$

Lemma 4.1 is proved. ■

Lemma 4.2. *Let $n \in \mathbb{N}$. There is a 1-periodic function $f \in M[0, 1]$ such that*

$$R_n^2(f) = \frac{2}{3} \tau_2 \left(f, \frac{1}{4n} \right)_{L_1[0,1]}.$$

Proof. Let $n = 1$ (for $n > 1$ the idea is the same if we assume that $[x_i, x_{i+1}] = [0, 1]$). We define the function

$$f(x) \stackrel{\text{def}}{=} \begin{cases} 1 & \text{if } x = 0 \text{ or } 1; \\ 3 & \text{if } x = \frac{1}{2}; \\ 2 & \text{if } x \in [\frac{1}{4}, \frac{1}{2}) \cup (\frac{1}{2}, \frac{3}{4}]; \\ 0 & \text{if } x \in (0, \frac{1}{4}) \cup (\frac{3}{4}, 1). \end{cases}$$

Obviously, $f \in M[0, 1]$, $R_n^2(f) = \frac{4}{3}$ and $\tau_2 \left(f, \frac{1}{4n} \right)_{L_1[0,1]} = 2$.

Lemma 4.2 is proved. ■

Lemma 4.3. *There is a sequence of 1-periodic functions $\{f_m\}_{m=1}^\infty$, $f_m \in C^\infty[0, 1]$ such that*

$$\lim_{m \rightarrow \infty} \frac{R_n^2(f_m)}{\tau_2 \left(f_m, \frac{1}{4n} \right)_{L_1[0,1]}} = \frac{2}{3}.$$

Proof. Let f be the function of Lemma 4.2 and for $m = 1, \dots, \infty$, $g_m(x)$ be the functions of Lemma 2.2. We set

$$f_m(x) \stackrel{\text{def}}{=} \begin{cases} 2 + g_m(x - \frac{1}{2}) & \text{if } x \in [\frac{1}{4}, \frac{3}{4}]; \\ 2g_m(x - \frac{1}{4}) + g_m(x) & \text{if } x \in [0, \frac{1}{4}]; \\ 2g_m(x - \frac{3}{4}) + g_m(1 - x) & \text{if } x \in (\frac{3}{4}, 1]. \end{cases}$$

For this functional sequence we have $f_m \in C^\infty[0, 1]$, $f_m(x) = f(x)$, if $x = 0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}$ or 1 and $\lim_{m \rightarrow \infty} f_m(x) = f(x)$ for every $x \in [0, 1]$.

Also for the error of the quadrature formulae and for the 2-th averaged modulus of f_m we have respectively,

$$\lim_{m \rightarrow \infty} R_n^2(f_m) = \frac{4}{3} \quad \text{and} \quad \lim_{m \rightarrow \infty} \tau_2 \left(f_m, \frac{1}{4n} \right)_{L_1[0,1]} = 2.$$

Lemma 4.3 is proved. ■

Summarizing again the statements from Lemma 4.1, Lemma 4.2 and Lemma 4.3, we obtain the following theorem.

Theorem 4.1. *For 1-periodic functions,*

$$c \left(2, \frac{1}{4} \right) = \tilde{c} \left(2, \frac{1}{4} \right) = \frac{2}{3}.$$

References

- [1] K. Ivanov, New estimates of errors of quadrature formulae, formulae of numerical differentiation and interpolation, *J. Analysis Mathematica, Hung. Acad. of Sci. Budapest*, **6** (1980), 281-303.
- [2] Bl. Sendov, V. A. Popov, *The Averaged Moduli of Smoothness*, John Willey & Sons, N. York (1988).

*Department of Mathematics
Higher Transport School
jk "Slatina", 1754 - Sofia, BULGARIA*

Received: 10.02.2001