

A New Modulus of Smoothness for Trigonometric Polynomial Approximation

BORISLAV R. DRAGANOV *

The rate of convergence of best trigonometric approximation in L_p and C -norm is characterized by a new modulus of smoothness. This modulus equals 0 for all trigonometric polynomials up to a given degree.

1. Introduction and Main Results

The approximation by trigonometric polynomials is well investigated. The rate of convergence in uniform and integral norm was described by the classical modulus of smoothness due to D. Jackson, S. N. Bernstein, A. Zygmund and S. B. Stechkin (see for example [4]). This result was extended by Shapiro [5] and Ditzian [2] to any Banach space of 2π -periodic functions for which translation is continuous isometry.

Let B be a homogeneous Banach space of 2π -periodic real-valued functions. We denote by T_n the set of all trigonometric polynomials of degree at most n and put

$$E_n^T(f)_B := \inf_{\tau \in T_n \cap B} \|f - \tau\|_B$$

for the best trigonometric approximation. It was shown by the authors mentioned above that for $f \in B$ we have

$$\begin{aligned} E_n^T(f)_B &\leq C_r \omega_r(f; n^{-1})_B, \\ \omega_r(f; t)_B &\leq C_r t^r \sum_{0 \leq k \leq 1/t} (k+1)^{r-1} E_k^T(f)_B, \quad 0 < t \leq t_0, \end{aligned}$$

where $\omega_r(f; t)_B$ is the classical modulus of smoothness defined as follows

$$\begin{aligned} \omega_r(f; t)_B &= \sup_{0 < h \leq t} \|\Delta_h^r f\|_B, \\ \Delta_h^r f(x) &= \sum_{k=0}^r (-1)^k \binom{r}{k} f(x + (r/2 - k)h). \end{aligned}$$

*The author is supported by the University of Sofia.

Let B denote $L_p^*[-\pi, \pi]$, $1 \leq p < \infty$, or $C^*[-\pi, \pi]$ where

$$\begin{aligned} L_p^*[-\pi, \pi] &:= \{f : \mathbb{R} \rightarrow \mathbb{R} : f(x + 2\pi) = f(x) \text{ a.e., } f|_{[-\pi, \pi]} \in L_p[-\pi, \pi]\} \\ C^*[-\pi, \pi] &:= \{f \in C(\mathbb{R}) : f(x + 2\pi) = f(x)\}. \end{aligned}$$

It is interesting to construct a modulus-like function $\omega_r^T(f; t)_p$ associated with best trigonometric approximation such that for $f \in B$ we have

$$\omega_r^T(f; t)_p \equiv 0 \iff f \in T_{r-1} \tag{1}$$

(here and further $f \in T_{r-1}$ in L_p -spaces means that f coincides a.e. with a trigonometric polynomial of degree at most $r - 1$; we use such a stipulation in similar cases as well) and $\omega_r^T(f; t)_p$ characterizes the best trigonometric approximation like the classical modulus does. Thus, this new modulus of smoothness describes more precisely (in the sense of (1)) the rate of convergence of best trigonometric approximation than the classical one. The definition of this new modulus, as we should expect, is more complicated. We shall show that the modulus of smoothness defined by

$$\omega_r^T(f; t)_p = \sup_{0 < h \leq t} \|\Delta_h^{2r-1} \mathcal{F}_{r-1} f\|_p, \quad r = 1, 2, \dots, \tag{2}$$

where

$$\mathcal{F}_{r-1}(f, x) = f(x) + \int_0^x \mathcal{K}_{r-1}(t) f(x-t) dt \tag{3}$$

and

$$\mathcal{K}_{r-1}(t) = \sum_{j=1}^{r-1} \frac{a_j^{(r-1)}}{(2j-1)!} t^{2j-1}, \quad a_j^{(r-1)} = \sum_{1 \leq l_1 < \dots < l_j \leq r-1} (l_1 \cdots l_j)^2,$$

satisfies (1) and the following theorem holds.

Theorem 1. *Let $f \in B$ where $B = L_p^*[-\pi, \pi]$, $1 \leq p < \infty$, or $B = C^*[-\pi, \pi]$. Then*

$$E_n^T(f)_p \leq C_r \omega_r^T(f; n^{-1})_p, \quad n \geq r - 1, \tag{4}$$

and

$$\omega_r^T(f; t)_p \leq C_r t^{2r-1} \sum_{r-1 \leq k \leq 1/t} (k+1)^{2r-2} E_k^T(f)_p, \quad 0 < t \leq \frac{1}{r}. \tag{5}$$

In particular, (4) gives the trigonometric analogue of Whitney theorem [6], as it was observed by Bl. Sendov.

Theorem 2. *Let $f \in B$ where $B = L_p^*[-\pi, \pi]$, $1 \leq p < \infty$, or $B = C^*[-\pi, \pi]$. Then*

$$E_{n-1}^T(f)_p \leq C_n \omega_n^T(f; n^{-1})_p.$$

Let us observe that although $\mathcal{F}_{r-1}f$ is not generally a 2π -periodic function, $\Delta_h^{2r-1}\mathcal{F}_{r-1}f$ is. Unlike the various moduli which describe the best algebraic approximation $\omega_r^T(f; t)_p$ is based on finite differences only of an odd order, and consequently, $\omega_r^T(f; t)_p$ is connected with the $(2r - 1)$ -th finite difference not the r -th one. This is due to the dimensions of the spaces T_{r-1} and Π_{r-1} , respectively. To state our next main result we define the K -functional

$$K_r^T(f; t)_p = \inf_{g \in B^{2r-1}} \{ \|f - g\|_p + t^{2r-1} \|\tilde{D}_r g\|_p \} \tag{6}$$

where we have put $B^s = \{g \in B : g, g', \dots, g^{(s-1)} \in AC^*[-\pi, \pi], g^{(s)} \in B\}$, $AC^*[-\pi, \pi]$ being the set of all 2π -periodic absolutely continuous functions, $\tilde{D}_r g = D_{r-1} \dots D_1 g'$ and $D_j g = g'' + j^2 g$. We have $\tilde{D}_r g = 0$ if and only if $g \in T_{r-1}$.

We write $\varphi(f; t) \sim \psi(f; t)$ if and only if there exists a constant $C > 0$ independent of f and t such that $C^{-1}\varphi(f; t) \leq \psi(f; t) \leq C\varphi(f; t)$. The following result holds.

Theorem 3. *For $f \in B$ where $B = L_p^*[-\pi, \pi]$, $1 \leq p < \infty$, or $B = C^*[-\pi, \pi]$, we have*

$$K_r^T(f; t)_p \sim \omega_r^T(f; t)_p$$

where $\omega_r^T(f; t)_p$ and $K_r^T(f; t)_p$ are defined in (2) and (6), respectively.

A detailed version of this research paper is accepted for publication in *East Journal on Approximations*.

2. A New Periodic Modulus of Smoothness

Let $[a, b]$ be an arbitrary finite subinterval of the real line such that $0 \in [a, b]$. We write $X = X[a, b]$ for any of the functional spaces $L_p[a, b]$, $1 \leq p < \infty$, or $C[a, b]$, and $X^r = X^r[a, b]$ for the Sobolev spaces $W_p^r[a, b]$, $1 \leq p < \infty$, or $C^r[a, b]$. We also write B for $L_p^*[-\pi, \pi]$, $1 \leq p < \infty$, or $C^*[-\pi, \pi]$. Finally, B^r will stand for $W_p^{*r}[-\pi, \pi]$, $1 \leq p < \infty$, or $C^{*r}[-\pi, \pi]$, where

$$\begin{aligned} W_p^{*r}[-\pi, \pi] &= \{f \in L_p^*[-\pi, \pi] : f, \dots, f^{(r-1)} \in AC^*[-\pi, \pi], f^{(r)} \in L_p^*[-\pi, \pi]\}, \\ C^{*r}[-\pi, \pi] &= \{f \in C^*[-\pi, \pi] : f^{(k)} \in C^*[-\pi, \pi], k = 1, \dots, r\}. \end{aligned}$$

The proof of the main result is based on several auxiliary propositions. First we consider the inverse of the bounded operator $\mathcal{F}_r : X \rightarrow X$.

Proposition 1. *We have*

$$\mathcal{F}_r^{-1}(g, x) = g(x) - \int_0^x \mathcal{L}_r(t)g(x - t) dt,$$

where

$$\mathcal{L}_r(t) = 2 \sum_{j=1}^r \frac{j^{2r}}{\omega_r'(j)} \sin jt \tag{7}$$

and $\omega_r(x) = \prod_{k=1}^r (x^2 - k^2)$. Hence $\mathcal{F}_r^{-1} : X \rightarrow X$ is bounded.

Formula (7) was pointed to the author by K. G. Ivanov. Let Π_s denote the set of all algebraic polynomials of degree at most s .

Proposition 2. *For \mathcal{F}_r , defined in (3), we have*

- (a) $(\mathcal{F}_r g)^{(2r+1)} = \tilde{D}_{r+1}g, g \in X^{2r+1};$
- (b) $\mathcal{F}_r \tau \in \Pi_{2r}, \tau \in T_r;$
- (c) $\mathcal{F}_r^{-1}P \in T_r, P \in \Pi_{2r}.$

After these preliminaries we can formulate the main result of the section. First we introduce the bounded linear operator $\tilde{\mathcal{F}}_{r-1} : B \rightarrow B, r \geq 2,$

$$\tilde{\mathcal{F}}_{r-1}(f, x) = \mathcal{F}_{r-1}(f, x) + P_{2r-2}(f, x),$$

where \mathcal{F}_{r-1} is defined in (3) and $P_{2r-2}(f, x) = -\sum_{k=1}^{2r-2} \alpha_k(x + \pi)^k/k!$ is the unique algebraic polynomial of degree $2r - 2$ such that we may have $(\tilde{\mathcal{F}}_{r-1}f)^{(s)}(-\pi) = (\tilde{\mathcal{F}}_{r-1}f)^{(s)}(\pi), s = 0, 1, \dots, 2r - 3,$ for any $f \in B^{2r-3}$ and hence $\tilde{\mathcal{F}}_{r-1} : B^s \rightarrow B^s, s \in \mathbb{N}.$

Let I be the identity. For the sake of consistency, we set $\mathcal{F}_0 = I (\mathcal{K}_0 = 0)$ and $P_0 = 0,$ and then $\tilde{\mathcal{F}}_0 = I.$

We introduce the following modulus of smoothness for a function $f \in B$ and $t > 0:$

$$\omega_r^T(f; t)_p = \omega_{2r-1}(\tilde{\mathcal{F}}_{r-1}f; t)_p, \quad r = 1, 2, \dots,$$

where $\omega_{2r-1}(F; t)_p$ is the classical periodic modulus of smoothness of order $2r - 1.$ Let us note that $\Delta_h^{2r-1}\mathcal{F}_{r-1}f \in B$ for any $f \in B$ and

$$\omega_r^T(f; t)_p = \sup_{0 < h \leq t} \|\Delta_h^{2r-1}\mathcal{F}_{r-1}f\|_p, \quad r = 1, 2, \dots$$

In the definition of $\omega_r^T(f; t)_p,$ the quantity $\Delta_h^{2r-1}\tilde{\mathcal{F}}_{r-1}f(x)$ depends only on the values of f in a neighbourhood of x whose diameter diminishes with $h.$ The point 0 in the integration limits of the integral operator used in the definition of $\tilde{\mathcal{F}}_{r-1}$ has been chosen only for convenience – any other value can be fixed and the definition of $\omega_r^T(f; t)_p$ is invariant of this choice.

Now (1) follows immediately from Proposition 2 (b)-(c) and the definition of $\omega_r^T(f; t)_p.$

The following equivalence result holds.

Theorem 4. For $f \in B$ where $B = L_p^*[-\pi, \pi]$, $1 \leq p < \infty$, or $B = C^*[-\pi, \pi]$ and any $l \in \mathbb{N}$, we have

$$\inf_{g \in B^{2r+l-1}} \{ \|f - g\|_p + t^{2r+l-1} \|(d/dx)^l \tilde{D}_r g\|_p \} \sim \omega_{2r+l-1}(\tilde{\mathcal{F}}_{r-1} f; t)_p.$$

The proof of this theorem is based on the properties of the operator $\tilde{\mathcal{F}}_{r-1}$ (see the propositions above) and those of another bounded linear operator $\mathcal{E}_{r-1} : B \rightarrow B$ defined by $\mathcal{E}_{r-1}(F, x) = \mathcal{F}_{r-1}^{-1}(F, x) + Q_{2r}(F, x)$ where $Q_{2r}(F, x)$ is an algebraic polynomial of the form $Q_{2r}(F, x) = \sum_{k=1}^{2r} \beta_k(x + \pi)^k/k!$, depending on F , such that $\mathcal{E}_{r-1}F \in B^s$ for any $F \in B^s$ for $s = 0, 1, \dots, 2r - 1$.

Now Theorem 3 follows from Theorem 4 for $l = 0$ and the definition of $\omega_r^T(f; t)_p$.

It is easy to verify, using the definition of $\omega_r^T(f; t)_p$ and some properties of the operator $\tilde{\mathcal{F}}_{r-1}$, that $\omega_r^T(f; t)_p$ possesses the properties:

1. $\omega_r^T(f + g; t)_p \leq \omega_r^T(f; t)_p + \omega_r^T(g; t)_p$ for $f, g \in B$.
2. $\omega_r^T(cf; t)_p = |c|\omega_r^T(f; t)_p$, c is a constant.
3. $\omega_r^T(f; t)_p \leq \omega_r^T(f; t')_p$, $t \leq t'$.
4. $\omega_r^T(f; t)_p \rightarrow 0$ as $t \rightarrow 0$.
5. $\omega_r^T(f; t)_p \leq (4 + (r - 1)^2 t^2) \omega_{r-1}^T(f; t)_p$, $r \geq 2$.
6. $\omega_1^T(f; t)_p \leq 2\|f\|_p$ and $\omega_1^T(f; t)_p \leq t\|f'\|_p$, $f \in B^1$ ($\omega_1^T(f; t)_p$ coincides with the ordinary modulus of continuity).
7. $\omega_r^T(f; \lambda t)_p \leq (\lambda + 1)^{2r-1} \omega_r^T(f; t)_p$.
8. $\omega_r^T(f; t)_p \leq t^2 \omega_{r-1}^T(D_{r-1} f; t)_p$, $f \in B^2$; $r \geq 2$.
9. Marchaud inequality

$$\omega_r^T(f; t)_p \leq C_r t^{2r-1} \left(\int_t^c \frac{\omega_{r+1}^T(f; u)_p}{u^{2r}} du + \|f\|_p \right), \quad 0 < t \leq c$$

where c is any fixed positive constant.

10. $\omega_r^T(f; t)_p = o(t^{2r-1}) \implies f \in T_{r-1}$ and $f \in T_{r-1} \implies \omega_r^T(f; t)_p \equiv 0$, $1 \leq p \leq \infty$.
11. $\omega_r^T(f; t)_p = \mathcal{O}(t^{2r-1}) \iff f \in W_p^{*2r-1}[-\pi, \pi]$, $1 < p \leq \infty$.
12. $\omega_r^T(f; t)_1 = \mathcal{O}(t^{2r-1}) \iff f^{(2r-3)} \in AC^*[-\pi, \pi]$, $f^{(2r-2)} \in BV[-\pi, \pi]$.

3. Best Trigonometric Approximation

Now we can sketch the proof of our main result concerning the rate of best trigonometric approximation in $L_p^*[-\pi, \pi]$, $1 \leq p < \infty$, and $C^*[-\pi, \pi]$.

Sketch of the proof of Theorem 1. To prove (4) we follow some methods demonstrated in [1] and [3] and use the combination of some modified Riesz operators

$$L_{r-1,n} = I - \prod_{j=0}^{r-1} (I - R_{j,n}) = \sum_{i=0}^{r-1} (-1)^i \sum_{0 \leq j_0 < \dots < j_i \leq r-1} R_{j_0,n} \cdots R_{j_i,n}$$

where I is the identity and

$$R_{j,n}(f, x) = \sum_{k=0}^{n-1} \left(1 - \frac{k^2 - j^2}{n^2 - j^2}\right) A_k(x), \quad n > j, \quad j = 1, 2, \dots,$$

$A_k(x) = A_k(f, x)$ being the k -th term in the Fourier expansion of f .

The inequality (5) can be proven similarly to its classical analogue.

Acknowledgements. I am grateful to Prof. K. Ivanov for his directions. I am also thankful to the referees for their corrections.

References

- [1] Z. DITZIAN, A K -functional and the rate of convergence of some linear polynomial operators, *Proc. Amer. Math. Soc.* **124** (1996), 1773–1781.
- [2] Z. DITZIAN, Some remarks on approximation theorems on various Banach spaces, *J. Math. Anal. Appl.* **77** (1980), 567–576.
- [3] Z. DITZIAN AND K. G. IVANOV, Strong converse inequalities, *J. D'Analyse Math.* **61** (1993), 61–111.
- [4] G. G. LORENTZ, “Approximation of Functions”, Chelsea Publishing Company, New York, 1986.
- [5] H. S. SHAPIRO, “Topics in Approximation Theory”, *Lecture Notes in Math.* **187**, Springer-Verlag, Berlin, 1970.
- [6] H. WHITNEY, On functions with bounded n -th differences, *J. Math Pures Appl.* **36**, 9 (1957), 67–95.

BORISLAV R. DRAGANOV
 Faculty of Mathematics and Informatics
 University of Sofia
 Blvd. J. Bourchier 5
 1164 Sofia
 BULGARIA
E-mail: bdraganov@fmi.uni-sofia.bg