

Provided for non-commercial research and educational use.
Not for reproduction, distribution or commercial use.

Mathematica Balkanica

Mathematical Society of South-Eastern Europe
A quarterly published by
the Bulgarian Academy of Sciences – National Committee for Mathematics

The attached copy is furnished for non-commercial research and education use only. Authors are permitted to post this version of the article to their personal websites or institutional repositories and to share with other researchers in the form of electronic reprints.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to third party websites are prohibited.

For further information on Mathematica Balkanica visit the website of the journal
<http://www.mathbalkanica.info>

or contact:

Mathematica Balkanica - Editorial Office;
Acad. G. Bonchev str., Bl. 25A, 1113 Sofia, Bulgaria
Phone: +359-2-979-6311, Fax: +359-2-870-7273,
E-mail: balmat@bas.bg

A Note on Lattice Horn Sentences with Three Variables

Gábor Czédli

Presented by P. Kenderov

Lattice Horn sentences with three variables are shown to be trivial within the theory of modular lattices.

In the theory of modular lattices, all the interesting Horn sentences that have been investigated have at least four variables (cf., e. g., G. Grätzer, H. Lakser and B. Jónsson [4] and [2, 3]). The aim of the present note is to explain this phenomenon by the following

Theorem. *Let χ be a lattice Horn sentence on three variables. Then either χ is a consequence of the modular law or χ together with modularity imply distributivity.*

By a lattice Horn sentence χ with three variables we mean a universally quantified formula

$$p_1 \leq q_1 \ \& \ \dots \ \& \ p_k \leq q_k \Rightarrow p \leq q$$

where $p_1, \dots, p_k, q_1, \dots, q_k, p, q$ are lattice terms on the set $\{x, y, z\}$ of variables and $k \geq 0$. (In case $k=0$ the premise is empty and χ is a lattice identity.) In what follows the reader is assumed to be familiar with some basic facts belonging to universal algebra and lattice theory and, first of all, with the diagram of $M = F_M(x, y, z)$, the free modular lattice generated by $\{x, y, z\}$ (cf. any book on lattice theory, e. g., G. Grätzer [3, p. 39]).

Proof. Without loss of generality we may assume that $p, p_1, \dots, p_k, q, q_1, \dots, q_k$ are elements of M . If our statement is true for $\chi_i, i=1, 2$, and χ is equivalent (modulo lattice theory) to the conjunction of χ_1 and χ_2 then the statement is also true for χ . On the other hand, in any lattice and for arbitrary lattice terms $r, r_1, r_2, r_1+r_2 \leq r$ holds iff $r_1 \leq r$ and $r_2 \leq r$ hold, and dually. This allows us to make the following assumption: $p, p_1, \dots, p_k \in M$ are join-irreducible elements and $q, q_1, \dots, q_k \in M$ are meet-irreducible elements. We may also suppose that $p_1 \not\leq q_1, \dots, p_k \not\leq q_k, p \not\leq q$ in M and, in $M, p \leq p_i$ simultaneously with $q_i \leq q$ hold for no $i \in \{1, \dots, k\}$ as otherwise χ would automatically hold in all modular lattices or some $p_i \leq q_i$ could be omitted from the premise of χ . Finally, we assume that χ holds in $\mathbf{2}$, the two-element lattice, as otherwise χ would trivially imply distributivity. Therefore, by this assumption, χ holds in every distributive lattice as each distributive lattice is a subdirect power of $\mathbf{2}$.

Now let $u = xy + xz + yz$, $v = (x + y)(x + z)(y + z)$, $a = u + xv$, $b = u + yv$, $c = u + zv$. Then $M_3 = [u, v] = \{u, a, b, c, v\}$ is the only diamond (i. e., five element non-distributive sublattice) in M . Let Θ denote the congruence of M corresponding to the partition $\{[x(y + z), x + yz], [y(x + z), y + xz], [z(x + y), z + xy], [0, u], [v, 1]\}$. We claim that, for any congruence Ψ of M , M/Ψ is distributive iff $\Psi \not\subseteq \Theta$. Indeed, if $\Psi \subseteq \Theta$ then $M_3 \cong M/\Theta$ is a homomorphic image of M/Ψ , whence M/Ψ is not distributive. Conversely, if $\Psi \not\subseteq \Theta$ then, as the blocks of Ψ are intervals, there are $d, e \in M$ such that $(d, e) \in \Psi \setminus \Theta$ and e covers d . Further, $\{d, e\} \subseteq M_3$ can be assumed as the only other case is symmetric and/or dual to $d = a + x$, $e = v + x$, whence $(a, v) = (vd, ve) \in \Psi \setminus \Theta$. Since $\Psi \cap M_3^2 \neq \emptyset$ and M_3 is a simple lattice, Ψ includes Θ_3 , the congruence of M generated by M_3^2 . Hence M/Ψ is a homomorphic image of M/Θ_3 , which is a distributive (moreover, a free distributive) lattice. Thus M/Ψ is distributive.

Now let Θ_i denote the congruence of M generated by $(p_i, p_i q_i)$. Suppose $\Theta_i \not\subseteq \Theta$ holds for some $i \in \{1, \dots, k\}$. If L is an arbitrary modular lattice, $x_1, y_1, z_1 \in L$ and $p_j(x_1, y_1, z_1) \leq q_j(x_1, y_1, z_1)$ for $j = 1, \dots, k$ then Θ_i is included in the kernel of the surjective homomorphism $M \rightarrow [x_1, y_1, z_1]$, $x \mapsto x_1, y \mapsto y_1, z \mapsto z_1$. Hence $[x_1, y_1, z_1]$, as a homomorphic image of M/Θ_i , is distributive, whereupon $p(x_1, y_1, z_1) \leq q(x_1, y_1, z_1)$. This shows that χ holds in L .

From now on let us assume that $\Theta_i \subseteq \Theta$ holds for all $i \in \{1, \dots, k\}$. Then the premise of χ holds for $a, b, c \in M_3$. Really, considering the homomorphism $\tau: M \rightarrow M_3$, $x \mapsto a, y \mapsto b, z \mapsto c$, from $(p_i, p_i q_i) \in \Theta = \ker \tau$ we obtain $p_i(a, b, c) = p_i(x\tau, y\tau, z\tau) = p_i(x, y, z)\tau = p_i\tau = (p_i q_i)\tau = (p_i\tau)(q_i\tau) = p_i(x\tau, y\tau, z\tau)q_i(x\tau, y\tau, z\tau) = p_i(a, b, c)q_i(a, b, c)$ for $i \in \{1, \dots, k\}$. If $p(a, b, c) \not\leq q(a, b, c)$ then χ fails to hold in M_3 . Hence χ and modularity imply distributivity. Therefore we assume that $p(a, b, c) \leq q(a, b, c)$. In virtue of our former assumptions, a quick glance at M shows that, apart from symmetry and duality, $p = x$ and $q = y + z$. Since χ holds in $\{0, 1\}$, a distributive sublattice of M , but $1 = p(1, 0, 0) \not\leq q(1, 0, 0) = 0$, there is an $i \in \{1, \dots, k\}$ such that $p_i(1, 0, 0) \not\leq q_i(1, 0, 0)$, i. e., $p_i(1, 0, 0) = 1$ and $q_i(1, 0, 0) = 0$. Considering the homomorphism $\varphi: M \rightarrow \{0, 1\}$, $x \mapsto 1, y \mapsto 0, z \mapsto 0$, $\ker \varphi$ has only two blocks: $[x, 1]$ and $[0, y + z]$. Since $p_i\varphi = p_i(x, y, z)\varphi = p_i(x\varphi, y\varphi, z\varphi) = p_i(1, 0, 0) = 1 = 1\varphi$ and $q_i\varphi = 0 = 0\varphi$, we have $p_i \in [x, 1]$ and $q_i \in [0, y + z]$. Thus the contradiction $p \leq p_i$ and $q_i \leq q$ completes the proof.

References

1. G. Czédli. Horn sentences in submodule lattices. *Acta. Sci. Math. (Szeged)*, **51**, 1987, 17-33.
2. G. Czédli, G. Hutchinson. An irregular Horn sentence in submodule lattices. *Acta Sci. Math. (Szeged)*, **51**, 1987, 35-38.
3. G. Grätzer. *General Lattice Theory*. Akademie-Verlag, Berlin, 1978.
4. G. Grätzer, H. Lakser, B. Jónsson. The amalgamation property in equational classes of modular lattices, *Pacific J. Math.*, **45**, 1973, 507-524.

JATE Bolyai Institute
Aradi vértanúk tere 1
H-6720 Szeged
HUNGARY

Received 29. 06. 1989