

ON HANKEL TRANSFORM OF GENERALIZED MATHIEU SERIES

Živorad Tomovski

Abstract

By using integral representations for several Mathieu type series, a number of integral transforms of Hankel type are derived here for general families of Mathieu type series. These results generalize the corresponding ones on the Fourier transforms of Mathieu type series, obtained recently by Elezovic et al. [4], Tomovski [19] and Tomovski and Vu Kim Tuan [20].

Mathematics Subject Classification: Primary 33E20, 44A10; Secondary 33C10, 33C20, 44A20

Key Words and Phrases: integral representations, Mathieu series, Hankel transform, Fourier transform, Bessel function, Fox-Wright function, Sonine-Schafheitlin formula, Fox H -function

1. Introduction and preliminaries

The following familiar infinite series

$$S(r) = \sum_{n=1}^{\infty} \frac{2n}{(n^2 + r^2)^2} \quad (r \in \mathbb{R}^+) \quad (1.1)$$

is named after Emile Leonard Mathieu (1835-1890), who investigated it in his 1890 work [8] on elasticity of solid bodies. An integral representation of (1.1) is given by (see [5])

$$S(r) = \frac{1}{r} \int_0^{\infty} \frac{t \sin(rt)}{e^t - 1} dt. \quad (1.2)$$

Several interesting results dealing with integral representations and bounds for a slight generalization of the Mathieu series with a fractional power, defined as

$$S_\mu(r) = \sum_{n=1}^{\infty} \frac{2n}{(n^2 + r^2)^\mu} \quad (r \in \mathbb{R}^+; \mu > 1), \quad (1.3)$$

can be found in the works by Diananda [2], Tomovski and Trencovski [16], Cerone and Lenard [1] and Tomovski [18]. Motivated essentially by the works of Cerone and Lenard [1] (and Qi [13]), Srivastava and Tomovski [14] defined a family of generalized Mathieu series

$$S_\mu^{(\alpha, \beta)}(r; a) = S_\mu^{(\alpha, \beta)}(r; \{a_k\}_{k=1}^\infty) = \sum_{n=1}^{\infty} \frac{2a_n^\beta}{(a_n^\alpha + r^2)^\mu} \quad (r, \alpha, \beta, \mu \in \mathbb{R}^+), \quad (1.4)$$

where it is tacitly assumed that the positive sequence

$$a = \{a_n\}_{n=1}^\infty = \{a_1, a_2, a_3, \dots\} \quad \left(\lim_{n \rightarrow \infty} a_n = \infty \right)$$

is so chosen that the infinite series in definition (1.4) converges, that is, that the following auxiliary series

$$\sum_{n=1}^{\infty} \frac{1}{a_n^{\mu\alpha - \beta}}$$

is convergent. Comparing the definitions (1.1), (1.3) and (1.4), we see that $S_2(r) = S(r)$ and $S_\mu(r) = S_\mu^{(2,1)}(r, \{k\})$. Furthermore, the special cases $S_2^{(2,1)}(r; \{a_k\})$, $S_\mu^{(2,1)}(r; \{k^\gamma\})$ and $S_\mu^{(\alpha, \alpha/2)}(r; \{k\})$ have been investigated by Qi [13], Diananda [2], Tomovski [17] and Cerone-Lenard [1].

2. Some definitions and formulas

In this section we give some definitions and formulas, needed for computation of the Hankel transforms of $S(r)$, $S_{\mu+1}(r)$, $S_\mu^{(\alpha, \beta)}(r; \{k^{2/\alpha}\})$ and $S_\mu^{(\alpha, \beta)}(r; \{k^\gamma\})$.

2.1. In order to evaluate the Hankel transform of $S(r)$, we first set $\nu = \frac{1}{2}$ in the Sonine-Schafheitlin formula (see for example, [6, p.692]):

$$\int_0^\infty t^{-\lambda} J_\mu(at) J_\nu(bt) dt = \frac{a^\mu \Gamma((\mu + \nu - \lambda + 1)/2)}{2^\lambda b^{\mu - \lambda + 1} \Gamma(\mu + 1) \Gamma((\lambda - \mu + \nu + 1)/2)}$$

$$\times {}_2F_1\left(\frac{\mu + \nu - \lambda + 1}{2}, \frac{\mu - \nu - \lambda + 1}{2}; \mu + 1; \frac{a^2}{b^2}\right) \quad (2.1)$$

$$(\Re(\mu + \nu + 1) > \Re(\lambda) > -1; 0 < a < b)$$

with a corresponding expression for the case when $0 < b < a$, which is obtained from (2.1) by interchanging a and b , and also μ and ν . In view of the relationship

$$J_{1/2}(x) = \sqrt{\frac{2}{\pi x}} \sin x, \quad (2.2)$$

we find from the Sonine-Schafheitlin formula (2.1) that

$$\int_0^\infty t^{-\lambda-1/2} J_\mu(at) \sin(bt) dt = \sqrt{\frac{\pi b}{2}} \frac{a^\mu \Gamma\left[\frac{1}{2}(\mu - \lambda + 3/2)\right]}{2^\lambda b^{\mu-\lambda+1} \Gamma(\mu+1) \Gamma\left[\frac{1}{2}(\lambda - \mu + 3/2)\right]}$$

$$\times {}_2F_1\left(\frac{\mu - \lambda + 3/2}{2}, \frac{\mu - \lambda + 1/2}{2}; \mu + 1; \frac{a^2}{b^2}\right) \quad (2.3)$$

$$\left(\Re(\mu + \frac{3}{2}) > \Re(\lambda) > -1; 0 < a < b\right),$$

together with the corresponding integral derived from the analogue of (2.1) for the case when $0 < b < a$. Thus, by further applying integral formula (2.3) and its companion when $0 < b < a$, we obtain (with $\lambda = 0$, $\mu \rightarrow \nu$):

$$\int_0^\infty r^{-1/2} \sin(rt) J_\nu(rx) dr = \sqrt{\frac{\pi}{2}} \begin{cases} \Omega_\nu^{(1)}(x; t) : 0 < x < t \\ \Omega_\nu^{(2)}(x; t) : 0 < t < x \end{cases} \quad \left(\Re(\nu) > -\frac{1}{2}\right), \quad (2.4)$$

where

$$\Omega_\nu^{(1)}(x; t) = \frac{x^\nu}{t^{\nu+1/2}} \frac{\Gamma(\frac{\nu}{2} + \frac{3}{4})}{\Gamma(\nu+1) \Gamma(\frac{3}{4} - \frac{\nu}{2})} {}_2F_1\left(\frac{\nu}{2} + \frac{3}{4}, \frac{\nu}{2} + \frac{1}{4}; \nu + 1; \frac{x^2}{t^2}\right)$$

$$(0 < x < t)$$

$$\Omega_\nu^{(2)}(x; t) = \frac{t}{x^{3/2}} \frac{\Gamma(\frac{\nu}{2} + \frac{3}{4})}{\Gamma(\frac{3}{2}) \Gamma(\frac{\nu}{2} + \frac{1}{4})} {}_2F_1\left(\frac{\nu}{2} + \frac{3}{4}, \frac{3}{4} - \frac{\nu}{2}; \frac{3}{2}; \frac{t^2}{x^2}\right)$$

$$(0 < t < x) \quad (\Re(\nu) > -\frac{1}{2}).$$

2.2. If we apply the Sonine-Schafheitlin formula (2.1), first with $\lambda = \mu - 1$, $\mu \rightarrow \mu - 1/2$, $a = t$, $b = x$, ($0 < a < b$), and then with $\lambda = \mu - 1$, $\mu \rightarrow \nu$, $\nu \rightarrow \mu - \frac{1}{2}$, $a = x$, $b = t$ ($0 < b < a$), we get

$$\int_0^\infty r^{1-\mu} J_{\mu-1/2}(rt) J_\nu(rx) dr = \begin{cases} \Theta_\nu^{(1)}(\mu; x, t) : 0 < t < x \\ \Theta_\nu^{(2)}(\mu; x, t) : 0 < x < t \end{cases} \quad \left(\Re(\mu) > 0, \Re(\nu) > -\frac{3}{2} \right), \quad (2.5)$$

where

$$\Theta_\nu^{(1)}(\mu; x, t) = \frac{t^{\mu-\frac{1}{2}} \Gamma\left(\frac{\nu}{2} + \frac{3}{4}\right)}{2^{\mu-1} x^{\frac{3}{2}} \Gamma(\mu + 1/2) \Gamma\left(\frac{\nu}{2} + \frac{1}{4}\right)} {}_2F_1\left(\frac{3}{4} + \frac{\nu}{2}, \frac{3}{4} - \frac{\nu}{2}; \mu + \frac{1}{2}; \frac{t^2}{x^2}\right) \quad \left(0 < t < x, \Re(\mu) > 0, \Re(\nu) > -\frac{3}{2}\right), \quad (2.6)$$

$$\Theta_\nu^{(2)}(\mu; x, t) = \frac{x^\nu \Gamma\left(\frac{\nu}{2} + \frac{3}{4}\right)}{2^{\mu-1} t^{\nu-\mu+2} \Gamma(\nu + 1) \Gamma\left(\mu - \frac{\nu}{2} - \frac{1}{4}\right)} \times {}_2F_1\left(\frac{3}{4} + \frac{\nu}{2}, \frac{\nu}{2} - \mu + \frac{5}{4}; \nu + 1; \frac{x^2}{t^2}\right) \quad \left(0 < x < t, \Re(\mu) > 0, \Re(\nu) > -\frac{3}{2}\right). \quad (2.7)$$

2.3. For the evaluation of the Hankel transform of the general Mathieu series $S_\mu^{(\alpha, \beta)}(r; \{k^{2/\alpha}\})$, we need the definition of the Meijer G -function and the Fox H -function (for more details, see e.g. in [3, Vol.1], [12]).

DEFINITION 1. By a Fox's H -function we mean a generalized hypergeometric function, defined by means of the Mellin-Barnes-type contour integral

$$H_{p,q}^{m,n} \left[\sigma \left| \begin{matrix} (a_p, \alpha_p) \\ (b_q, \beta_q) \end{matrix} \right. \right] = \frac{1}{2\pi i} \int_{\mathcal{L}} \frac{\prod_{k=1}^m \Gamma(b_k - s\beta_k) \prod_{j=1}^n \Gamma(1 - a_j + s\alpha_j)}{\prod_{k=m+1}^q \Gamma(1 - b_k + s\beta_k) \prod_{j=n+1}^p \Gamma(a_j - s\alpha_j)} \sigma^s ds, \quad (2.8)$$

where \mathcal{L} is a suitable contour in \mathbb{C} , the orders (m, n, p, q) are integers, $0 \leq m \leq q$, $0 \leq n \leq p$ and the parameters $a_j \in \mathbb{R}$, $\alpha_j > 0$, $j = 1, 2, \dots, p$, $b_k \in \mathbb{R}$, $\beta_k > 0$, $k = 1, 2, \dots, q$ are such that $\alpha_j(b_k + l) \neq \beta_k(a_j - l' - 1)$, $l, l' = 0, 1, 2, \dots$

Many special functions are particular cases of the H -function. For example, if $\alpha_l = \beta_j = 1$ ($l = 1, 2, \dots, p$; $j = 1, 2, \dots, q$), it reduces to the Meijer G -function $G_{p,q}^{m,n}(\sigma)$ (for definition and properties, see [3, Vol.1]):

$$H_{p,q}^{m,n} \left[\sigma \left| \begin{matrix} (a_p, \alpha_p) \\ (b_q, \beta_q) \end{matrix} \right. \right] = G_{p,q}^{m,n} \left[\sigma \left| \begin{matrix} (a_p) \\ (b_q) \end{matrix} \right. \right]. \quad (2.9)$$

On the other hand, the Fox-Wright Ψ -function is the following special case of the H -function, see for example in [15]:

$${}_p\Psi_q[(a_p, \alpha_p); (b_q, \beta_q); x] = H_{p,q+1}^{1,p} \left[-x \left| \begin{matrix} (1-a_p, \alpha_p) \\ (0, 1), (1-b_q, \beta_q) \end{matrix} \right. \right]. \quad (2.10)$$

Next we use of the following Mellin transform of a product of two hypergeometric functions, proven by Miller and Srivastava [9] (see also [12, Sect.2.22, p.333])

$$\begin{aligned} F(s) &= \int_0^\infty r^{s-1} {}_0F_1(-; 1+\mu; -a^2r^2) {}_1F_2(\alpha; \beta, 1+\nu; -b^2r^2) dr \\ &= \frac{1}{2a^s} \frac{\Gamma(1+\mu)\Gamma(1+\nu)\Gamma(\beta)}{\Gamma(\alpha)} G_{3,3}^{1,2} \left[\frac{b^2}{a^2} \left| \begin{matrix} 1-s/2, 1-\alpha, 1+\mu-s/2 \\ 0, -\nu, 1-\beta \end{matrix} \right. \right] \end{aligned} \quad (2.11)$$

$$(a > 0, b > 0, 0 < \Re(s) < \Re(2+\mu+\nu+\beta-\alpha), 0 < \Re(s) < \Re(\frac{3}{2}+2\alpha+\mu)).$$

Substituting the relation (see [11, p.727])

$${}_0F_1(-; 1+\mu; -a^2r^2) = \frac{\Gamma(1+\mu) J_\mu(2ar)}{(ar)^\mu} \quad (2.12)$$

into the integral formula (2.11) with $\mu \rightarrow \nu, s = \mu + \frac{3}{2} \rightarrow \nu + \frac{3}{2}, a \rightarrow \frac{x}{2}, \alpha \rightarrow \mu, \beta \rightarrow \mu - \frac{\beta}{\alpha}, \nu = \mu - \frac{\beta}{\alpha} - \frac{1}{2}, b = \frac{t}{2}$, we get

$$\begin{aligned} &\int_0^\infty r^{1/2} J_\nu(rx) {}_1F_2\left(\mu; \mu - \frac{\beta}{\alpha}, \mu - \frac{\beta}{\alpha} + \frac{1}{2}; -\frac{r^2t^2}{4}\right) dr = \frac{\sqrt{2}}{x^{3/2}} \\ &\times \frac{\Gamma\left(\mu - \frac{\beta}{\alpha} + \frac{1}{2}\right) \Gamma\left(\mu - \frac{\beta}{\alpha}\right)}{\Gamma(\mu)} G_{3,3}^{1,2} \left[\frac{t^2}{x^2} \left| \begin{matrix} 1/4 - \mu/2, 1-\mu, 1/4 + \nu - \mu/2 \\ 0, -\mu + \beta/\alpha + 1/2, 1-\mu + \eta\alpha \end{matrix} \right. \right] \\ &(\Re(\nu) > -\frac{3}{2}, \Re(\mu) > 0, \Re(\mu - 2\frac{\beta}{\alpha}) > 0, t > 0, x > 0). \end{aligned} \quad (2.13)$$

2.4. In order to evaluate the Hankel transform of $S_\mu^{(\alpha, \beta)}(r; \{k^\gamma\})$, we apply the known integral formula from [11, p.355] with $p = 1, q = 2, r = 2, \alpha = 3/2$:

$$\begin{aligned}
& \int_0^\infty x^{1/2} J_\nu(\sigma x) H_{1,2}^{1,1} \left[\omega x^2 \left| \begin{matrix} (1-\mu, 1) \\ (0, 1), (1-\gamma(\mu\alpha-\beta), \gamma\alpha) \end{matrix} \right. \right] dx \\
&= \frac{\sqrt{2}}{\sigma^{3/2}} H_{3,2}^{1,2} \left[\frac{4\omega}{\sigma^2} \left| \begin{matrix} (1/4-\nu/2, 1), (1-\mu, 1), (1/4+\nu/2, 1) \\ (0, 1), (1-\gamma(\mu\alpha-\beta), \gamma\alpha) \end{matrix} \right. \right] \quad (2.14) \\
&\quad \left(\omega \in \mathbb{R}, \sigma \in \mathbb{R}^+, \Re(\mu) > 0, \Re(\nu) > -\frac{3}{2} \right).
\end{aligned}$$

Using relation (2.10) with $p = 1, q = 1$, by (2.14) we get the following integral formula:

$$\begin{aligned}
& \int_0^\infty x^{1/2} J_\nu(\sigma x)_1 \Psi_1[(\mu, 1); (\gamma(\mu\alpha-\beta), \gamma\alpha); -x^2 t^{\gamma\alpha}] dx \\
&= \frac{\sqrt{2}}{\sigma^{3/2}} H_{3,2}^{1,2} \left[\frac{4t^{\gamma\alpha}}{\sigma^2} \left| \begin{matrix} (1/4-\nu/2, 1), (1-\mu, 1), (1/4+\nu/2, 1) \\ (0, 1), (1-\gamma(\mu\alpha-\beta), \gamma\alpha) \end{matrix} \right. \right] \quad (2.15) \\
&\quad \left(t \in \mathbb{R}^+, \sigma \in \mathbb{R}^+, \Re(\mu) > 0, \Re(\nu) > -\frac{3}{2} \right).
\end{aligned}$$

3. Evaluation of the Hankel transform

The Hankel transform of order ν is defined by

$$\mathbf{H}_\nu(f(r))(x) = \int_0^\infty f(r) J_\nu(rx) \sqrt{rx} dr, \quad (3.1)$$

where J_ν is the Bessel function of the first kind of order ν with $\Re(\nu) \geq -\frac{1}{2}$.

In view of the relationship (2.1) and

$$J_{-1/2}(x) = \sqrt{\frac{2}{\pi x}} \cos x, \quad (3.2)$$

the Hankel transform reduces to the sin-Fourier and cos-Fourier transforms:

$$(\mathcal{F}_s f)(x) = \sqrt{\frac{2}{\pi}} \int_0^\infty \sin(rx) f(r) dr, \quad (\mathcal{F}_c f)(x) = \sqrt{\frac{2}{\pi}} \int_0^\infty \cos(rx) f(r) dr.$$

For a generalization of the Hankel type transform as well as of the sin-Fourier and cos-Fourier transforms, see Luchko and Kiryakova [7].

3.1. Using the integral formula (2.4), we first evaluate the Hankel transform of $S(r)$:

$$\begin{aligned} \mathbf{H}_\nu(S(r))(x) &= \int_0^\infty S(r) J_\nu(rx) \sqrt{rx} dr = \sqrt{x} \int_0^\infty \frac{1}{\sqrt{r}} \left(\int_0^\infty \frac{t \sin(rt)}{e^t - 1} dt \right) J_\nu(rx) dr \\ &= \sqrt{x} \int_0^\infty \frac{t}{e^t - 1} \left(\int_0^\infty r^{-1/2} \sin(rt) J_\nu(rx) dr \right) dt \\ &= \sqrt{\frac{\pi x}{2}} \left(\int_0^x \frac{t}{e^t - 1} \Omega_\nu^{(2)}(x; t) dt + \int_x^\infty \frac{t}{e^t - 1} \Omega_\nu^{(1)}(x; t) dt \right). \end{aligned} \quad (3.3)$$

Using the relation [6, p.1041]

$${}_2F_1\left(1, \frac{1}{2}; \frac{3}{2}; z^2\right) = \frac{1}{2z} \ln \frac{1+z}{1-z},$$

we get

$$\begin{aligned} \Omega_{1/2}^{(1)}(x; t) &= \frac{\sqrt{x}}{t \Gamma\left(\frac{3}{2}\right) \Gamma\left(\frac{1}{2}\right)} {}_2F_1\left(1, \frac{1}{2}; \frac{3}{2}; \frac{x^2}{t^2}\right) = \frac{1}{\pi \sqrt{x}} \ln \frac{t+x}{t-x} \quad (0 < x < t) \\ \Omega_{1/2}^{(2)}(x; t) &= \frac{t}{x \sqrt{x} \Gamma\left(\frac{3}{2}\right) \Gamma\left(\frac{1}{2}\right)} {}_2F_1\left(1, \frac{1}{2}; \frac{3}{2}; \frac{t^2}{x^2}\right) = \frac{1}{\pi \sqrt{x}} \ln \frac{x+t}{x-t} \quad (0 < t < x). \end{aligned}$$

Hence

$$\begin{aligned} \mathbf{H}_{1/2}(S(r))(x) &= (\mathcal{F}_s S(r))(x) \\ &= \sqrt{\frac{\pi x}{2}} \left(\int_0^x \frac{t}{e^t - 1} \frac{1}{\pi \sqrt{x}} \ln \frac{x+t}{x-t} dt + \int_x^\infty \frac{t}{e^t - 1} \frac{1}{\pi \sqrt{x}} \ln \frac{t+x}{t-x} dt \right) \\ &= \frac{1}{\sqrt{2\pi}} \left(\int_0^x \frac{t}{e^t - 1} \ln \frac{x+t}{x-t} dt + \int_x^\infty \frac{t}{e^t - 1} \ln \frac{t+x}{t-x} dt \right) \\ &= \frac{1}{\sqrt{2\pi}} PV \left(\int_0^\infty \frac{t}{e^t - 1} \ln \left| \frac{t+x}{t-x} \right| dt \right) \quad (x > 0), \end{aligned} \quad (3.4)$$

where the Cauchy Principal Value (PV) of the last integral is assumed to exist. By a direct computational, the same formula (3.4) was recently proved by Elezovic et al [4].

3.2. The integral representation of $S_{\mu+1}(r)$, obtained by Cerone and Lenard in [1], is given by

$$S_{\mu+1}(r) = \frac{\sqrt{\pi}}{(2r)^{\mu-1/2} \Gamma(\mu+1)} \int_0^\infty \frac{t^{\mu+1/2}}{e^t - 1} J_{\mu-1/2}(rt) dt \quad (r, \mu \in \mathbb{R}^+). \quad (3.5)$$

Applying integral formula (2.5), we obtain

$$\begin{aligned} \mathbf{H}_\nu(S_{\mu+1}(r))(x) &= \int_0^\infty S_{\mu+1}(r) J_\nu(rx) \sqrt{rx} dr \\ &= \frac{\sqrt{\pi x}}{2^{\mu-1/2} \Gamma(\mu+1)} \int_0^\infty r^{1-\mu} \left(\int_0^\infty \frac{t^{\mu+1/2}}{e^t - 1} J_{\mu-1/2}(rt) dt \right) J_\nu(rx) dr \\ &= \frac{\sqrt{\pi x}}{2^{\mu-1/2} \Gamma(\mu+1)} \int_0^\infty \frac{t^{\mu+1/2}}{e^t - 1} \left(\int_0^\infty r^{1-\mu} J_{\mu-1/2}(rt) J_\nu(rx) dr \right) dt \\ &= \frac{\sqrt{\pi x}}{2^{\mu-1/2} \Gamma(\mu+1)} \left(\int_0^x \frac{t^{\mu+1/2}}{e^t - 1} \Theta_\nu^{(1)}(\mu; x, t) dt + \int_x^\infty \frac{t^{\mu+1/2}}{e^t - 1} \Theta_\nu^{(2)}(\mu; x, t) dt \right) \end{aligned} \quad (3.6)$$

$$(\mu \in \mathbb{R}^+, \Re(\nu) > -\frac{1}{2}, x > 0).$$

3.3. In order to evaluate the Hankel transform of $S_\mu^{(\alpha, \beta)}(r; \{k^{2/\alpha}\})$, we apply its integral representation from [14]:

$$\begin{aligned} S_\mu^{(\alpha, \beta)}(r; \{k^{2/\alpha}\}) &= \frac{2}{\Gamma\left(2\left[\mu - \frac{\beta}{\alpha}\right]\right)} \int_0^\infty \frac{x^{2(\mu - \frac{\beta}{\alpha})-1}}{e^x - 1} \\ &\quad \times {}_1F_2\left(\mu; \mu - \frac{\beta}{\alpha}, \mu - \frac{\beta}{\alpha} + \frac{1}{2}; -\frac{r^2 x^2}{4}\right) dx \\ &\quad \left(r, \alpha, \beta \in \mathbb{R}^+, \mu - \frac{\beta}{\alpha} > \frac{1}{2}\right) \end{aligned} \quad (3.7)$$

and integral formula (2.13). Thus, we have

$$\mathbf{H}_\nu\left(S_\mu^{(\alpha, \beta)}(r; \{k^{2/\alpha}\})\right)(x) = \int_0^\infty S_\mu^{(\alpha, \beta)}(r; \{k^{2/\alpha}\}) J_\nu(rx) \sqrt{rx} dr$$

$$\begin{aligned}
&= \frac{2\sqrt{x}}{\Gamma\left(2\left[\mu - \frac{\beta}{\alpha}\right]\right)} \int_0^\infty \frac{t^{2(\mu - \frac{\beta}{\alpha})-1}}{e^t - 1} \\
&\times \left(\int_0^\infty r^{1/2} {}_1F_2\left(\mu; \mu - \frac{\beta}{\alpha}, \mu - \frac{\beta}{\alpha} + \frac{1}{2}; -\frac{r^2 t^2}{4}\right) J_\nu(rx) dr \right) dt \\
&= \frac{2\sqrt{2\pi}}{2^{2(\mu - \frac{\beta}{\alpha})-1} x \Gamma(\mu)} \left(\int_0^\infty \frac{t^{2(\mu - \frac{\beta}{\alpha})-1}}{e^t - 1} G_{3,3}^{1,2}\left(\frac{t^2}{x^2} \middle| \begin{matrix} \frac{1}{4} - \frac{\mu}{2}, 1 - \mu, \frac{1}{4} + \nu - \frac{\mu}{2} \\ 0, -\mu + \frac{\beta}{\alpha} + \frac{1}{2}, 1 - \mu + \frac{\beta}{\alpha} \end{matrix} \right) dt \right) \\
&\left(\Re(\nu) > -\frac{1}{2}, \mu - 2\frac{\beta}{\alpha} > 0, \mu - \frac{\beta}{\alpha} > \frac{1}{2}, x > 0 \right). \quad (3.8)
\end{aligned}$$

3.4. The integral representation of $S_\mu^{(\alpha, \beta)}(r; \{k^\gamma\}_{k=1}^\infty)$ obtained by Srivastava and Tomovski [14] is given by

$$\begin{aligned}
S_\mu^{(\alpha, \beta)}(r; \{k^\gamma\}_{k=1}^\infty) &= \frac{2}{\Gamma(\mu)} \int_0^\infty \frac{x^{\gamma(\mu\alpha - \beta) - 1}}{e^x - 1} \\
&\times {}_1\Psi_1[(\mu, 1); (\gamma(\mu\alpha - \beta), \gamma\alpha); -r^2 x^{\gamma\alpha}] dx \\
&(r, \alpha, \beta, \gamma \in \mathbb{R}^+, \gamma(\mu\alpha - \beta) > 1).
\end{aligned} \quad (3.9)$$

Using this integral representation and (2.15), we get

$$\begin{aligned}
\mathbf{H}_\nu\left(S_\mu^{(\alpha, \beta)}(r; \{k^\gamma\})\right)(x) &= \int_0^\infty S_\mu^{(\alpha, \beta)}(r; \{k^\gamma\}) J_\nu(rx) \sqrt{rx} dr = \frac{2\sqrt{x}}{\Gamma(\mu)} \\
&\times \int_0^\infty \frac{t^{\gamma(\mu\alpha - \beta) - 1}}{e^t - 1} \left(\int_0^\infty r^{1/2} J_\nu(rx) \Psi_1[(\mu, 1); (\gamma(\mu\alpha - \beta), \gamma\alpha); -r^2 t^{\gamma\alpha}] dr \right) dt \\
&= \frac{2\sqrt{2}}{x \Gamma(\mu)} \int_0^\infty \frac{t^{\gamma(\mu\alpha - \beta) - 1}}{e^t - 1} H_{3,2}^{1,2}\left(\frac{4t^{\gamma\alpha}}{x^2} \middle| \begin{matrix} (\frac{1}{4} - \frac{\nu}{2}, 1), (1 - \mu, 1), (\frac{1}{4} + \frac{\nu}{2}, 1) \\ (0, 1), (1 - \gamma(\mu\alpha - \beta), \gamma\alpha) \end{matrix} \right) dt \\
&\left(\Re(\mu) > 0, \Re(\nu) > -\frac{3}{2}, \gamma(\mu\alpha - \beta) > 1, x > 0 \right). \quad (3.10)
\end{aligned}$$

References

- [1] P. Cerone, C.T. Lenard, On integral forms of generalized Mathieu series. *J. Inequal. Pure Appl. Math.* **4**, No 5 (2003), Article 100, 1-11 (electronic).
- [2] P.H. Diananda, Some inequalities related to an inequality of Mathieu. *Math. Ann.* **250** (1980), 95-98.
- [3] A. Erdélyi, W. Magnus, F. Oberhettinger and F.G. Tricomi (Ed-s), *Higher Transcendental Functions*, Vols. 1-2-3, Mc Graw-Hill, N. York - Toronto - London (1953).
- [4] N. Elezovic, H.M. Srivastava, Ž. Tomovski, Integral representations and integral transforms of some families of Mathieu type series. *Integral Transforms and Special Functions* **19**, No 7 (2008), 481-495.
- [5] O. Emersleben, Über die Reihe. *Math. Ann.* **125** (1952), 165-171.
- [6] I.S. Gradshteyn, I.M. Ryzhik, *Table of Integrals, Series and Products*. Academic Press, N. York - London (1965).
- [7] Yu.F. Luchko, V.S. Kiryakova, Generalized Hankel transforms for hyper-Bessel differential operators. *C.R. Acad. Bulgare Sci.* **53**, No 8 (2000), 17-20.
- [8] E.L. Mathieu, *Traité de Physique Mathématique. VI-VII: Théorie de l'Elasticité des Corps Solides (Part 2)*. Gauthier-Villars, Paris (1890).
- [9] A.R. Miller and H.M. Srivastava, On the Mellin transform of a product of hypergeometric functions, *J. Austral. Math. Soc. Ser. B* **40** (1998), 222-237.
- [10] T.K. Pogany, H.M. Srivastava, Ž. Tomovski, Some families of Mathieu a-series and alternating Mathieu a-series. *Appl. Math. Computation* **173** (2006), 69-108.
- [11] A.P. Prudnikov, Yu.A. Brychkov, O.I. Marichev, *Integrals and Series, Vol. 1: Elementary Functions; Vol. 2: Special Functions*. Gordon and Breach Sci. Publ., N. York (1992); Russian original: Nauka, Moscow (1981).

- [12] A.P. Prudnikov, Yu.A. Brychkov, O.I. Marichev, *Integrals and Series (More Special Functions)*. Gordon and Breach Sci. Publ., N. York (1990); Russian original: Nauka, Moscow (1981).
- [13] F. Qi, An integral expression and some inequalities of Mathieu type series. *Rostock. Math. Kolloq.* **58** (2004), 37-46.
- [14] H.M. Srivastava, Ž. Tomovski, Some problems and solutions involving Mathieu's series and its generalizations. *J. Inequal. Pure Appl. Math.* **5**, No 2 (2004), Article 45, 1-13 (electronic).
- [15] H.M. Srivastava and H.L. Manocha, *A Treatise of Generating Functions*. Halsted Press (Ellis Horwood Ltd, Chichester), J. Wiley & Sons, N. York - Chichester - Brisbane -Toronto (1984).
- [16] Ž. Tomovski, K. Trencevski, On an open problem of Bai-Ni Guo and Feng Qi. *J. Inequal. Pure Appl. Math.* **4**, No 2 (2003), Article 29, 1-7 (electronic).
- [17] Ž. Tomovski, New double inequality for Mathieu series, *Univ. Beograd Publ. Elektrotehn. Fak. Ser. Mat.* **15** (2004), 79-83.
- [18] Ž. Tomovski, Integral representations of generalized Mathieu series via Mittag-Leffler type functions. *Fract. Calc. and Appl. Anal.* **10**, No 2 (2007), 127-138.
- [19] Ž. Tomovski, New integral and series representations of the generalized Mathieu series. *Appl. Anal. Discrete Math.* **2**, No 2 (2008), 205-212.
- [20] Ž. Tomovski, Vu Kim Tuan, On Fourier transforms and summation formulas of generalized Mathieu series. *Math. Science Res. J.*, To appear (2009).

Received: August 25, 2008

University "St. Cyril and Methodius"
Faculty of Natural Sciences and Mathematics
Institute of Mathematics
P.O. Box 162, 1000 – Skopje, Republic of MACEDONIA
e-mail: tomovski@iunona.pmf.ukim.edu.mk