

3. Hörmander, L.: *The Analysis of Linear Partial Differential Operators IV*, Springer, Berlin 1985.
4. Isozaki, H.: Differentiability of generalized Fourier transforms associated with Schrödinger operators, *J. Math. Kyoto Univ.* **25**, 789–806 (1985).
5. Isozaki, H. and H. Kitada: Modified wave operators with time-independent modifiers, *J. Fac. Sci. Univ. Tokyo, Sect. IA, Math.* **32**, 77–104 (1985).
6. Isozaki, H. and H. Kitada: Scattering matrices for two-body Schrödinger operators, *Sci. Papers College Arts Sci. Univ. Tokyo* **35**, 81–107 (1985).
7. Jensen, A.: Commutator methods and a smoothing property of the Schrödinger evolution group, *Math. Z.* **191**, 53–59 (1986).
8. Jensen, A. and H. Kitada: Fundamental solutions and eigenfunction expansions for Schrödinger operators II. Eigenfunctional expansions, *Math. Z.* **199**, 1–13 (1988).
9. Jensen, A., and S. Nakamura: Mapping properties of wave and scattering operators for two-body Schrödinger operators, preprint, Aalborg University, 1991.
10. Poeschke, T. and G. Stolz: On eigenfunction expansions and scattering theory, preprint, Johann Wolfgang Goethe-Universität, Frankfurt am Main, 1990.
11. Reed, M. and B. Simon: *Methods of Modern Mathematical Physics III: Scattering Theory*, Academic Press, New York 1979.
12. Saito, Y.: *Spectral Representation of Schrödinger Operators with Long-Range Potentials*, Lecture Notes Math. **727**, Springer, Berlin 1979.
13. Simon, B.: Schrödinger semigroups, *Bull. Amer. Math. Soc.* **7**, 447–526 (1982).

Received September 13, 1991.

## HARMONICITY MODULUS AND APPLICATIONS TO THE APPROXIMATION BY POLYHARMONIC FUNCTIONS

O. I. KOUNCHEV

*Institute of Mathematics, Bulgarian Academy of Sciences,  
Acad. G. Bonchev St. 8, 1113 Sofia, Bulgaria*

**Abstract.** In the present paper we introduce the notion of harmonicity modulus and harmonicity  $K$ -functional and apply these notions to prove a Jackson type theorem for approximation of continuous functions by polyharmonic functions. For corresponding results on approximation by polynomials see [3, 7].

**Key words:** Harmonicity modulus,  $K$ -functional, polyharmonic functions, Jackson type theorem.

### 0. Notions and Notations

Suppose that  $D \subset \mathbb{R}^n$  is an open, connected and bounded set ( $n \geq 2$ ). We shall work with functions  $f$  in the space  $HC^r(\overline{D})$ ,  $r \geq 0$ , consisting of all functions  $f$  such that  $\Delta^r f$ , the  $r$ th power of the Laplacian, exists and is continuous in  $D$ . In the space  $HC^0(\overline{D}) = C(\overline{D})$  of functions which are continuous in  $\overline{D}$  the usual norm is

$$\|f\| := \max_{x \in \overline{D}} |f(x)|.$$

By  $B(x; t)$  we will denote an open ball in  $\mathbb{R}^n$ :

$$B(x; t) := \{y \in \mathbb{R}^n : |x - y| < t\}.$$

For the function  $f$ , any point  $x \in D$ , and a sufficiently small positive number  $h$  we will consider the spherical mean

$$(1) \quad \mu_0(x, h) := \mu_0(f; x, h) := \frac{1}{\omega_n} \int_{\Omega_\xi} f(x + h\xi) d\omega_\xi;$$

here  $\Omega_\xi$  denotes the unit sphere in  $\mathbb{R}^n$ ,  $\omega_n$  denotes its area and  $d\omega_\xi$  is the area element on  $\Omega_\xi$ .

Further we define the quantity

$$(2) \quad \Delta_h(f; x) := \mu_0(f; x, h) - f(x).$$

Throughout the paper we shall use the symbol  $C$  as an universal constant.

## 1. Harmonicity Modulus

DEFINITION 1. The harmonicity modulus of the function  $f$  in the domain  $D$  is defined by

$$(3) \quad \omega^h(u) := \omega^h(f; u) := \sup |\Delta_t(f; x)|,$$

where the sup is taken over  $0 < t \leq u$ , and  $B(x; t) \subset D$ .

REMARK. It is clear that

$$\omega^h(f; u) \leq \omega_1(f; u),$$

where  $\omega_1$  is the usual first modulus of continuity (see [5, 9]).

It is easy to see that we have the representation

$$\Delta_h(f; x) = \frac{1}{2\omega_n} \int_{\Omega_\xi} (f(x + h\xi) - 2f(x) + f(x - h\xi)) d\omega_\xi.$$

This implies

$$\omega^h(f; u) \leq \omega_2(f; u)$$

for the usual second modulus of continuity (cf. [5, 9]).

PROPOSITION 1. For every function  $f$ , continuous in  $\bar{D}$ , the harmonicity modulus has the following properties:

$$1) \quad \lim_{t \rightarrow 0} \omega^h(f; t) = 0;$$

2)  $\omega^h(f; u)$  is a monotone increasing function;

3) for every positive  $u$  the inequality

$$\omega^h(f + g; u) \leq \omega^h(f; u) + \omega^h(g; u)$$

holds;

4) for every positive number  $u$  the inequality

$$\omega^h(f; u) \leq 2\|f\|$$

holds.

Proof: Property 1) follows from the definition of  $\mu_0$  and the continuity of the function  $f$ . Properties 2) and 3) are evident. Property 4) follows from the easy-to-check representation

$$(4) \quad \Delta_h(f; x) = \frac{1}{\omega_n} \int_{\Omega_\xi} (f(x + h\xi) - f(x)) d\omega_\xi.$$

Let us introduce the integral operator  $J_0$  by

$$(5) \quad J_0[\phi; R] := \int_0^R (r - r^{n-1} R^{-n+2}) \phi(r) dr$$

for  $n \geq 3$ , and by

$$(6) \quad J_0[\phi; R] := \int_0^R r \log(R/r) \phi(r) dr$$

for  $n = 2$ .

Further we will need different forms of the classical Pizzetti formula for the representation of the spherical means (see [2, 8]).

THEOREM 1. Let the function  $f$  have a continuous Laplacian  $\Delta f$  in the domain  $D$ . Then the following representation holds:

$$(7) \quad \mu_0(f; x, R) = f(x) + l_n J_0[\mu_0(\Delta f; x, \cdot); R],$$

where  $l_n = \frac{1}{(n-2)}$  for  $n \geq 3$  and  $l_2 = 1$ .

The remainder can also be written as

$$J_0[\mu_0(\Delta f; x, \cdot); R] = \mu_0(\Delta f; x; \vartheta R) J_0[1, R]$$

with some number  $\vartheta = \vartheta(x; R)$  such that  $0 < \vartheta < 1$ . Since  $J_0[1; R] = c_n R^2$ , where  $c_2 = \frac{1}{4}$ ,  $c_n = \frac{(n-2)}{(2n)}$  for  $n \geq 3$ , we have the representation

$$(8) \quad \mu_0(f; x, R) = f(x) + d_n R^2 \Delta f(\xi),$$

where the point  $\xi = \xi(x, R) \in B(x; R)$  and  $d_n = c_n l_n = \frac{1}{2n}$ .

PROPOSITION 2. Suppose that the function  $f$ , defined and continuous in  $\bar{D}$ , has a continuous Laplacian  $\Delta f$  in  $\bar{D}$  satisfying the inequality

$$|\Delta f(x)| \leq M, \quad x \in \bar{D}.$$

Then the following inequality holds for every positive number  $u$ :

$$\omega^h(f; u) \leq M d_n u^2.$$

The proof follows immediately from Pizzetti's formula (8). ■

The harmonicity modulus plays a role similar to that of the second modulus of continuity in the one-dimensional case (see [3]). This is well seen from the following classical result (cf. [8]).

**THEOREM 2.** *Let  $u$  be a function defined and integrable in the domain  $D$  in  $\mathbb{R}^n$ . Then, if  $t > 0$ , we have*

$$\omega^h(f; t) = 0$$

*if and only if  $f$  is harmonic in  $D$ , i.e.*

$$\Delta f(x) = 0, \quad x \in D.$$

Theorem 2 is the motivation for calling  $\omega^h$  the harmonicity modulus. We also recall that harmonic functions are considered to be a multivariate analogue to the linear functions in one dimension.

### 2. Harmonicity $K$ -Functional

Here we introduce the notion of harmonicity  $K$ -functional which provides a basic tool for studying the important properties of the harmonicity modulus.

**DEFINITION 2.** *For every function  $f \in C(\overline{D})$  and every number  $t > 0$  we define the harmonicity  $K$ -functional by*

$$(9) \quad K^h(f; t) := \inf\{\|f - g\| + t^2\|\Delta g\|\},$$

*where the infimum is taken over all functions  $g \in HC^1(\overline{D})$ .*

### 3. Harmonicity Modulus and Harmonicity $K$ -Functional

The main technical result of the paper is proved in the present and the next sections. Roughly speaking, it states that the harmonicity modulus and the harmonicity  $K$ -functional are equivalent on compact subdomains of  $D$ . The simple part of the equivalence is the following

**LEMMA 1.** *Let  $D$  be an open set in  $\mathbb{R}^n$ . For all  $t$  with  $0 < t < \infty$  and  $f \in C(\overline{D})$ , the inequality*

$$(10) \quad \omega^h(f; t) \leq CK^h(f; t)$$

*holds with some constant  $C > 0$ .*

*Proof:* The proof is based on a standard argument. We split  $f = f - g + g$ , and apply Propositions 1 and 2 to obtain the inequality

$$\begin{aligned} \omega^h(f; t) &\leq \omega^h(f - g; t) + \omega^h(g; t) \leq 2\|f - g\| + d_n t^2 \|\Delta g\| \\ &\leq \max\{2, d_n\} (\|f - g\| + t^2 \|\Delta g\|). \end{aligned}$$

Since  $g \in HC^1(\overline{D})$  is arbitrary, the statement (10) is proved. ■

The domination of  $K^h$  by  $\omega^h$  will be established only on compact subdomains of  $D$  in the sense that the  $K$ -functional of the subdomain  $D_1, K_{D_1}^h$ , will be proved to be dominated by the harmonicity modulus  $\omega_{D_1}^h$  with respect to the domain  $D$ .

The problem is that for every  $R$  (possibly such that  $R < R_1$  for some sufficiently small positive number  $R_1$ ) we have to find a function  $g_R \in HC^1(\overline{D})$  such that

$$(11) \quad \|f - g_R\| + R^2 \|\Delta g_R\| \leq C\omega^h(f; R),$$

where the constant  $C$  does not depend on  $f$  and  $R$ .

Following the scheme given in [5], taking some spherical means of the function  $f$ , we succeed in constructing the function  $g_R$  not on the whole of  $D$  but on every subdomain  $D_1$ , such that  $\overline{D_1} \subset D$  and  $R_1 \leq \text{dist}(D_1, \partial D)$ . In such a way we can prove the inequality (11) over subdomains where the norm  $\|\cdot\|$  is in fact  $\|\cdot\|_{D_1}$ .

### 4. Domination of $K^h$ by $\omega^h$ on Compact Subdomains

Having in mind Pizzetti's formula (7) in Theorem 1, we consider the function

$$(12) \quad g_{R,t}(x) = v(t) J_{0,s}[\mu_0(f; x, R_s); t].$$

Here  $J_0$  is the operator given by (5) and (6) and  $J_{0,s}$  means that  $s$  is the input variable for  $J_0$ ; the output variable is  $t$ ;  $v(t)$  is equal to  $(J_0[1; t])^{-1}$ , where  $J_0[1; t]$  is the value of the functional for  $\phi(t) = 1$ , so in fact  $\frac{1}{v(t)} = t^2(\frac{1}{2} - \frac{1}{n})$  for  $n \geq 3$ ,  $\frac{1}{v(t)} = \frac{2}{4}$  for  $n = 2$ .

The operator  $J_0$  changes the output in a specific way described by

**PROPOSITION 3.** *For every integrable function  $\phi$  and positive numbers  $s$  and  $R$  we have*

$$(13) \quad J_{0,t}[\phi(st); R] = \frac{1}{s^2} J_{0,t}[\phi(t); sR].$$

*Proof:* We give the proof for  $n \geq 3$ . Then  $J_0$  is given by formula (5). The case  $n = 2$  is similar.

By changing the variables we obtain

$$\begin{aligned} J_{0,t}[\phi(st); R] &= \int_0^R (t - t^{n-1} R^{-n+2}) \phi(st) dt \\ &= \frac{1}{s^2} \int_0^{sR} [t - t^{n-1} (sR)^{-n+2}] \phi(t) dt \\ &= \frac{1}{s^2} J_{0,t}[\phi(t); sR]. \quad \blacksquare \end{aligned}$$

Proposition 3 shows that (12) becomes

$$(14) \quad g_{R,t}(x) = v(t) \frac{1}{R^2} J_0[\mu_0(f; x, \cdot); tR].$$

The following is the main technical result of the paper.

**THEOREM 3.** For every subdomain  $D_1$  such that  $\overline{D_1} \subset D$ , the inequality

$$(15) \quad K^h(f; R)_{D_1} \leq C\omega^h(f; R)_D$$

holds for every number  $R$  with

$$(16) \quad 0 < R < d = \text{dist}(D_1, \partial D);$$

here the constant  $C$  does not depend on  $f$  and  $R$ , and  $K^h(f; R)_{D_1}$  denotes the harmonicity  $K$ -functional for the domain  $D_1$ , while  $\omega^h(f; R)_D$  denotes the harmonicity modulus on  $D$ .

*Proof:* Let us notice first that the function  $g_{R,t}$ , given by (12), is well defined in  $D_1$  for  $R$  satisfying (16) and every number  $t$  with  $0 \leq t \leq 1$ . Since  $v(t)J_0[1; t] = 1$ , we obtain

$$g_{R,t}(x) - f(x) = v(t)J_{0,s}[\mu_0(f; x, Rs) - f(x); t]$$

for every number  $R$  satisfying (16) and every number  $t$  with  $0 \leq t \leq 1$ . Hence, for every  $x \in D_1$  we obtain the inequalities

$$(17) \quad |g_{R,t}(x) - f(x)| \leq v(t)J_{0,t}[1; t]\omega^h(f; Rt) \leq \omega^h(f; R).$$

Consequently, we have proved the domination of the first term of  $K^h$  by  $\omega^h$ :

$$\|g_{R,t} - f\| \leq \omega^h(f; R).$$

For proving the domination of the second term, we will check the value of  $\Delta g_{R,t}(x)$  for  $x \in D_1$ .

First let us suppose that  $f$  is twice differentiable in  $D$ , i.e.  $f \in C^2(D)$ . By formula (14) the Laplacian of  $g_{R,t}(x)$  is then equal to

$$(18) \quad \Delta g_{R,t}(x) = v(t) \frac{1}{R^2} J_{0,s}[\mu_0(\Delta f; x, s); tR].$$

Hence, combining with formula (7), we obtain

$$(19) \quad \Delta g_{R,t}(x) = v(t) \frac{1}{l_n R^2} [\mu_0(f; x, tR) - f(x)].$$

This implies the inequalities

$$(20) \quad R^2 |\Delta g_{R,t}(x)| \leq v(t) \frac{1}{l_n} \omega^h(f; tR) \leq C\omega^h(f; R)$$

for every  $x \in D_1$  and every  $R < \text{dist}(D_1, \partial D)$ , where  $C$  is a constant given by

$$C := |\nu(t_1)|/l_n$$

and  $t_1$  is an arbitrary number with  $0 < t_1 < 1$  satisfying  $v(t_1) \neq 0$ .

Inequality (20) implies that the second term of  $K^h$  is dominated by  $\omega^h$ . This ends the proof for  $f \in C^2(D)$ .

In the case of an arbitrary continuous function  $f$ , let us take an approximation to  $f$ , say  $f_\delta \in C^\infty$ ,  $\delta > 0$ , such that  $f_\delta$  converges to  $f$  uniformly on  $D_1$  for  $\delta \rightarrow 0$  (see this construction in [1, paragraph 5]).

Since  $\overline{D_1}$  is a compact set and  $f$  is continuous in  $D$ , we obtain by a standard limiting argument that formula (7) takes for an arbitrary integrable function  $f$  the form

$$\mu_0(f; x, R) = f(x) + l_n \Delta J_0[\mu_0(f; x, \cdot); R].$$

This implies that the relations (19) and (20) hold as well. ■

Now we are ready to prove an important property of the harmonicity modulus.

For an arbitrary subdomain  $D_1$  of  $D$  let us denote by  $\omega^h(\cdot; D_1)$  the harmonicity modulus for the set  $D_1$ .

**THEOREM 4.** For every subdomain  $D_1$  of  $D$  such that  $\overline{D_1} \subset D$ , the following inequalities hold:

$$(i) \quad \omega^h(f; \lambda R)_{D_1} \leq C(\lambda + 1)^2 \omega^h(f; \tau)D$$

for every number  $R \leq d = \text{dist}(D_1, \partial D)$  and every number  $\lambda \geq 0$  such that  $\lambda R \leq d$ ;

$$(ii) \quad \omega^h(f; a + b)_{D_1} \leq C[\omega^h(f; a)_D + \omega^h(f; b)_D]$$

for all positive real numbers  $a$  and  $b$ .

*Proof:* Inequality (i) follows from a similar inequality for the harmonicity  $K$ -functional. Indeed, since

$$\|f - g\|_{D_1} + (\lambda u)^2 \|\Delta g\|_{D_1} \leq (\lambda + 1)^2 (\|f - g\|_{D_1} + u^2 \|\Delta g\|_{D_1})$$

for an arbitrary function  $g \in H^{C^1}(\overline{D_1})$ , by the definition of the harmonicity  $K$ -functional we obtain the inequality

$$K^h(f; \lambda u)_{D_1} \leq (\lambda + 1)^2 K^h(f; u)_{D_1},$$

for every number  $\lambda \geq 0$ .

Lemma 1 gives

$$\omega^h(f; \lambda R)_{D_1} \leq C K^h(f; \lambda R)_{D_1},$$

and Theorem 3 implies

$$K^h(f; R)_{D_1} \leq C \omega^h(f; R)_{D_1}$$

for every number  $R \leq d$ . These inequalities imply the inequality (i).

In order to prove inequality (ii) let us note that

$$(a + b)^2 \leq 2(a^2 + b^2)$$

for all real numbers  $a$  and  $b$ . This implies

$$\begin{aligned} \|f - g\|_{D_1} + (a + b)^2 \|\Delta g\|_{D_1} &\leq \|f - g\|_{D_1} + 2(a^2 + b^2) \|\Delta g\|_{D_1} \\ &\leq 2\|f - g\|_{D_1} + 2a^2 \|\Delta g\|_{D_1} + 2\|f - g\|_{D_1} + 2b^2 \|\Delta g\|_{D_1}. \end{aligned}$$

The definition of  $K^h$  implies the inequality

$$K^h(f; a + b)_{D_1} \leq 2(K^h(f; a)_{D_1} + K^h(f; b)_{D_1}).$$

Now inequality (ii) follows by arguments similar to those used for (i). ■

### 5. Polyharmonic Kernels

Here we introduce kernels which are polyharmonic functions and arise naturally from the Jackson type kernels used in approximation theory [3].

Let us recall that the function  $f$  is called polyharmonic of order  $p$  in an open set  $D$ , where  $p$  is a nonnegative integer, if it satisfies the equation

$$\Delta^p f(x) = 0, \quad x \in D;$$

here the iterated Laplacian of order  $p$  is defined inductively by the equations  $\Delta_{k+1} := \Delta(\Delta^k)$  for  $k \geq 0$  and  $\Delta^0 := id$  (see [8]).

Let us remind the notion of Jackson type kernel (cf. [3]).

DEFINITION 3. A kernel of Jackson type of order  $\nu$ , where  $\nu = 1, 2, \dots$ , is defined to be the function given by

$$J_{k,\nu}(t) := (\gamma_k, \nu)^{-1} [\sin(\nu t/2) / \sin(t/2)]^{2k},$$

where  $k$  is a natural number and the constant is

$$\gamma_{k,\nu} := \frac{1}{\pi} \int_{-\pi}^{\pi} [\sin(\nu t/2) / \sin(t/2)]^{2k} dt.$$

For the properties of these kernels we refer to [3]. Through the substitution

$$x = 2 \sin(t/2), \quad t \in [-\pi, \pi], \quad x \in [-2, 2],$$

we obtain the nonperiodic Jackson type kernels:

$$\bar{J}_{k,\nu}(x) = \gamma_{k,\nu} (\bar{\gamma}_{k,\nu})^{-1} J_{k,\nu}[\arccos(1 - x^2/2)];$$

here the constant is

$$\bar{\gamma}_{k,\nu} = \int_{-1}^1 \gamma_{k,\nu} J_{k,\nu}[\arccos(1 - x^2/2)] dx,$$

for  $\nu \in \mathbb{N}$ .

Finally, we define the polyharmonic Jackson type kernels of order  $p$  by the equation

$$\bar{J}_{k,p}(x) = \bar{J}_{k,p}(1/x) = \bar{\gamma}_{k,p} (\check{\gamma}_{k,p})^{-1} \bar{J}_{k,p}(x),$$

for  $p \in \mathbb{N}$ , and for every  $x \in \mathbb{R}^n$  such that  $|x| \leq 2$ ; here the constant is given by

$$\check{\gamma}_{k,p} = \int_0^1 r^{n-1} \bar{\gamma}_{k,p} \bar{J}_{k,p}(r) dr = \int_0^1 r^{n-1} [\sin(\nu t/2) / \sin(t/2)]^{2k} dr,$$

where  $t = \arccos(1 - r^2/2)$ .

THEOREM 5. The polyharmonic Jackson type kernels have the following properties:

- (i) For all natural numbers  $p$  and  $k$ , the kernel  $\bar{J}_{k,p}$  is a nonnegative polyharmonic function of order  $k(p-1) + 1$  and  $\bar{J}_{k,p}(x)$  is defined for every  $x \in \mathbb{R}^n$  satisfying  $|x| \leq 2$ ;
- (ii)  $\int_{B(0,1)} \bar{J}_{k,p}(x) dx = 1$ ;
- (iii) If  $I_i$  is defined by

$$I_i := \int_0^1 t^{i+n-1} \bar{J}_{k,p}(t) dt \quad \text{for nonnegative integers } i,$$

then for  $i < 2k - n$  we have the inequality

$$I_i \leq C p^{-i},$$

and for  $i = 2k - n$  we have the inequality

$$I_i \leq C(\ln p) p^{-i}.$$

The proof of Theorem 5 is based on standard arguments [3] and is given in detail in a forthcoming paper [6].

## 6. A Direct Theorem of Jackson Type

Here we prove an approximation theorem which is analogous to the direct theorem of Jackson for the approximation by polynomials in the one-dimensional case, where the rate of approximation is estimated by the first and the second modulus of continuity (see [3, 7]).

In the multivariate case we approximate by polyharmonic functions and the rate of approximation is estimated by the harmonicity modulus.

Let us first give some necessary notations. Let  $K$  be a polyharmonic function on  $\{x \in \mathbb{R}^n : |x| \leq 1\}$ . Then for every function  $f$ , defined and continuous in the domain  $D$ , we can define the operator

$$(21) \quad T_K[f](x) := \int_{B(x;1)} K(x-u)f(u)du$$

for every  $x \in D$  such that  $\text{dist}(x, \partial D) \leq 1$ .

Let the domain  $D$  be regular in the sense of solvability of the Dirichlet problem (see [4]), and let the function  $f$  be continuous  $D$ . Then there exists a harmonic function  $h_f$  solving the Dirichlet problem in  $\bar{D}$ , i.e.

$$\Delta h_f(x) = 0, \quad x \in D,$$

$$h_f(x) = f(x), \quad x \in \partial D.$$

We shall consider the function  $F_0$  given by the following conditions:

$$F_0(x) := f(x) - h_f(x) \quad \text{for } x \in \bar{D},$$

and

$$F_0(x) := 0 \quad \text{for } x \notin \bar{D}.$$

The function  $F_0$  is evidently continuous on the whole space and it makes sense to consider its harmonicity modulus there or in domains containing  $D$ .

Another interesting feature of the function  $F_0$  is that its harmonicity modulus in  $D$  satisfies

$$\omega^h(F_0; t)_D = \omega^h(f; t)_D.$$

This follows immediately from the Gauss mean value theorem, which states that

$$\mu_0(h_f; x, t) = h_f(x)$$

for every  $x \in D$  and  $t > 0$  such that  $B(x; t) \subset D$ .

Notice that for every domain  $D_1$  such that  $\bar{D} \subset D_1$  we have

$$\omega^h(F_0; t)_{D_1} = \omega^h(F_0; t)_{\mathbb{R}^n}$$

for every positive number  $t \leq \text{dist}(D, \partial D_1)$ . Here  $\omega^h(F_0; t)_{\mathbb{R}^n}$  denotes the harmonicity modulus of the function  $F_0$  in the whole space.

Let  $D_2$  be a domain such that  $\bar{D}_1 \subset D_2$ . Then we can apply Theorem 3 to obtain the following inequalities

$$(22) \quad C_1 \omega^h(F_0; t)_D \leq K^h(F_0; t)_D \leq C_2 \omega^h(F_0; t)_{D_2}$$

for sufficiently small numbers  $t > 0$  and appropriate constants  $C_1, C_2$ , which do not depend on  $f$  and  $t$ .

Next suppose that, for some nonnegative  $r$ , the function  $f$  is in  $H^{Cr}(\bar{D})$ . Then, inductively in  $r$ , we obtain a solution  $h_f$  to the following boundary value problem:

$$\Delta^{r+1} h_f(x) = 0, \quad x \in D;$$

$$\Delta^j h_f(x) = \Delta^j f(x), \quad x \in \partial D,$$

for  $j = 0, 1, \dots, r$ .

We shall consider the function  $F_r$  given by

$$(23) \quad F_r(x) := f(x) - h_f(x) \quad \text{for } x \in \bar{D};$$

$$F_r(x) := 0 \quad \text{for } x \notin \bar{D}.$$

Note that the function  $F_r$  is continuous on the whole space together with  $\Delta^r F_r$  and we can apply to it all properties of the harmonicity modulus, a fact which will be used below.

Now we are ready to state the following result which is the main application of the harmonicity modulus in the present paper.

**THEOREM 6.** *Let the domain  $D$  be regular in the sense of solvability of the Dirichlet problem. Let for some integer  $r \geq 0$ , the function  $f \in H^{Cr}(D)$ . Let us denote by  $F_r$  the function given by (23). Then, for every natural number  $p$  satisfying  $p \geq r + 1$ , there exists a polyharmonic function  $T_p$  of order  $p$  in  $D$  satisfying the inequality*

$$(24) \quad |f(x) - T_p(x)| \leq C \omega^h(\Delta^r F_r; \frac{1}{p})_{\frac{1}{p^2x}}$$

for every  $x \in \bar{D}$ , where the constant  $C$  depends on the domain  $D$  and on  $r$ .

*Proof:* (1) By a similarity transform we can suppose that the domain  $D$  is contained in the ball  $B(0; 1/2)$ . Obviously, this transform preserves the polyharmonic functions. To find the harmonicity modulus for the function  $G(x) = f(\lambda x)$ , where  $\lambda$  is a positive real number, let us compute the harmonicity difference given by (2):

$$\Delta_\lambda(G; x) = \mu_0(G; x, t) - G(x) = \mu_0(f; \lambda x; \lambda t) - f(\lambda x) = \Delta_\lambda(f; \lambda x).$$

Hence, we obtain

$$\omega^h(G; t)_D = \omega^h(f; \lambda t)_{\lambda D},$$

where  $\lambda D$  is the domain given by

$$\lambda D = \{y \in \mathbb{R}^n : y = \lambda x, \quad x \in D\}.$$

So for a domain  $D_1$  such that  $\bar{D}_1 \subset D$  we have

$$\omega^h(G; t)_{D_1} = \omega^h(f; \lambda t)_{\lambda D_1},$$

which proves the inequality

$$\omega^h(G; t)_{D_1} \leq (\lambda + 1)^2 \omega^h(f; t)_{\lambda D}$$

for every number  $t \leq \text{dist}(\lambda D_1, \partial \lambda D)$ .

This shows that the harmonicity modulus is at most multiplied by a constant as a result of a similarity transform. Applied to the modulus  $\omega^h(\Delta^r F_0; p^{-1})_{\mathbb{R}^n}$  we see that by (22) it only changes up to a constant multiple.

(2) We will define the polyharmonic function  $T_p(x) = T_p(f; \tau, x)$  of order  $p$  inductively by the following recurrence relation:

$$(25) \quad \begin{aligned} T_p(x) &:= T_p(F_\tau; m, x) \\ &:= T_p(F_\tau; m - 1, x) + T_{k,\nu}[F_\tau(\cdot) - T_p(F_\tau; m - 1, \cdot)](x), \end{aligned}$$

for every  $x \in \bar{D}$  and every  $m$  with  $1 \leq m \leq \tau$ .

Here  $T_{k,\nu}$  is a short notation for the operator given by formula (21) for the Jackson Type kernel  $\tilde{J}_{k,\nu}$ , where we take  $k$  big enough to satisfy  $2k - n \geq 3$ , and put  $\nu := [(p - 1)/k] + 1$  (here  $[y]$  denotes, as usually, the greatest integer which does not exceed  $y$ ). The choice of such  $\nu$  provides that the order of the polyharmonic function  $\tilde{J}_{k,p}$  be equal to  $k(\nu - 1) + 1 \leq p$ .

Note that the operator  $T_{k,\nu}$  is well defined and produces a polyharmonic function since  $F_\tau$  is a finite function, the kernels are defined in  $B(0; 1)$  and we have the inclusion  $D \subset B(0; 1/2)$ .

(3) Let us check the Theorem for  $\tau = 0$ . In this case we have  $f \in C(\bar{D})$ . Due to Theorem 5 the following holds:

$$\begin{aligned} D(x) &:= F_0(x) - T_{k,\nu}[F_0](x) \\ &= \int_{B(x;1)} [F_0(x) - F_0(u)] \tilde{J}_{k,\nu}(x - u) du \\ &= \int_0^1 \left\{ \int_{\Omega_\xi} [F_0(x) - F_0(x - r\xi)] d\omega_\xi \right\} r^{\nu-1} \tilde{J}_{k,\nu}(r) dr. \end{aligned}$$

By the properties of the harmonicity modulus (see Theorem 4) this gives the following estimate

$$\begin{aligned} |D(x)| &\leq \omega_n \int_0^1 r^{\nu-1} \tilde{J}_{k,\nu}(r) \omega^h(F_0; r) \mathbb{R}^n dr \\ &\leq C \omega^h(F_0; p^{-1})_{\mathbb{R}^n} \omega_n \int_0^1 r^{\nu-1} \tilde{J}_{k,\nu}(r) (pr + 1)^2 dr \end{aligned}$$

for every  $p \geq 1$  and some constant  $C > 0$ .

Again, applying Theorem 5, (iii), since  $2k - n \geq 3$ , we have the inequality

$$\int_0^1 r^{\nu-1} \tilde{J}_{k,\nu}(r) (pr + 1)^2 dr \leq C \int_0^1 r^{\nu-1} \tilde{J}_{k,\nu}(r) (\nu r + br + 1)^2 dr \leq C_1$$

for appropriate constants  $C, C_1$  and  $b$ . The last gives, finally, that

$$(26) \quad |D(x)| \leq C \omega^h(F_0; p^{-1}), \quad x \in \bar{D},$$

for some constant  $C > 0$ . From this estimate we get the statement for  $\tau = 0$ .

(4) Before proceeding by induction on  $\tau$ , let us note the following. If for some function  $\phi$  on some domain  $D$ , such that  $\Delta \phi$  is continuous on  $D$ , the inequality

$$|\Delta \phi(x)| \leq M, \quad x \in D,$$

holds, then by Proposition 2 we obtain the inequality

$$\omega^h(\phi; t) \leq M d_n t^2$$

for every number  $t > 0$ . Hence, by (26), we obtain the inequality

$$(27) \quad |\phi(x) - T_{k,\nu}[\phi](x)| \leq C M \frac{1}{p^2}$$

for an appropriate constant  $C$ .

(5) Let us suppose that the Theorem is true for the classes of functions  $HC^0, HC^1, \dots, HC^r, r \geq 0$ . Then, if  $f \in HC^{r+1}$ , it follows that  $\Delta f \in HC^r$ , and equality (25) implies that

$$\Delta T_p(F_{\tau+1}; \tau, x) = T_p(\Delta F_{\tau+1}; \tau, x).$$

Applied to the function  $\Delta F_{\tau+1}$ , the induction hypothesis (24) gives

$$\begin{aligned} |\Delta[F_{\tau+1}(x) - T_p(F_{\tau+1}; \tau, x)]| &= |\Delta F_{\tau+1}(x) - T_p(\Delta F_{\tau+1}; \tau, x)| \\ &\leq C \omega^h(\Delta^{\tau+1} F_{\tau+1}; p^{-1})_D p^{-2\tau}. \end{aligned}$$

Let us put

$$\phi(x) := F_{r+1}(x) - T_p(F_{r+1}; r, x)$$

and apply inequality (27) to this function  $\phi$ . We obtain the following inequalities:

$$(28) \quad \begin{aligned} |\phi(x) - T_{k,\nu}[\phi](x)| &= |F_{r+1}(x) - T_p(F_{r+1}; r, x) \\ &\quad - T_{k,\nu}[F_{r+1}(\xi) - T_p(F_{r+1}; r, \xi)](x)| \\ &\leq CG_1 p^{-2} \omega^h(\Delta^{r+1} F_{r+1}; p^{-1}) D \frac{1}{p^{2r}} \\ &= CG_1 \omega^h(\Delta^{r+1} F_{r+1}; p^{-1}) D \frac{1}{p^{2(r+1)}}. \end{aligned}$$

On the other hand, by (25) we have

$$\phi(x) - T_{k,\nu}[\phi](x) = F_{r+1}(x) - T_p(F_{r+1}; r+1, x),$$

which shows that the inequality in (28) is exactly inequality (24) for  $r+1$ . This yields the statement of the Theorem for  $r+1$ . ■

**COROLLARY.** *In view of the Remark after Definition 1, in Theorem 6 we can replace inequality (24) by the following inequalities:*

$$|f(x) - T_p(x)| \leq C\omega_1(\Delta^r F_r; \frac{1}{p}) \frac{1}{p^{2r}}$$

or

$$|f(x) - T_p(x)| \leq C\omega_2(\Delta^r F_r; \frac{1}{p}) \frac{1}{p^{2r}}$$

for  $x \in \bar{D}$ , where  $\omega_1$  and  $\omega_2$  are the usual first and second moduli of continuity (see [5]).

#### Acknowledgements

The author wishes to thank Professor Walter K. Hayman and the reviewer for their invaluable remarks.

#### References

1. Besov, O. V., V. P. Il'in and S. M. Nikol'skii: *Integral Representation of Functions and Imbedding Theorems*, Nauka, Moscow 1975.
2. Courant, R. and D. Hilbert: *Methoden der mathematischen Physik II*, Springer, Berlin-Heidelberg-New York 1968.
3. Dzjadzyk, V. K.: *Introduction to the Theory of Uniform Approximation of Functions by Polynomials*, Nauka, Moscow 1977.
4. Helms, L. L.: *Einführung in die Potentialtheorie*, de Gruyter, Berlin-New York 1973.
5. Johnen, H., and K. Scherer: On the equivalence of the  $K$ -functional and moduli of continuity and some applications, in: *Const. Theory of Funct. of Several Variables*, Lecture Notes Math. 571, pp. 119-140, Springer, Berlin-Heidelberg-New York (1977).
6. Kounchev, O. I.: Harmonicity moduli and Jackson type theorems for the approximation through polyharmonic functions (preprint).
7. Meinardus, G.: *Approximation of Functions: Theory and Numerical Methods*, Springer, Berlin-Heidelberg-New York 1967.
8. Nicolescu, M.: *Opera Matematică. Funcții Poliharmonice*, Editura Academiei, București 1980.
9. Nikol'skii, S. M.: *Approximation of Functions of Several Variables and Imbedding Theorems*, Springer, Berlin-Heidelberg-New York 1975.

Received September 16, 1991.