

The approximation order of polysplines

O. Kounchev and H. Render

April 2, 2002

Abstract

We show that the scaling spaces defined by the polysplines of order p provide approximation order $2p$. For that purpose we refine the results on one-dimensional approximation order by L -splines obtained in [2].

AMS (MOS) Classification. Primary 41A15; Secondary 35J40, 31B30.

Key words and phrases. cardinal splines, cardinal L -splines, polysplines, approximation order of splines, polyharmonic functions, cardinal polysplines.

Acknowledgement. The first author was partially sponsored by the Fulbright Program during his stay at the University of Wisconsin-Madison. The second author thanks the Alexander von Humboldt-Foundation for supporting him in the framework of the Feodor-Lynen-program.

1 Introduction

In the last decade the approximation order of shift-invariant subspaces of the space $L^2(\mathbb{R}^n)$ of all square-integrable functions on the euclidean space \mathbb{R}^n has been investigated extensively, e.g. in the survey paper [10] approximately 100 references are given. The problem can be formulated in a rather general way: suppose that $(V_h)_{h \in I}$ is a family of subspaces of $L^2(\mathbb{R}^n)$ (not necessarily shift-invariant) where I is subset of $(0, \infty)$ having 0 as an accumulation point. One has to estimate the rates of decay of the approximation error

$$E(f, V_h) := \inf \left\{ \|f - s\|_{L^2(\mathbb{R}^n)} : s \in V_h \right\} \quad (1)$$

for h tending to 0. If W is a subspace of $L^2(\mathbb{R}^n)$ endowed with a norm $\|\cdot\|_W$ we say that $(V_h)_{h \in I}$ provides approximation order m with respect to the norm $\|\cdot\|_W$ if there exists a constant c_W such that for every $f \in W$ and for every $h \in I$

$$E(f, V_h) \leq c_W \cdot h^m \cdot \|f\|_W . \quad (2)$$

Usually W is the potential space $W_2^m(\mathbb{R}^n)$ for $m \in (0, \infty)$ defined as the subspace of those $f \in L^2(\mathbb{R}^n)$ such that

$$\|f\|_{W_2^m(\mathbb{R}^n)} := (2\pi)^{-\frac{n}{2}} \left\| (1 + |\xi|)^m \widehat{f}(\xi) \right\|_{L^2(\mathbb{R}^n)} < \infty. \quad (3)$$

In this note we want to prove that cardinal polysplines of order p provide approximation order $2p$.

The motivation for the present work comes from the fact that polysplines are useful for solving multivariate interpolation problems [4], [5], [6] and they are of importance for the multivariate Wavelet Analysis, cf. the monograph [9]. Recall that a function $S : \mathbb{R}^n \setminus \{0\} \rightarrow \mathbb{C}$ is called a *cardinal polyspline*¹ (on annuli) of order p if S is a $(2p - 2)$ -times continuously differentiable and the restriction of S to each open annulus $\{x \in \mathbb{R}^n : e^l < |x| < e^{l+1}\}$ is a polyharmonic function² of order p for $l \in \mathbb{Z}$. The reason for calling such polysplines "cardinal" is found in Theorem 3 where it is seen that after expanding S in a Fourier–Laplace series of spherical harmonics the coefficients $S_{k,l}(\log r)$ are cardinal L -splines in the usual sense of the word, cf. Michelli's paper of 1976 [8].

Introducing a parameter $h > 0$, by P_h we denote the set of all functions $S : \mathbb{R}^n \setminus \{0\} \rightarrow \mathbb{C}$ which are $(2p - 2)$ -times continuously differentiable and whose restriction to each open annulus $A_{h,l} := \{x \in \mathbb{R}^n : e^{hl} < |x| < e^{h(l+1)}\}$ is a polyharmonic function of order p for $l \in \mathbb{Z}$. Then the *scaling spaces of polysplines of order p* , shortly PV_h , are defined as the L^2 -closure of $P_h \cap L^2(\mathbb{R}^n)$, $h > 0$.

The main result is the following:

Theorem 1 *The sequence $(PV_h)_{h>0}$ provides approximation order $2p$ where p denotes the order of the polysplines. More precisely, there exists a constant*

¹The first author introduced in 1991 polysplines in a more general setting with arbitrary interfaces, see [3] and [9].

²Recall that a function f defined on an open set U in the euclidean space \mathbb{R}^n is *polyharmonic of order p* if f is $2p$ -times continuously differentiable and $\Delta^p f(x) = 0$ for all $x \in U$ where Δ is the Laplace operator and Δ^p its p -th iterate.

$C > 0$ such that for all h with $0 < h < 1$ and $f \in L^2(\mathbb{R}^n)$ the following inequality holds

$$\inf \left\{ \|f - g\|_{L^2(\mathbb{R}^n)} : g \in PV_h \right\} \leq C \cdot h^{2p} \cdot \left(\int_{\mathbb{R}^n} | |x|^{2p} \cdot \Delta^p f(x) |^2 dx \right)^{\frac{1}{2}}.$$

Note that in place of the norm (3) we have a semi-norm on the right-hand side which is zero on the polyharmonic functions of order p .

The paper is organized as follows: in the next Section we discuss the approximation order of cardinal L -splines by using important results from [2]. In Section 3 the main result will be proven.

2 Approximation order of cardinal L -splines

Let us recall Theorem 4.3 in the fundamental paper [2]: Suppose that for every $h > 0$, the space S_h is the $L^2(\mathbb{R}^n)$ -closure of the linear space generated by the shifts $\varphi_h(\cdot - m)$, $m \in \mathbb{Z}^n$ of the functions $\varphi_h \in L^2(\mathbb{R}^n)$ (so S_h is the shift-invariant space generated by φ_h) and that $V_h = \{s(\frac{\cdot}{h}) : s \in S_h\}$. Then the family $(V_h)_{h \in I}$ provides approximation order m with respect to the norm $\|\cdot\|_{W_2^m(\mathbb{R}^n)}$ defined in (3) if and only if there exists $D > 0$ such that for all $h \in I$ and for almost all $x \in C := [-\pi, \pi]^n$

$$|\Lambda_{\varphi_h}(x)| \leq D \cdot (h + |x|^m), \quad (4)$$

where

$$(\Lambda_{\varphi_h}(\xi))^2 := \frac{\sum_{\alpha \in \mathbb{Z}^n, \alpha \neq 0} |\widehat{\varphi}_h(\xi + 2\pi\alpha)|^2}{\sum_{\beta \in \mathbb{Z}^n} |\widehat{\varphi}_h(\xi + 2\pi\beta)|^2} \leq 1.$$

We will need a refinement of that result. For our purposes it will be useful to consider instead of (3) different norms. In the following we replace the function $(1 + |x|)^m$ by a measurable function $Q(x)$ with the following properties: (i) the zero set $Q^{-1}(0)$ of Q is a set of Lebesgue measure zero and (ii) there exists a constant $D_1 > 0$ such that

$$\left| Q\left(\frac{x}{h}\right) \right| \geq D_1 \frac{1}{h^m} \quad \text{for all } x \notin C := [-\pi, \pi]^n. \quad (5)$$

Suppose further that there exists a constant $D_2 > 0$ such that for all $x \in C$ and for all $0 < h < 1$

$$|\Lambda_{\varphi_h}(hx)| \leq h^m D_2 |Q(x)|. \quad (6)$$

An analysis of the proof in [2] shows that then the following inequality holds (for us the constants D_1 and D_2 defined in the formula will be very important!)

$$E(f, V_h) \leq \left(D_2 (2\pi)^{\frac{n}{2}} + \frac{1}{D_1 (2\pi)^{\frac{n}{2}}} \right) \cdot h^m \cdot \left\| Q(\xi) \widehat{f}(\xi) \right\|_{L_2(\mathbb{R}^n)}. \quad (7)$$

Let us recall some facts about L -splines: Let L be a linear differential operator with constant coefficients of order $N + 1$, say

$$L = M_\Lambda := \prod_{j=1}^{N+1} \left(\frac{d}{dv} - \lambda_j \right) \quad \text{where } \Lambda := (\lambda_1, \dots, \lambda_{N+1}). \quad (8)$$

Then a function $u : \mathbb{R} \rightarrow \mathbb{R}$ is called *cardinal L -spline on the mesh $h\mathbb{Z}$* ($h > 0$) if u is $(N - 1)$ -times continuously differentiable and if for every $l \in \mathbb{Z}$ there exists $f_l \in U_L := \{f \in C^\infty(\mathbb{R}) : Lf = 0\}$ such that $u(t) = f_l(t)$ for all $t \in (lh, (l + 1)h)$. The set of all cardinal L -splines for the operator $L = M_\Lambda$ on $h\mathbb{Z}$ will be denoted by $\mathcal{S}_{h\mathbb{Z}}(\Lambda)$. The scaling spaces $V_h(\Lambda)$ are defined by

$$V_h(\Lambda) = L^2(\mathbb{R}) \text{-closure of } \mathcal{S}_{h\mathbb{Z}}(\Lambda) \cap L^2(\mathbb{R}). \quad (9)$$

Let Q_Λ be the basic spline which can be defined by its Fourier transform by

$$\widehat{Q}_\Lambda(\xi) = \frac{\prod_{j=1}^{N+1} (e^{-\lambda_j} - e^{-i\xi})}{\prod_{j=1}^{N+1} (i\xi - \lambda_j)}. \quad (10)$$

Theorem 2 *Let $N \in \mathbb{N}$ be fixed. Then there exists a constant $D > 0$ such that for all $\Lambda = (\lambda_1, \dots, \lambda_{N+1}) \in \mathbb{R}^{N+1}$ and for all $f \in L_2(\mathbb{R})$ the following inequality holds*

$$E(f, V_h(\Lambda)) \leq h^{N+1} \cdot D \left\| P_\Lambda(\xi) \widehat{f}(\xi) \right\|_{L_2(\mathbb{R})}, \quad (11)$$

where the polynomial $P_\Lambda(x) = \prod_{j=1}^{N+1} (ix - \lambda_j)$.

Remark 3 *Note that if we used the usual Sobolev norm (3) then we could not be able to obtain the sharp constant D of inequality (11); the last is the main virtue of Theorem 2.*

Proof. By the above we have to check (5) and (6). Note that for $Q := P_\Lambda$ we have the estimate

$$\left| P_\Lambda \left(\frac{x}{h} \right) \right|^2 = \prod_{j=1}^{N+1} \left(\left(\frac{x}{h} \right)^2 + \lambda_j^2 \right) \geq \pi^{2(N+1)} \frac{1}{h^{2(N+1)}} \quad (12)$$

for all $|x| \geq \pi$ and for all $h > 0$. Hence it suffices to show that

$$|\Lambda_{\varphi_h}(h\xi)|^2 \leq h^{2(N+1)} |P_\Lambda(\xi)|^2 \sum_{\alpha \in \mathbb{Z}, \alpha \neq 0} \frac{1}{(\pi |\alpha|)^{2(N+1)}} \quad (13)$$

The trivial inequality $(\Lambda_{\varphi_h}(\xi))^2 \leq \frac{\sum_{\alpha \in \mathbb{Z}, \alpha \neq 0} |\widehat{\varphi_h}(\xi + 2\pi\alpha)|^2}{|\widehat{\varphi_h}(\xi)|^2}$ and the estimate

$$\frac{|\widehat{\varphi_h}(\xi + 2\pi\alpha)|^2}{|\widehat{\varphi_h}(\xi)|^2} = \frac{|\widehat{Q_{h\Lambda}}(\xi + 2\pi\alpha)|^2}{|\widehat{Q_{h\Lambda}}(\xi)|^2} = \prod_{j=1}^{N+1} \left| \frac{i\xi - h\lambda_j}{i(\xi + 2\pi\alpha) - h\lambda_j} \right|^2$$

yields

$$|\Lambda_{\varphi_h}(h\xi)|^2 \leq h^{2(N+1)} \prod_{j=1}^{N+1} (\xi^2 + \lambda_j^2) \sum_{\alpha \in \mathbb{Z}, \alpha \neq 0} \prod_{j=1}^{N+1} \frac{1}{(h\xi + 2\pi\alpha)^2 + h^2\lambda_j^2}.$$

Since $(h\xi + 2\pi\alpha)^2 + h^2\lambda_j^2 \geq (h\xi + 2\pi\alpha)^2 \geq (2\pi|\alpha| - |h\xi|)^2$ we obtain for $0 < h < 1$ and $|\xi| \leq \pi$ the estimate $2\pi|\alpha| - |h\xi| \geq \pi|\alpha|$ (since $\alpha \neq 0$) arriving at (13). ■

3 The approximation order of Polysplines

Let $\mathbb{S}^{n-1} = \{x \in \mathbb{R}^n; |x| = 1\}$ be the unit sphere. Each $x \in \mathbb{R}^n$ will be written in spherical coordinates $x = r\theta$ with $r \geq 0$ and $\theta \in \mathbb{S}^{n-1}$. Recall that a function $Y : \mathbb{S}^{n-1} \rightarrow \mathbb{C}$ is a *spherical harmonic* of degree $k \in \mathbb{N}_0$ if there exists a homogeneous harmonic polynomial $P(x)$ of degree k such that $P(\theta) = Y(\theta)$ for all $\theta \in \mathbb{S}^{n-1}$. The set \mathfrak{H}_k of all spherical harmonics of degree exactly k is a linear space of dimension $a_k := \dim \mathfrak{H}_k = \binom{n+k-1}{k} - \binom{n+k-3}{k-2}$. We denote by $Y_{k,l}$ with $l = 1, 2, \dots, a_k$ a base of \mathfrak{H}_k . For a detailed account we refer to *Stein-Weiss* [12].

Let $u : (R_1, R_2) \rightarrow \mathbb{C}$ be infinitely differentiable and $Y_k \in \mathfrak{H}_k$. Then it is well known that $\Delta(u(r)Y_k(\theta)) = Y_k(\theta)L_{(k)}u(r)$ where we have put

$$L_{(k)} = \frac{d^2}{dr^2} + \frac{n-1}{r} \frac{d}{dr} - \frac{k(k+n-2)}{r^2}. \quad (14)$$

By iteration we have $\Delta^p u = Y_k(\theta) \cdot [L_{(k)}]^p u(r)$. Let us put for convenience

$$\begin{aligned} \Lambda_+(k, p) &:= \{k, k+2, \dots, k+2p-2\}, \\ \Lambda_-(k, p) &:= \{-k-n+2, -k-n+4, \dots, -k-n+2p\}. \end{aligned}$$

The space of solutions of the equation $L_{(k)}^p f(r) = 0$ which are C^∞ for $r > 0$ is generated by a simple basis: for $j \in \Lambda_+(k, p) \cup \Lambda_-(k, p)$ the function r^j is clearly a solution, while for $j \in \Lambda_+(k, p) \cap \Lambda_-(k, p)$ we obtain a second solution $r^j \log r$. It will be convenient to make a transform of the variable r to $v = \log r$. Then a solution of the form r^j will be transformed to e^{jv} and a solution of the form $r^j \log r$ is transformed to ve^{jv} . We see immediately that all solutions to the equation $L_{(k)}^p f(r) = 0$ are transformed to solutions of the equation $M_{\Lambda(k)}g(v) = 0$ where $M_{\Lambda(k)}$ is defined by (8) with respect to the vector

$$\Lambda_k := (k, k+2, \dots, k+2(p-1), -(k+n)+2, \dots, -(k+n)+2p). \quad (15)$$

The dependence on the parameter p and n will be suppressed throughout the paper.

A proof of the following can be found in [6], [9, Theorem 9.7].

Theorem 4 *Let $S : \mathbb{R}^n \setminus \{0\} \rightarrow \mathbb{R}$ be a polyspline of order p . Then the Laplace-Fourier coefficient $S_{k,l} : \mathbb{R} \rightarrow \mathbb{R}$ defined by*

$$S_{k,l}(v) := \int_{\mathbb{S}^{n-1}} S(e^v \theta) Y_{k,l}(\theta) d\theta \quad (16)$$

is a cardinal L -spline with respect to the linear differential operator $M_{\Lambda(k)}$.

We want to characterize the $L^2(\mathbb{R}^n)$ -closure PV_h . It is a temptation to assume that for $S \in PV_h$ the Fourier-Laplace coefficient defined through formula (16) will be in $V_h(\Lambda_k)$, i.e. in the closure of $\mathcal{S}_{h\mathbb{Z}}(\Lambda_k) \cap L_2(\mathbb{R})$. This is *not true* since the transformation rule will give us an additional weight for $f \in L_2(\mathbb{R}^n)$:

$$\int_{\mathbb{R}^n} |f(x)|^2 dx = \int_0^\infty \int_{\mathbb{S}^{n-1}} |f(r\theta)|^2 r^{n-1} d\theta dr. \quad (17)$$

Fortunately, this problem can be easily solved, see e.g. [7].

Theorem 5 Define $\overline{\Lambda}_k = \left(\frac{n}{2}, \dots, \frac{n}{2}\right) + \Lambda_k$. Then for each $k \in \mathbb{N}_0, l = 1, \dots, a_k$, the following map, defined on $P_h \cap L^2(\mathbb{R}^n)$ by

$$S \mapsto \overline{S}_{k,l}(v) := e^{\frac{n}{2}v} \int_{\mathbb{S}^{n-1}} S(e^v \theta) Y_{k,l}(\theta) d\theta, \quad (18)$$

maps into $\mathcal{S}_{h\mathbb{Z}}(\overline{\Lambda}_k) \cap L^2(\mathbb{R}, dv)$, and by continuity it can be extended to a map from $PV_j \rightarrow V_j(\overline{\Lambda}_k)$. Further, PV_h is isomorphic to $V_h := \bigoplus_{k \in \mathbb{N}_0, l=1, \dots, a_k} V_h(\overline{\Lambda}_k)$.

Proof of Theorem 1. Let $f \in L^2(\mathbb{R}^n)$ and $g \in PV_h$. Then by the tranformation rule (17)

$$\|f - g\|_{L^2(\mathbb{R}^n)}^2 = \int_0^\infty \int_{\mathbb{S}^{n-1}} |f(r\theta) - g(r\theta)|^2 r^{n-1} d\theta dr. \quad (19)$$

Let $f_{k,l}$ and $g_{k,l}$ be the Laplace Fourier coefficients of f and g respectively as defined in (16). Note that $v \mapsto \overline{g_{k,l}}(e^v) := e^{\frac{n}{2}v} g_{k,l}(e^v)$ is in $V_h(\overline{\Lambda}_k)$. Since $Y_{k,l}(\theta)$ constitutes an orthonormal basis we obtain

$$\|f - g\|_{L^2(\mathbb{R}^n)}^2 = \sum_{k=0}^\infty \sum_{l=1}^{a_k} \int_{-\infty}^\infty |f_{k,l}(e^v) - g_{k,l}(e^v)|^2 e^{nv} dv. \quad (20)$$

Taking $g \in PV_h$ such that $\|f - g\|_{L^2(\mathbb{R}^n)}^2$ attains the ifimum is equivalent to take for each $k \in \mathbb{N}, l = 1, \dots, a_k$ the expression

$$\int_{-\infty}^\infty |e^{\frac{n}{2}v} f_{k,l}(e^v) - \overline{g_{k,l}}(e^v)|^2 dv$$

to be minimal, where $\overline{g_{k,l}} \in V_j(\overline{\Lambda}_k)$. Theorem 2 applied to $\Lambda = \overline{\Lambda}_k$ (hence $N + 1 = 2p$) shows that there exists a constant $C_p > 0$ which only depends on p (and not on the values λ_j in $\overline{\Lambda}_k$) such that

$$E(e^{\frac{n}{2}v} f_{k,l}(e^v), V_h(\overline{\Lambda}_k)) \leq h^{2p} \cdot C_p \left\| P_{\overline{\Lambda}_k} \cdot \widehat{e^{\frac{n}{2}v} f_{k,l}(e^v)} \right\|_{L_2(\mathbb{R})}. \quad (21)$$

Put $G_{k,l}(v) := e^{\frac{n}{2}v} f_{k,l}(e^v)$. A simple computation (using Parseval's identity and the fact that differentiation becomes multiplication via Fourier transform) shows that

$$\frac{1}{2\pi} \left\| P_{\overline{\Lambda}_k} \cdot \widehat{G_{k,l}} \right\|_{L_2(\mathbb{R})}^2 = \int_{-\infty}^\infty |M_{\overline{\Lambda}_k} G_{k,l}(v)|^2 dv.$$

A calculation shows that $M_{\Lambda_k}^{-1}(e^{\frac{n}{2}v} f_{k,l}(e^v)) = e^{\frac{n}{2}v} M_{\Lambda_k}(f(e^v))$. Then (20) and (21) yield

$$E(f, PV_h)^2 \leq h^{4p} \cdot 2\pi C_p^2 \sum_{k=0}^{\infty} \sum_{l=1}^{a_k} \left\| e^{\frac{n}{2}v} M_{\Lambda_k}(f(e^v)) \right\|_{L_2(\mathbb{R}^n)}^2.$$

The next theorem applied to the case $p = q$ finishes the proof. ■

Theorem 6 *Let $p, q \in N_0$ and define $\|f(x)\|_{q,p}^2 := \int | |x|^{2q} \cdot \Delta^p f(x) |^2 dx$ for $f \in L_2(\mathbb{R}^n)$. Then*

$$\|f(x)\|_{q,p}^2 = \sum_{k=0}^{\infty} \sum_{l=1}^{a_k} \int |e^{v(2q-2p+\frac{n}{2})} M_{\Lambda_k}(f_{k,l}(e^v))|^2 dv$$

where $f_{k,l}(r)$ are the Laplace-Fourier coefficients of f defined as in equality (16).

Proof. Assume that $f(r\theta) = f_{k,\ell}(r) Y_{k,\ell}(\theta)$. Since $\Delta^p f(x) = L_k^p f_{k,\ell}(r) Y_{k,\ell}(\theta)$ we obtain

$$\|f(x)\|_{q,p}^2 = \int_0^{\infty} \int_{\mathbb{S}^{n-1}} \left| r^{2q} L_{(k)}^p f_{k,\ell}(r) Y_{k,\ell}(\theta) \right|^2 r^{n-1} dr d\theta.$$

The integration over θ only gives a factor 1. Now we change the variable $r = e^v$ and apply the identity $(L_k^p f_{k,l})(e^v) = e^{-2vp} M_{\Lambda_k}(f_{k,l}(e^v))$, see e.g. Theorem 10.34 in [9]. Then

$$\|f(x)\|_{q,p}^2 = \int |e^{2vq} e^{-2vp} M_{\Lambda_k}(f_{k,l}(e^v))|^2 e^{nv} dv.$$

Finally we see that for arbitrary $f \in L^2(\mathbb{R}^n)$ the result follows via the orthogonal decomposition of f in spherical harmonics. ■

References

- [1] Adams R., *Sobolev Spaces*, Academic Press, New York–San Francisco–London, 1975.
- [2] de Boor, C., DeVore, R.A., Ron, A., *Approximation from shift-invariant subspaces of $L_2(\mathbb{R}^d)$* . Trans. Amer. Math. Soc. 341 (1994), pp. 787–806.

- [3] Kounchev, O.I., *Definition and basic properties of polysplines, I and II*, C.R. Acad. bulg. sci., 44 (1991), No. 7 and 8, pp. 9-11, pp.13-16.
- [4] Kounchev, O.I., Render, H., *Multivariate cardinal splines via spherical harmonics*. Submitted.
- [5] Kounchev, O.I., Render, H., *The interpolation problem for cardinal splines*. Submitted.
- [6] Kounchev, O.I., Render, H., *Symmetry of interpolation polysplines and L -splines*, Trends in Approximation Theory, K. Kopotun, T. Lyche, and M. Neamtu (eds.), Vanderbilt University Press, Nashville, TN, 2001.
- [7] Kounchev, O.I., Render, H., *Wavelet Analysis of cardinal L -splines and Construction of multivariate prewavelets*, Proceedings of Tenth Approximation Theory Conference, St. Louis, 2001, to appear.
- [8] Micchelli, Ch., *Cardinal L -splines*, In: Studies in Spline Functions and Approximation Theory, Eds. S. Karlin et al., Academic Press, NY, 1976, pp. 203-250.
- [9] Kounchev, O.I., *Multivariate Polysplines. Applications to Numerical and Wavelet Analysis*, Academic Press, Boston 2001.
- [10] Jetter, K., Plonka G., A survey on L_2 -approximation order from shift-invariant spaces, In: Multivariate Approximation and Applications (N. Dyn, D. Leviatan, D. Levin, and A. Pinkus, eds.), pp. 73-111. Cambridge University Press, 2001.
- [11] Meyer, Y. Wavelets and Operators, Cambridge University Press, 1992.
- [12] Stein, E.M., Weiss, G., *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton University Press, Princeton, 1971.

Address for correspondence:

1. Hermann Render

e-mail: render@math.uni-duisburg.de;

Institute of Mathematics, University of Duisburg, Lotharstr. 65, 47048
Duisburg, Germany

2. Ognyan Kounchev

e-mail addresses: kounchev@cblink.net; kounchev@math.uni-duisburg.de;
kounchev@math.bas.bg

Institute of Mathematics, Bulgarian Academy of Sciences, Acad. G.
Bonchev St. 8, 1113 Sofia, Bulgaria;