

1. COMPLEX NUMBERS

Notations: \mathbf{N} – the set of the natural numbers, \mathbf{Z} – the set of the integers, \mathbf{R} – the set of real numbers, $\mathbf{Q} :=$ the set of the rational numbers.

Given a quadratic equation

$$ax^2 + bx + c = 0,$$

we know that it is not always solvable; for example, the simple equation

$$x^2 = -1 \tag{1}$$

cannot be satisfied for any real number. But we can expand our number system \mathbf{R} by appending a symbol for a solution of (1); customary the symbol used is i , e.g.

$$i^2 = -1. \tag{2}$$

Definition: A complex number z is an expression of the form $z := a + ib$, where $a, b \in \mathbf{R}$. Two complex numbers $a+ib$ and $c+id$ are equal ($a+ib = c+id$) if and only if $a = c, b = d$. \aleph

1.1. The algebra of the complex numbers

Set \mathbf{C} for the set of complex numbers. Let $z_j = a_j + ib_j$. Following (2), we define

Addition by:

$$z_1 + z_2 := (a_1 + a_2) + i(b_1 + b_2);$$

Multiplication by;

$$z_1 z_2 := (a_1 + ib_1)(a_2 + ib_2) = (a_1 a_2 - b_1 b_2) + i(a_1 b_2 + a_2 b_1). \tag{3}$$

The Division of the complex numbers $\frac{z_1}{z_2}, z_2 \neq 0$ is given by

$$\frac{a_1 + ib_1}{a_2 + ib_2} := \frac{a_1 + ib_1}{a_2 + ib_2} \frac{a_2 - ib_2}{a_2 - ib_2} = \frac{a_1 a_2 + b_1 b_2 + i(a_2 b_1 - a_1 b_2)}{a_2^2 + b_2^2}.$$

We easily prove that addition and multiplication are *commutative* and *distributive*, as well as that the *Distributive Law* takes place, that is:

$$(z_1 + z_2)z_3 = z_1z_3 + z_2z_3.$$

Definition: The **real part** $\Re z$ of the complex number $z = a + ib$ is the (real) number a , its **imaginary part** $\Im z$ is the (real) number b . If a is zero, the number is said to be a **pure imaginary number**. \aleph .

1.2. Point representation of complex numbers, absolute value and complex conjugate.

A convenient way to represent complex numbers as points in the xy -plane is suggested by the *Cartesian coordinate system*; namely, to each complex number $z = a + ib$ we associate that point in the xy -plane which has the coordinates (a, b) (the projection of the $0x$ - axis is a , and the one on the Oy - axis is b). Obviously, the correspondence between the set of the complex numbers and the set of ordered pairs (x, y) is one-to-one.

When the xy -plane is used to describe complex numbers it is referred to as *complex plane* or **\mathbf{C} plane**. The Ox - axis is called the *real axis*, whereas the oy - axis - the *imaginary axis*.

Definition: The **absolute value** or the **modulus** of the number $z = a + ib$ is denoted by $|z|$ and is given by

$$|z| := \sqrt{a^2 + b^2} \tag{4}$$

\aleph .

Remark: $|z|$ is always nonnegative; $|z| = 0$ iff $\Re z = \Im z = 0$. The *distance* between $z_i = a_i + ib_i$, $i = 1, 2$ is given by

$$|z_1 - z_2| = \sqrt{(a_1 - a_2)^2 + (b_1 - b_2)^2}.$$

Definition: The complex conjugate of the number $z = a + ib$ is denoted by \bar{z} and is given by

$$\bar{z} := a - ib. \quad (5)$$

ℵ.

As we see, the complex conjugate of z is its reflection with respect to the real axis.

One easily can show that

$$\overline{(z_1 + z_2)} = \bar{z}_1 + \bar{z}_2,$$

$$\overline{(z_1 z_2)} = \bar{z}_1 \bar{z}_2,$$

$$\frac{\bar{z}_1}{z_2} = \frac{\bar{z}_1}{\bar{z}_2},$$

$$|z| = |\bar{z}|, |z|^2 = z\bar{z},$$

$$\Re z = \frac{z + \bar{z}}{2}, \Im z = \frac{z - \bar{z}}{2}.$$

1.3. Vectors and polar forms.

Definition: The Vector determined by the point z (the vector from the origin to the point z) in the complex plane \mathbf{C} will be called **the vector \mathbf{z}** . ℵ.

Addition: Let $z_i, i = 1, 2$ be two vectors, $z_i = \Re a_i + i b_i$. Hereafter, the sum of z_1 and z_2 is presented by the vector sum of both vectors, e.g., by the *parallelogram law*; $z_1 + z_2 = (a_1 + a_2, b_1 + b_2)$.

Subtraction: $z_1 - z_2 = (a_1 - a_2, b_1 - b_2)$.

Theorem 1.1 The Triangle Inequality. *For any two complex numbers z_1 and z_2 , the inequalities*

$$||z_1| - |z_2|| \leq |z_1 + z_2| \leq |z_1| + |z_2| \quad (6)$$

are valid.

There is another set of parameters that characterize the vector from the origin to the point z . This is the set of polar coordinates r — the modulus, and Θ — the argument of z . The coordinate r is the distance from the origin to

the point z ; $r := |z|$. *Theta* is an angle of inclination of the vector z measured positively in a counterclockwise sense from the positive real axis (and thus measured negative when clockwise). Let x, y be the Cartesian (rectangular) coordinates of z ; then

$$x = r \cos \Theta \quad (7)$$

and

$$y = r \sin \Theta \quad (8)$$

(recall that $r = |z|$.) Further,

$$\cos \Theta = \frac{x}{r}, \sin \Theta = \frac{y}{r}. \quad (9)$$

Remark: *Although it is certainly true that $\tan \Theta = x/y$, the natural conclusion*

$$\Theta = \arctan(y/x)$$

is not true in the second and the third quadrants.

On the "uniqueness" of Θ_0 . Let Θ satisfy (9). Then so does each

$$\Theta_0 + 2k\pi, k \in \mathbf{Z}.$$

We shall call the value of any of these angles an *argument* and denote it by $\arg z$. That value of the argument which belongs to the interval $(-\pi, \pi]$ will be called *the Principal Part of the argument* and denoted by $\text{Arg } z$.

For instance,

$$\begin{aligned} \arg 1 &= 0, & 2\pi, & -2\pi, \dots, \\ \arg i &= \frac{\pi}{2}, & \frac{5\pi}{2}, & \frac{-3\pi}{2}, \dots \\ \arg (1 - i) &= \frac{-\pi}{4}, & \frac{7\pi}{4}, & \frac{-9\pi}{4}, \dots \end{aligned}$$

With these convention in hand, one can now write $z = x + iy$ in a *polar form*

$$z = x + iy = r(\cos \Theta + i \sin \Theta). \quad (10)$$

Let now $z_i = r_i(\cos \Theta_i + i \sin \Theta_i), i = 1, 2$. Applying (3), we get

$$z_1 z_2 = r_1 r_2 (\cos(\Theta_1 + \Theta_2) + i \sin(\Theta_1 + \Theta_2)).$$

So we conclude that

$$|z_1 z_2| = |z_1| |z_2| \quad (11)$$

and

$$\arg(z_1 z_2) = \arg z_1 + \arg z_2. \quad (12)$$

Also,

$$\arg(1/z) = -\arg z \quad (13)$$

$$\arg \bar{z} = -\arg z. \quad (14)$$

1.4. The complex exponential - Euler's equation.

Definition: Euler's equation:

$$e^{iy} := \cos y + i \sin y.$$

We leave to the reader the proof that the function $e^{i\alpha}$ preserves all known characteristics of the exponential function e^x , x — a real variable, i.e. :

$$e^0 = 1, e^{x+y} = e^x e^y, (e^x)^y = e^{xy}.$$

The new definition enables us to rewrite (10) in a *trigonometric form* that is:

$$z = |z| e^{\text{argument} z}.$$

Further, we now may define the exponential function e^z , $z \in \mathbf{C}$:

Definition: if $z = x + iy$, then e^z is defined to be the complex number

$$e^z = e^{x+iy} = e^x (\cos y + i \sin y).$$

EXERCISES:

1. Write in the form $a + ib$.

a): $-3(i/2)$; $2/i$; $(-1 + i)^2$, $i^3(i + 1)^2$.

b) Show that $\Re(iz) = -\Im(z)$ for every complex number z .

2.

a) Let $z = 3 - 2i$. Plot the points z , $-z$, \bar{z} and $1/z$ in the complex plane.

- b) Describe the set of points z in \mathbf{C} that satisfy $\Im z = -2$, $|z - 2| \leq 1$, $\Re z > 2$, $|z| = \Re z - 2$.
- c) Prove that $|\Re z| \leq |z|$, $|\Im z| \leq |z|$.
- d) Let $a_i, i = 1, \dots, n$ are real numbers. Show that if z_0 is a root of the polynomial $z^n + a_1 z^{n-1} + \dots + a_n = 0$, then so is \bar{z}_0 .

3.

- a) Write down in a polar form $1, -1, i, -i, 1 \pm i, \frac{\pm 1 + i \pm \sqrt{3}}{2}, \frac{\pm 1 + i \pm 1}{2}$.
- b) Is it true or not: $\text{Arg}(z_1 z_2) = \text{Arg} z_1 + \text{Arg} z_2$; $\text{Arg} z = -\text{Arg}(-z)$, $\arg z = \text{Arg} z + 2k\pi, k \in \mathbf{Z}$

4. Prove Moivre's formula

$$(\cos \Theta + i \sin \Theta)^n = \cos n\Theta + i \sin n\Theta.$$

Show that $|e^{i\alpha}| = 1$, $e^{2k\pi i} = 1, k \in \mathbf{Z}$, $e^{\pi i/2} = i$, $e^{(2k+1)\pi/2} = -1$.

$$\sum_{k=0}^n z^k = \frac{z^{n+1} - 1}{z - 1}, z \neq 1.$$

5.

Write down in a trigonometric form the numbers

- a) $1 - \cos \alpha + i \sin \alpha$;
 b) $1 + \cos \alpha + i \sin \alpha$;
 c) $1 + \sin \alpha + i \cos \alpha$, with $\alpha \in [0, 2\pi]$.

5. Write down the sum $1 + 2 \sum \cos n\alpha$ as a product.

6. Given the complex number $z_i, i = 1, 2, 3, z_i \neq z_j, i, j = 1, 2, 3$ verify the equivalency

$$|z_1 - z_3|^2 = |z_1 - z_2|^2 + |z_2 - z_3|^2 \iff (z_3 - z_2) = \beta(z_2 - z_1),$$

β — a pure imaginary number.

7. Show that

a) the equation

$$|z|^2 + lz + \bar{l}\bar{z} + m = 0, m - \text{ a real number, } m < |l|^2$$

describes a circle.

b) the equation

$$Az + \bar{A}\bar{z} + B = 0, A \neq 0, B - \text{ a pure imaginary number, } m < |l|^2$$

describes a line.