

DISTINCTNESS OF THE "LIFTED" KLOOSTERMAN SUMS OVER THE PRIME FIELD \mathbb{F}_p

LYUBOMIR BORISSOV

ABSTRACT. In this talk I consider the Kloosterman sums over the finite field \mathbb{F}_q of characteristic p , defined by

$$\mathcal{K}_q(u) = \sum_{x \in \mathbb{F}_q^*} \omega^{Tr(x+ux^{-1})},$$

where $\omega = e^{\frac{2\pi i}{p}}$ is a primitive p -th root of unity, and $Tr(a)$ is the absolute trace of $a \in \mathbb{F}_q$ over \mathbb{F}_p .

The focus of special attention are the so-called "lifted" Kloosterman sums over \mathbb{F}_q (see, [2]), i.e., $\mathcal{K}_{q^n}(u)$, $u \in \mathbb{F}_q$, where \mathbb{F}_{q^n} is the finite field of order q^n , $n > 1$.

It is well-known that the Kloosterman sums play an important role in algebraic coding theory and cryptography (see, e.g., the surveys [3]-[4]).

Firstly I clashed with them in the problem of enumerating the elements of a finite field having prescribed trace and co-trace:

<https://arxiv.org/pdf/1711.08306.pdf>

The issue of their distinctness is considered and partly solved for the first time by Benjamin Fisher in 1992 [5]. In particular, this author has proved that fact for the simplest sums, i.e., over the prime fields.

Recently, in a personal communication with us, Daqing Wan has announced that as a co-product of his research [6] (based on deep algebraic number theory such as Stickelberger's theorem) it follows the distinctness of "lifted" Kloosterman sums over any prime field \mathbb{F}_p whenever the extension degree is not a multiple of p . This statement generalizes our result for the fields whose extension degree is a power of 2:

<https://link.springer.com/article/10.1007/s12095-020-00443-1>

The case $p = 3$ I considered in

Here I give a complete proof that all "lifted" Kloosterman sums over each prime field of characteristic $p > 3$ and any extension degree, are distinct. I gave a talk on international seminar <https://boolean.w.uib.no/bfa-2024/> about this and the result is still in process to be published.

REFERENCES

- [1] ON THE DISTINCTNESS OF SOME TERNARY KLOOSTERMAN SUMS Lyubomir Borissov Proceedings of the Fiftieth Spring Conference of the Union of Bulgarian Mathematicians, (2021)
- [2] L. Carlitz, "Kloosterman sums and finite field extensions", Acta Arithmetica vol. XVI.2 (1969), pp. 179-193.
- [3] N. E. Hurt, "Exponential sums and coding theory: a review", Acta Appl. Math., vol. 46.1 (1997), pp. 49-91.
- [4] V. A. Zinoviev, "On classical Kloosterman sums", Cryptogr. and Commun., 11.3 (2019), pp. 461-496.
- [5] B. Fischer, "Distinctness of Kloosterman sums", Contemporary Mathematics, vol. 133 (1992), pp. 81-102.
- [6] D. Wan, "Minimal polynomials and distinctness of Kloosterman sums", Finite Fields Appl., 1 (1995), pp. 189-203.

INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES, SOFIA,
BULGARIA