

Vector Schroedinger Equation. Generalized Wave Equation. Boussinesq Paradigm. Nonintegrability and Conservativity of Soliton Solutions

Michail D. Todorov

Faculty of Applied Mathematics and Informatics
Technical University of Sofia, Bulgaria

NEW SEMINAR OF THE FACULTY OF APPLIED MATHEMATICS
AND INFORMATICS
MARCH 15, 2012

- Problem Formulation. Integrability and Nonintegrability
- Conservation Laws and Boussinesq Paradigm
- Choice or Generation of Initial Conditions
- Quasi-Particle Dynamics and Polarization
- Numerics
- Some Results and Discussion

Problem Formulation: Equations

CNLSE is system of nonlinearly coupled Schrödinger equations (called the Gross-Pitaevskii or Manakov-type system):

$$\begin{aligned}i\psi_t &= \beta\psi_{xx} + [\alpha_1|\psi|^2 + (\alpha_1 + 2\alpha_2)|\phi|^2]\psi(+\Gamma\phi), \\i\phi_t &= \beta\phi_{xx} + [\alpha_1|\phi|^2 + (\alpha_1 + 2\alpha_2)|\psi|^2]\phi(+\Gamma\psi),\end{aligned}\tag{1}$$

where:

β is the dispersion coefficient;

α_1 describes the self-focusing of a signal for pulses in birefringent media;

$\Gamma = \Gamma_r + i\Gamma_i$ is the magnitude of linear coupling. Γ_r governs the oscillations between states termed as breathing solitons, while Γ_i describes the gain behavior of soliton solutions.

α_2 (called cross-modulation parameter) governs the nonlinear coupling between the equations.

Problem Formulation: Equations

When $\alpha_2 = 0$, no nonlinear coupling is present despite the fact that “cross-terms” proportional to α_1 appear in the equations. For $\alpha_2 = 0$, the solutions of the two equations are identical, $\psi \equiv \phi$, and equal to the solution of single NLSE with nonlinearity coefficient $\alpha = 2\alpha_1$.

For $\Gamma = 0$, CNSE is alternately called the Gross-Pitaevskii equation or an equation of Manakov type. It was derived independently by Gross and Pitaevskii to describe the behavior of Bose-Einstein condensates as well as optic pulse propagation. It was solved analytically for the case $\alpha_2 = 0$, $\beta = \frac{1}{2}$ by Manakov via Inverse Scattering Transform who generalized an earlier result by Zakharov & Shabat for the scalar cubic NLSE (i.e. Eq.(2- ψ) with $\varphi(x, t) = 0$). Recently, Chow, Nakkeeran, and Malomed studied periodic waves in optic fibers using a version of CNSE with $\Gamma \neq 0$.

The single NLSE has the form

$$i\psi_t + \beta\psi_{xx} + \alpha|\psi|^2\psi = 0 \quad (2)$$

As far as the applications in nonlinear optics are concerned, the above equation describes the single-mode wave propagation in a fiber. Depending on the sign of (2) admits single and multiple sech-solutions (*bright solitons*), as well as tanh-profile, or *dark soliton* solutions. In this paper we concentrate on the case of bright solitons.

A dynamical system with infinite number of conservation laws (integrals) is called integrable. NSE is an integrable dynamical system. CNSE is a nonintegrable generalisation of NSE.

Problem Formulation: Conservation Laws

Define “mass”, M , (pseudo)momentum, P , and energy, E :

$$M \stackrel{\text{def}}{=} \frac{1}{2\beta} \int_{-L_1}^{L_2} (|\psi|^2 + |\phi|^2) dx, \quad P \stackrel{\text{def}}{=} - \int_{-L_1}^{L_2} \mathcal{I}(\psi \bar{\psi}_x + \phi \bar{\phi}_x) dx, \\ E \stackrel{\text{def}}{=} \int_{-L_1}^{L_2} \mathcal{H} dx, \quad \text{where} \quad (3)$$

$$\mathcal{H} \stackrel{\text{def}}{=} \beta (|\psi_x|^2 + |\phi_x|^2) - \frac{1}{2} \alpha_1 (|\psi|^4 + |\phi|^4) \\ - (\alpha_1 + 2\alpha_2) (|\phi|^2 |\psi|^2) - 2\Gamma [\Re(\bar{\psi}\bar{\phi})]$$

is the Hamiltonian density of the system. Here $-L_1$ and L_2 are the left end and the right end of the interval under consideration.

The following conservation/balance laws hold, namely

$$\frac{dM}{dt} = 0, \quad \frac{dP}{dt} = \mathcal{H}|_{x=L_2} - \mathcal{H}|_{x=-L_1}, \quad \frac{dE}{dt} = 0, \quad (4)$$

Boussinesq Paradigm. Quasi-particles Concept

Boussinesq 's equation (BE) was the first model for the propagation of surface waves over shallow inviscid layer which contains dispersion. Boussinesq found an analytical solution of his equation and thus proved that the balance between the steepening effect of the nonlinearity and the flattening effect of the dispersion maintains the shape of the wave. This discovery can be properly termed 'Boussinesq Paradigm'. Apart from the significance for the shallow water flows, this paradigm is very important for understanding the particle-like behavior of localized waves which behavior was discovered in the 1960ies (the so-called 'collision property'), and the localized waves were called *solitons* (Zabusky & Kruskal). The localized waves which can retain their identity during interaction appear to be a rather pertinent model for particles, especially if some mechanical properties are conserved by the governing equations (such as mass, energy, momentum). For this reason they are called quasi-particles.

Boussinesq Paradigm. Quasi-particles Concept

The equation derived by Boussinesq (1871) referred as “Original Boussinesq Equation”

$$\frac{\partial^2 h}{\partial t^2} = \frac{\partial^2}{\partial x^2} \left[(gH)h + \frac{3g}{2}h^2 + \frac{gH^3}{3} \frac{\partial^2 h}{\partial x^2} \right] \quad (5)$$

is fully integrable but incorrect in the sense of Hadamard due to the positive sign of the fourth spatial derivative.

Including the effect of surface tension as was done by Korteweg & de Vries one arrives so called Proper Boussinesq Equation

$$\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2}{\partial x^2} \left[u - \alpha u^2 + \beta \frac{\partial^2 u}{\partial x^2} \right] \quad (6)$$

which is correct in the sense of Hadamard (well-posed as IVP) only when $\beta < 0$.

Boussinesq Paradigm. Quasi-particles Concept

The ill-posedness of OBE can be removed if the Boussinesq assumption that $\frac{\partial}{\partial t} \approx \frac{\partial}{\partial x}$ is used in “reverse”:

$$\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2}{\partial x^2} \left[u - \alpha u^2 + \beta \frac{\partial^2 u}{\partial t^2} \right]. \quad (7)$$

This equation is called “improper” or RLWE. All versions of the Boussinesq equations considered here possess solitary-wave solutions of *sech* type. For example, for the BE the solution reads

$$u = -\frac{3}{2} \frac{c^2 - 1}{\alpha} \operatorname{sech}^2 \left(\frac{x - ct}{2} \sqrt{\frac{c^2 - 1}{\beta}} \right)$$

Boussinesq Paradigm. Quasi-particles Concept

The overwhelming majority of the analytical and numerical results so far are for one spatial dimension, while in multidimension much less is possible to achieve analytically, and almost nothing is known about the unsteady solutions, especially when the Boussinesq equations contain different dispersions and nonlinearities are involved. An exception is the case of so called KPE, which has fourth derivative only in one spatial directions, while in the other direction, the highest order is second.

Boussinesq Paradigm Equation (BPE)

$$u_{tt} = \Delta[u - \alpha u^2 + \beta_1 u_{tt} - \beta_2 \Delta u] \quad (8)$$

where u is the surface elevation, $\beta_1, \beta_2 > 0$ are two dispersion coefficients, α is an amplitude parameter.

Boussinesq Paradigm. Quasi-particles Concept

Cubic-Quintic Boussinesq Paradigm Equation (CQBPE)

$$u_{tt} = \Delta[u - \alpha(u^3 - \sigma u^5) + \beta_1 u_{tt} - \beta_2 \Delta u] \quad (9)$$

where u again is the surface elevation, $\beta_1, \beta_1 > 0$ are two dispersion coefficients. The parameter σ accounts for the relative importance of the quintic nonlinearity term. Then energy law reads

$$\frac{dE}{dt} = 0 \quad \text{with}$$

$$E = \frac{1}{2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} [(\nabla u_t)^2 + u^2 - \frac{1}{2} u^4 + \frac{\sigma}{3} u^6 + \beta_1 u^2 + \beta_2 (\nabla u)^2] dx dy.$$

Unlike the BPE with quadratic nonlinearity, when the amplitude increases, the quintic term in CQBPE for reasonably large σ will dominate and will make the energy functional positive, which limits the increase of the amplitude. All this means that no blow-up can be expected for CQBPE when $\sigma > \frac{3}{16}$.

Boussinesq Paradigm. (Energy)-Consistent Boussinesq System

This a further generalization of the Boussinesq Paradigm equation. The following system was derived rigorously from the main Boussinesq assumption

$$\frac{\partial \chi}{\partial t} = -\beta \nabla \cdot \chi \nabla f - \Delta f + \frac{\beta}{6} \Delta^2 f - \frac{\beta}{2} \frac{\partial^2 \Delta f}{\partial t^2}, \quad (10a)$$

$$\frac{\partial f}{\partial t} = -\frac{\beta}{2} (\nabla f)^2 - \chi, \quad (10b)$$

Boussinesq Paradigm. (Energy)-Consistent Boussinesq System

Upon multiplying the left-hand side of (10a) by the right-hand side of Eq. (10b) and the right-hand side of Eq. (10a) by the left-hand side of Eq. (10b), adding the results and integrating over the surface region D under consideration one gets the following energy balance law

$$E = \frac{1}{2} \int_D \left[\chi^2 + (1 + \beta\chi)(\nabla f)^2 + \frac{\beta}{6}(\Delta f)^2 + \frac{\beta}{2}(\nabla f_t)^2 \right] dx$$
$$\frac{dE}{dt} = \oint_{\partial D} \left[(1 + \beta\chi)f_t \frac{\partial f}{\partial n} + \frac{\beta}{2}f_t \frac{\partial f_{tt}}{\partial n} + \frac{\beta}{2}f_t \frac{\partial \Delta f}{\partial n} - \frac{\beta}{2}\Delta f \frac{\partial f_t}{\partial n} \right] ds, \quad (11)$$

which allows us to call the Eqs. (10) 'Energy Consistent Boussinesq Paradigm.'

Problem Formulation: Choice of Initial Conditions

We concern ourselves with the soliton solutions whose modulation amplitude is of general form (non-sech) and which are localized envelopes on a propagating carrier wave. This allows us to play various scenario of initial polarization. Unfortunately in sech-case the initial polarization can be only linear. Then we assume that for each of the functions ϕ, ψ the initial condition is of the form of a single propagating soliton, namely

$$\begin{aligned} \begin{Bmatrix} \psi(x, t) \\ \phi(x, t) \end{Bmatrix} &= \begin{Bmatrix} A^\psi \\ A^\phi \end{Bmatrix} \operatorname{sech} [b(x - X - ct)] \exp \left\{ i \left[\frac{c}{2\beta} (x - X) - nt \right] \right\}. \\ b^2 &= \frac{1}{\beta} \left(n + \frac{c^2}{4\beta} \right), \quad A = b \sqrt{\frac{2\beta}{\alpha_1}}, \quad u_c = \frac{2n\beta}{c}, \end{aligned} \quad (12)$$

where X is the spatial position (center of soliton), c is the phase speed, n is the carrier frequency, and b^{-1} – a measure of the support of the localized wave.

Problem Formulation: Choice of Initial Conditions

We assume that for each of the functions ϕ, ψ the initial condition has the general type

$$\begin{aligned}\psi &= A_\psi(x + X - c_\psi t) \exp \left\{ i \left[n_\psi t - \frac{1}{2} c_\psi (x - X - c_\psi t) + \delta_\psi \right] \right\} \\ \phi &= A_\phi(x + X - c_\phi t) \exp \left\{ i \left[n_\phi t - \frac{1}{2} c_\phi (x - X - c_\phi t) + \delta_\phi \right] \right\},\end{aligned}\quad (13)$$

where c_ψ, c_ϕ are the phase speeds and X 's are the initial positions of the centers of the solitons; n_ψ, n_ϕ are the carrier frequencies for the two components; δ_ψ and δ_ϕ are the phases of the two components. Note that the phase speed must be the same for the two components ψ and ϕ . If they propagate with different phase speeds, after some time the two components will be in two different positions in space, and will no longer form a single structure. For the envelopes (A_ψ, A_ϕ) , $\theta \equiv \arctan(\max |\phi| / \max |\psi|)$ is a polarization angle.

Problem Formulation: Generation of Initial Conditions

Generally the carrier frequencies for the two components $n_\psi \neq n_\phi$ – elliptic polarization. When $n_\psi = n_\phi$ – circular polarization. If one of them vanishes – linear polarization (sech soliton, $\theta = 0; 90^\circ$). In general case the initial condition is solution of the following system of nonlinear conjugated equations

$$\begin{aligned} A''_\psi + \left(n_\psi + \frac{1}{4}c_\psi^2\right) A_\psi + [\alpha_1 A_\psi^2 + (\alpha_1 + 2\alpha_2)A_\phi^2] A_\psi &= 0 \\ A''_\phi + \left(n_\phi + \frac{1}{4}c_\phi^2\right) A_\phi + [\alpha_1 A_\phi^2 + (\alpha_1 + 2\alpha_2)A_\psi^2] A_\phi &= 0. \end{aligned} \quad (14)$$

The system admits bifurcation solutions since the trivial solution obviously is always present.

Initial Conditions and Initial Polarization

We solve the auxiliary conjugated system (14) with asymptotic boundary conditions using Newton method and the initial approximation of sought nontrivial solution is sech-function. The final solution, however, is not obligatory sech-function. It is a two-component polarized soliton solution.




Figure: 1. Amplitudes A_ψ and A_ϕ for $c_l = -c_r = 1$, $\alpha_1 = 0.75$, $\alpha_2 = 0.2$. Left: $n_\psi = -0.68$; middle: $n_\psi = -0.55$; right: $n_\psi = -0.395$.

Initial Conditions and Initial Phase Difference

Another dimension of complexity is introduced by the phases of the different components. The initial difference in phases can have a profound influence on the polarizations of the solitons after the interaction and the magnitude of the full energy. The relative shift of real and imaginary parts is what matters in this case.




Figure: 2. Real and imaginary parts of the amplitudes from the case shown in the middle panel of Figure 1 and the dependence on phase angle.

Problem Formulation: Initial Conditions

After completing the initial conditions our aim is to understand better the influence of the initial polarization and initial phase difference on the particle-like behavior of the localized waves. We call a localized wave a quasi-particle (QP) if it survives the collision with other QPs (or some other kind of interactions) without losing its identity.

Linearly Coupled Problem Formulation: Equations and Initial Conditions

For the linearly coupled system of NLSE the magnitude of linear coupling Γ_r generates breathing the solitons although noninteracting The initial conditions must be

$$\Psi = \psi \cos(\Gamma t) + i\phi \sin(\Gamma t), \quad \Phi = \phi \cos(\Gamma t) + i\psi \sin(\Gamma t), \quad (15)$$

where ϕ and ψ are assumed to be sech-solutions of (1) for $\alpha_2 = 0$. Hence (1) posses solutions, which are combinations of interacting solitons oscillating with frequency Γ_r and their motion gives rise to the so-called rotational polarization.

Problem Formulation: Conservation Laws

In all considered cases we found that a conservation of the total polarization is present. Only for the linearly CNLSE ($\alpha_2 = 0$) the total polarizations breathe with an amplitude evidently depending on the initial phase difference but is conserved within one full period of the breathing.

$\delta_r - \delta_l$	θ_l^i	θ_r^i	$\theta_l^i + \theta_r^i$	θ_l^f	θ_r^f	$\theta_l^f + \theta_r^f$
45°	45°	45°	90°	33°48'	56°12'	90°
90°	45°	45°	90°	24°06'	65°54'	90°
0°	20°	20°	40°	20°00'	20°00'	40°
90°	20°	20°	40°	28°48'	2°02'	30°50'
0°	36°	36°	72°	36°00'	36°00'	72°
90°	36°	36°	72°	53°00'	13°20'	66°20'
0°	10°	80°	90°	21°05'	68°54'	89°59'
90°	10°	80°	90°	9°27'	80°30'	89°57'

To solve the main problem numerically, we use an implicit conservative scheme in complex arithmetic.

$$\begin{aligned} i \frac{\psi_i^{n+1} - \psi_i^n}{\tau} &= \frac{\beta}{2h^2} (\psi_{i-1}^{n+1} - 2\psi_i^{n+1} + \psi_{i+1}^{n+1} + \psi_{i-1}^n - 2\psi_i^n + \psi_{i+1}^n) \\ &+ \frac{\psi_i^{n+1} + \psi_i^n}{4} \left[\alpha_1 (|\psi_i^{n+1}|^2 + |\psi_i^n|^2) + (\alpha_1 + 2\alpha_2) (|\phi_i^{n+1}|^2 + |\phi_i^n|^2) \right] \\ &- \frac{1}{2} \Gamma (\phi_i^{n+1} + \phi_i^n), \end{aligned}$$

$$\begin{aligned} i \frac{\phi_i^{n+1} - \phi_i^n}{\tau} &= \frac{\beta}{2h^2} (\phi_{i-1}^{n+1} - 2\phi_i^{n+1} + \phi_{i+1}^{n+1} + \phi_{i-1}^n - 2\phi_i^n + \phi_{i+1}^n) \\ &+ \frac{\phi_i^{n+1} + \phi_i^n}{4} \left[\alpha_1 (|\phi_i^{n+1}|^2 + |\phi_i^n|^2) + (\alpha_1 + 2\alpha_2) (|\psi_i^{n+1}|^2 + |\psi_i^n|^2) \right] \\ &- \frac{1}{2} \Gamma (\psi_i^{n+1} + \psi_i^n). \end{aligned}$$

$$\begin{aligned}
 i \frac{\psi_i^{n+1,k+1} - \psi_i^n}{\tau} &= \frac{\beta}{2h^2} \left(\psi_{i-1}^{n+1,k+1} - 2\psi_i^{n+1,k+1} + \psi_{i+1}^{n+1,k+1} \right. \\
 &\quad \left. + \psi_{i-1}^n - 2\psi_i^n + \psi_{i+1}^n \right) \\
 &+ \frac{\psi_i^{n+1,k} + \psi_i^n}{4} \left[\alpha_1 (|\psi_i^{n+1,k+1}| |\psi_i^{n+1,k}| + |\psi_i^n|^2) \right. \\
 &\quad \left. + (\alpha_1 + 2\alpha_2) (|\phi_i^{n+1,k+1}| |\phi_i^{n+1,k}| + |\phi_i^n|^2) \right] \\
 i \frac{\phi_i^{n+1,k+1} - \phi_i^n}{\tau} &= \frac{\beta}{2h^2} \left(\phi_{i-1}^{n+1,k+1} - 2\phi_i^{n+1,k+1} + \phi_{i+1}^{n+1,k+1} \right. \\
 &\quad \left. + \phi_{i-1}^n - 2\phi_i^n + \phi_{i+1}^n \right) \\
 &+ \frac{\phi_i^{n+1,k} + \phi_i^n}{4} \left[\alpha_1 (|\phi_i^{n+1,k+1}| |\phi_i^{n+1,k}| + |\phi_i^n|^2) \right. \\
 &\quad \left. + (\alpha_1 + 2\alpha_2) (|\psi_i^{n+1,k+1}| |\psi_i^{n+1,k}| + |\psi_i^n|^2) \right].
 \end{aligned}$$

Numerical Method: Conservation Properties

It is not only convergent (consistent and stable), but also conserves mass and energy, i.e., there exist discrete analogs for (4), which arise from the scheme.

$$\begin{aligned} M^n &= \sum_{i=2}^{N-1} (|\psi_i^n|^2 + |\phi_i^n|^2) = \text{const}, \\ E^n &= \sum_{i=2}^{N-1} \frac{-\beta}{2h^2} (|\psi_{i+1}^n - \psi_i^n|^2 + |\phi_{i+1}^n - \phi_i^n|^2) + \frac{\alpha_1}{4} (|\psi_i^n|^4 + |\phi_i^n|^4) \\ &\quad + \frac{1}{2}(\alpha_1 + 2\alpha_2) (|\psi_i^n|^2 |\phi_i^n|^2) - \Gamma \Re[\bar{\phi}_i^n \psi_i^n] = \text{const}, \\ &\quad \text{for all } n \geq 0. \end{aligned}$$

These values are kept constant during the time stepping. The above scheme is of Crank-Nicolson type for the linear terms and we employ internal iterations to achieve implicit approximation of the nonlinear terms, i.e., we use its linearized implementation.

Results and Discussion: Initial Circular Polarizations of 45° , $\alpha_2 = 0$




Figure: 3. $\delta_l = 0^\circ$, $\delta_r = 0^\circ$

Results and Discussion: Initial Circular Polarizations of 45° , $\alpha_2 = 0$




Figure: 4. $\delta_l = 0^\circ$, $\delta_r = 45^\circ$

Results and Discussion: Initial Circular Polarizations of 45° , $\alpha_2 = 0$

- When both of QPs have zero phases (Fig. 3), the interaction perfectly follows the analytical Manakov two-soliton solution.
- The surprise comes in Fig. 4 where is presented an interaction of two QPs, the right one of which has a nonzero phase $\delta_r = 45^\circ$. After the interaction, the two QPs become different Manakov solitons than the original two that entered the collision. The outgoing QPs have polarizations $33^\circ 48'$ and $56^\circ 12'$. Something that can be called a ‘shock in polarization’ takes place. All the solutions are perfectly smooth, but because the property called polarization cannot be defined in the cross-section of interaction and for this reason, it appears as undergoing a shock.

Results and Discussion: Initial Circular Polarizations of 45° , $\alpha_2 = 0$

Here is to be mentioned that when rescaled the moduli of ψ and ϕ from Fig. 4 perfectly match each other which means that the resulting solitons have circular polarization (see left panel of Fig. 5 below). The Manakov solution is not unique. There exists a class of Manakov solution and in the place of interaction becomes a bifurcation between them.

Results and Discussion: Initial Circular Polarization and Nonuniqueness of the Manakov Solution




Figure: 5. Circular polarization (left); Elliptic Polarization (right).

Equal Elliptic Initial Polarizations of $50^{\circ}08'$ for $\alpha_2 = 2$




Figure: 6. $\delta_l = 0^\circ, \delta_r = 0^\circ$




Figure: 7. $\delta_l = 0^\circ, \delta_r = 180^\circ$

Equal Elliptic Initial Polarizations of $50^{\circ}08'$ for $\alpha_2 = 2$




Figure: 8. $\delta_l = 0^{\circ}, \delta_r = 130^{\circ}$




Figure: 9. $\delta_l = 0^{\circ}, \delta_r = 135^{\circ}$




Figure: 10. $\delta_l = 0^{\circ}, \delta_r = 140^{\circ}$

Equal Elliptic Initial Polarizations of $50^{\circ}08'$ for $\alpha_2 = 2$.

We choose $n_{l\psi} = n_{r\psi} = -1.5$, $n_{l\phi} = n_{r\phi} = -1.1$, $c_l = -c_r = 1$, $\alpha_1 = 0.75$, and focus on the effects of α_2 and $\vec{\delta}$.

One sees that the desynchronisations of the phases leads in the final stage to a superposition of two one-soliton solutions but with different polarizations from the initial polarization. Yet, for $\delta_r = 130^{\circ} \div 140^{\circ}$ one of the QPs loses its energy contributing it to the other QP during the collision and then virtually disappears: kind of energy trapping (Figs.8, 9, 10).

For $\delta_r = 180^{\circ}$ another interesting effect is seen, when the right outgoing QP is circularly polarized (Fig. 7).

All these interactions are accompanied by changes of phase speeds. The total polarization exhibits some kind of conservation.

Strong Nonlinear Interaction: $\alpha_2 = 10$




Figure: 11. $\alpha_2 = 10$, $c_l = 1$, $c_r = -0.5$.

Strong Nonlinear Interaction: $\alpha_2 = 10$

- Two new solitons are born after the collision.
- The kinetic energies of the newly created solitons correspond their phase speeds and masses, but the internal energy is very different for the different QP.
- the total energy of the QPs is radically different from the total energy of the initial wave profile. The differences are so drastic that the sum of QPs energies can even become negative. This means that the energy was carried away by the radiation.
- The predominant part of the energy is concentrated in the left and right forerunners because of the kinetic energies of the latter are very large. This is due to the fact that the forerunners propagate with very large phase speeds, and span large portions of the region.
- All four QPs have elliptic polarizations.
- Energy transformation is a specific trait of the coupled system considered here.

Linear Coupling: Initial Elliptic Polarizations: $\theta = 23^\circ 44'$,
 $\alpha_2 = 0$, $c_l = -c_r = 1$, $n_{l\psi} = n_{r\psi} = -1.1$,
 $n_{l\phi} = n_{r\phi} = -1.5$, $\Gamma = 0.175$




Figure: 12. $\delta_l = 0^\circ$, $\delta_r = 0^\circ$




Figure: 13. $\delta_l = 0^\circ$, $\delta_r = 90^\circ$

Linear Coupling: Initial Elliptic Polarizations: $\theta = 23^\circ 44'$,
 $\alpha_2 = 0$, $c_l = -c_r = 1$, $n_{l\psi} = n_{r\psi} = -1.1$,
 $n_{l\phi} = n_{r\phi} = -1.5$, $\Gamma = 0.175$




Figure: 14. Influence of the initial phase difference on the total energy:

$$\begin{aligned} \delta_l = 0^\circ, \delta_r = 0^\circ - E &= -0.262; \\ \delta_l = 0^\circ, \delta_r = 90^\circ - E &= -0.821; \\ \delta_l = 0^\circ, \delta_r = 135^\circ - E &= -0.206; \\ \delta_l = 0^\circ, \delta_r = 180^\circ - E &= 0.640 \end{aligned}$$

Linear Coupling: Initial Elliptic Polarizations: $\theta = 23^\circ 44'$,
 $\alpha_2 = 0$, $c_l = -c_r = 1$, $n_{l\psi} = n_{r\psi} = -1.1$,
 $n_{l\phi} = n_{r\phi} = -1.5$, $\Gamma = 0.175$




Figure: 15. $\delta_l = 0^\circ$, $\delta_r = 90^\circ$, $P = 10^{-3} \div 10^{-5}$

Linear Coupling: Initial Elliptic Polarizations: $\theta = 23^\circ 44'$,
 $\alpha_2 = 0$, $c_l = -c_r = 1$, $n_{l\psi} = n_{r\psi} = -1.1$,
 $n_{l\phi} = n_{r\phi} = -1.5$, $\Gamma = 0.175$




Figure: 16. $\delta_l = 0^\circ$, $\delta_r = 0^\circ$




Figure: 17. $\delta_l = 0^\circ$, $\delta_r = 90^\circ$

Linear Coupling: Initial Linear Polarizations: $\theta_l = 0^\circ$,
 $\theta_r = 90^\circ$, $\alpha_2 = 0$, $c_l = 1.5$, $c_r = 0.6$, $\Gamma = 0.175 + 0.005i$




Figure: 18. $\delta_l = 0^\circ$, $\delta_r = 90^\circ$

Linear Coupling: Initial Elliptic Polarizations: $\theta = 23^\circ 44'$,
 $\alpha_2 = 0$, $c_l = -c_r = 1$, $n_{l\psi} = n_{r\psi} = -1.1$,
 $n_{l\phi} = n_{r\phi} = -1.5$, $\Gamma = 0.175$

- We have found that the phases of the components play an essential role on the full energy of QPs. The magnitude of the latter essentially depends on the choice of initial phase difference (Figure 14);
- The pseudomomentum is also conserved and it is trivial due to the symmetry (Figure 15);
- The individual masses, however, breathe together with the individual (rotational) polarizations. Their amplitude and period do not influenced from the initial phase difference (Figure 15);

Linear Coupling: Initial Elliptic Polarizations: $\theta = 23^\circ 44'$,
 $\alpha_2 = 0$, $c_l = -c_r = 1$, $n_{l\psi} = n_{r\psi} = -1.1$,
 $n_{l\phi} = n_{r\phi} = -1.5$, $\Gamma = 0.175$

- The total mass is constant while the total polarization oscillates and suffers a 'shock in polarization' when QPs enter the collision. The polarization amplitude evidently depends on the initial phase difference (Figures 16,17);
- Due to the real linear coupling the polarization angle of QPs can change independently of the collision.
- Complex parameter of linear coupling: Along with the oscillations of the energy and masses the (negative) energy decreases very fast, while the masses M_ψ and M_ϕ increase all of them oscillating. The pseudomomentum P increases without appreciable oscillation (Figure 18).

C. I. Christov, S. Dost, and G. A. Maugin. Inelasticity of soliton collisions in system of coupled NLS equations. *Physica Scripta*, 50:449–454, 1994.

S. V. Manakov. On the theory of two-dimensional stationary self-focusing of electromagnetic waves. *Soviet Physics JETP*, 38:248–253, 1974.

W. J. Sonnier and C. I. Christov. Strong coupling of Schrödinger equations: Conservative scheme approach. *Mathematics and Computers in Simulation*, 69:514–525, 2005.

C. Sophocleous and D. F. Parker. Pulse collisions and polarization conversion for optical fibers. *Opt. Commun.*, 112:214–224, 1994.

T. R. Taha and M. J. Ablowitz. Analytical and numerical aspects of certain nonlinear evolution equations. ii. numerical, Schrödinger equation. *J. Comp. Phys.*, 55:203–230, 1984.

M. D. Todorov and C. I. Christov. Conservative scheme in complex arithmetic for coupled nonlinear Schrödinger equations. *Discrete and Continuous Dynamical Systems*”, Supplement:982–992, 2007

M. D. Todorov and C. I. Christov. Collision Dynamics of Elliptically Polarized Solitons in Coupled Nonlinear Schrodinger Equations.

Mathematics and Computers in Simulation, doi:

10.1016/j.matcom.2010.04.022, 2010.

M. D. Todorov and C. I. Christov. Collision Dynamics of Polarized Solitons in Linearly Coupled Nonlinear Schrödinger Equations, in K.Sekigawa et al (Eds.), *The International Workshop on Complex Structures, Integrability, and Vector Fields, 2010, Sofia, Bulgaria*, American Institute of Physics CP1340, pages 144–153, 2011.

V. E. Zakharov and A. B. Shabat. Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in nonlinear media. *Soviet Physics JETP*, 34:62–9, 1972: 37:823, 1973.

C. I. Christov, M. D. Todorov, and M. A. Christou, Perturbation Solution for the 2D Shallow-water Waves. in *Application of Mathematics in Technical and Natural Sciences, 2011, Albena, Bulgaria*, American Institute of Physics CP1404, pages 49–56, 2012.

Thank you for your kind attention !