

Proving Explicit Statements

Problem-1. Let d, k, q be positive integers and k being odd, find greatest power of 2, in the decomposition of $\sum_{n=1}^{2^d k} n^q$.

AMM

Solution: we prove that if d being an even number or $d = 1$ the answer is $d - 1$ otherwise the answer is $2(d - 1)$.
if $q = 1$ it is obvious, if $d = 1$ the resulting sum is odd, now assume that $d > 1$ we can write the sum as the following:

$$(2^d k)^q + (2^{d-1} k)^q + \sum_{i=0}^{2^{d-1} k} (i^q + (2^d k - i)^q)$$

For even d , take the sum modulo 2^d , thus we find that $\sum_{n=1}^{2^d k} n^q \equiv 2 \sum_{n=1}^{2^{d-1} k} n^q \pmod{2^d}$, thus by use of induction on d we find that $\sum_{n=1}^{2^{d-1} k} n^q$ is congruent to 2^{d-2} modulo 2^{d-1} so $\sum_{n=1}^{2^d k} n^q \equiv 2^{d-1} \pmod{2^d}$ and it is the desired result.

For odd d , notice that by use of binomial theorem, $i^q + (2^d k - i)^q \equiv 2^d q k n^{q-1} \pmod{2^{2d-1}}$, so $\sum_{n=1}^{2^d k} n^q \equiv 2^d q k \sum_{n=1}^{2^{d-1} k} n^{q-1} \pmod{2^{2d-1}}$ since $q - 1$ is even by use of the former part we find that $\sum_{n=1}^{2^{d-1} k} n^{q-1} \equiv 2^{d-2} \pmod{2^{d-1}}$ so $\sum_{n=1}^{2^d k} n^q \equiv 2^{2d-2} \pmod{2^{2d-1}}$ and our proof is complete.

Variant :

Prove that $1^n + 2^n + \dots + (2^k - 1)^n$ is divisible by 2^k . **Kvant-M2277**

Solution: let's name the sum as A , if n is odd, then rewrite A as the form $(1^n + (2^k - 1)^n) + (2^n + (2^k - 2)^n) + \dots + ((2^{k-1} - 1)^n + (2^{k-1} + 1)^n) + 2^{n(k-1)}$ so it is obvious that A is divisible by 2^k .

If n is even, we made an induction on k assume the statement holds true for k then we must prove it for $k + 1$ implies that, we know that $a^n \equiv (2^{k+1} - a)^n \pmod{2^{k+1}}$ thus:

$$1^n + 2^n + \dots + (2^{k+1} - 1)^n = 1^n + (2^{k+1} - 1)^n + (2^n + (2^{k+1} - 2)^n) + \dots + ((2^{k+1} - 1)^n + (2^{k+1} + 1)^n) + 2^{kn} \equiv 2(1^n + 2^n + \dots + (2^k - 1)^n) + 2^{kn} \pmod{2^{k+1}}$$

By induction hypothesis $(1^n + 2^n + \dots + (2^k - 1)^n)$ is divisible by 2^k , thus $1^n + 2^n + \dots + (2^{k+1} - 1)^n$ must be divisible by 2^{k+1} so our proof is complete.

Problem-2. Prove that $1 + 3^3 + 5^5 + \dots + (2^n - 1)^{2^n - 1} \equiv 2^n \pmod{2^{n+1}}$

Kvant-M2252

Solution: at first we use two facts:

Fact-1: for odd k , $k^{2^n} \equiv 1 \pmod{2^{n+2}}$

Fact-2: $(k + 2^n)^k \equiv k^k (1 + 2^n) \pmod{2^{n+2}}$

The former , could be proven by the identity $k^{2^n} - 1 = (k - 1)(k + 1)(k^2 + 1) \dots (k^{2^{n-1}} + 1)$ or by induction on n , and the later could be proven by use of binomial theorem. Lets define $S_n = 1 + 3^3 + 5^5 + \dots + (2^n - 1)^{2^n - 1}$, then $S_{n+1} = S_n + R_n$, where $R_n = (2^n + 1)^{2^{n+1}} + \dots + (2^{n+1} - 1)^{2^{n+1} - 1}$, all the $2^n - 1$ terms of R_n , are of the form $m = 2^n + k$, $k < 2^n$ so one can write that:

$$m^m \equiv m^{2^n} \cdot m^k \equiv m^k \equiv k^k (1 + 2^n) \pmod{2^{n+2}}$$

Implies that $R_n \equiv (1 + 2^n)(1 + 3^3 + 5^5 + \dots + (2^n - 1)^{2^n - 1}) \equiv (1 + 2^n)S_n \pmod{2^{n+2}}$

Thus $S_{n+1} \equiv 2S_n (1 + 2^{n-1}) \pmod{2^{n+2}}$, now we take an induction on n by induction hypothesis we must have $S_n = 2^{n+1}k + 2^n$ for some integer k , thus :

$$S_{n+1} \equiv (2^{n+2}k + 2^{n+1})(1 + 2^{n-1}) \equiv 2^{n+1} \pmod{2^{n+2}}$$

Ensure that , our proof is complete.

Problem-3. Prove that $\sum_{k=0}^{\lfloor \frac{n}{p} \rfloor} (-1)^k \binom{n}{kp}$ is divisible by $p^{\lfloor \frac{n-1}{p-1} \rfloor}$, where p is prime and n is an integer , .

GMA-2013

Solution: assume that $S_n = \sum_{k=0}^{\lfloor \frac{n}{p} \rfloor} (-1)^k \binom{n}{kp}$, now we first prove preceding lemma:

Lemma: for all $\geq p$: $S_n - \binom{p}{1}S_{n-1} + \binom{p}{2}S_{n-2} + \dots + \binom{p}{p-1}S_{n-p+1} = 0$.

Proof: let ω be primitive p - th root of unity , we know that $\sum_{i=0}^{p-1} \omega^i = \begin{cases} 0 & p \nmid i \\ p & p \mid i \end{cases}$, thus one can see that $S_n = \frac{1}{p} \sum_{i=0}^{p-1} (1 - \omega^i)^n$, since ω^i , $i = 1, 2, \dots, p-1$ are roots of the polynomial $\frac{x^p - 1}{x - 1}$, then $1 - \omega^i$ are roots of the polynomial $\frac{1 - (1-x)^p}{x} = x^{p-1} - \binom{p}{1}x^{p-2} + \dots + \binom{p}{p-1}$, by setting $x = 1 - \omega^i$, $i = 0, \dots, p-1$ and adding up , for all $n \geq p$ one can find that:

$$S_n - \binom{p}{1}S_{n-1} + \binom{p}{2}S_{n-2} + \dots + \binom{p}{p-1}S_{n-p+1} = 0$$

Our proof is complete.

Back to the problem , we prove the statement by use of the induction on n , it is clear that for $n = 1, 2, \dots, p-1$. the statement holds , if $n \geq p$ then for all $1 \leq j \leq p-1$, $p \mid \binom{p}{j}$, by induction hypothesis , we now that for all $1 \leq j \leq p-1$, $p^{\lfloor \frac{n-j-1}{p-1} \rfloor}$ divides S_{n-j} , now we know that for all $1 \leq j \leq p-1$ $\left\lfloor \frac{n-j-1}{p-1} \right\rfloor + 1 = \left\lfloor \frac{n+p-j-2}{p-1} \right\rfloor \geq$

$\left\lfloor \frac{n-1}{p-1} \right\rfloor$. Thus all the terms of the sum $\binom{p}{1}S_{n-1} - \binom{p}{2}S_{n-2} + \dots - \binom{p}{p-1}S_{n-p+1}$ is divisible by $p^{\left\lfloor \frac{n-1}{p-1} \right\rfloor}$ and then that is true for S_n .

Induction and sequences

Problem-1. We know about the sequence $\{x_n\}$ such that $x_1 = \frac{2}{3}$, $x_{n+1} = \frac{3x_n+2}{3-2x_n}$ Is this sequence is eventually periodic?

Bulgarian TST-2011

Solution: Assume the answer is yes, then there are N, T such that for all $i \geq N$ we have $x_{i+T} = x_i$, we can assume that $N = 1$, indeed since $x_n = \frac{3x_{n+1}-2}{3-2x_{n+1}}$, by $x_{n+1+T} = x_{n+1}$ we deduce that $x_{n+T} = x_n$. lets assume $x_n = \frac{p_n}{q_n}$ in its reduced form, then $x_{n+1} = \frac{p_{n+1}}{q_{n+1}} = \frac{3p_n+2q_n}{3q_n-2p_n}$, lets $d = \gcd(3p_n+2q_n, 3q_n-2p_n)$ then $d = 1, 13$. lets define two sequence a_n, b_n such that $a_1 = 2, b_1 = 3$, and $a_{n+1} = 3a_n + 2b_n, b_{n+1} = 3b_n - 2a_n$, since these sequences are periodic modulo 13, easy calculation shows that, these sequences hasn't ever divisible by 13, thus we can say that $p_{n+1} = 3p_n + 2q_n, q_{n+1} = 3q_n - 2p_n$, now one can prove inductively that $p_{n+1}^2 + q_{n+1}^2 = 13(p_n^2 + q_n^2) = \dots = 13^n$, now if $x_{n+1} = \frac{2}{3}$ then $x_n = 0$ thus $13|p_n$ and then $13|q_n$, a contradiction.

Variant:

We know about the sequence $\{a_n\}$, such that a_0 is a positive integer and we have:

$$a_{n+1} = \begin{cases} \frac{a_n-1}{2} & \text{if } a_n \geq 1 \\ \frac{2a_n}{1-a_n} & \text{if } a_n < 1 \end{cases}$$

We know that $a_n \neq 2$, for $n = 1, 2, \dots, 2001$ and $a_{2002} = 2$. Find the value of a_0 .

Saint Petersburg

Solution-1: let $a_n = \frac{p}{q}$, then $a_{n+1} = \frac{p-q}{2q}$ or $a_{n+1} = \frac{2q}{q-p}$, since $\gcd(p-q, 2q) = 1$ or 2 thus in each step, the sum of nominator and denominator either $p+q$, $\frac{p+q}{2}$ thus sum of num and den of a_i doesn't increased, since $a_{2002} = 2$, so sum of num and den are equal to 3 we can assume that sum of num and den of a_0 is equal to $a_0 + 1$, so $a_0 + 1 = 3 \cdot 2^k$, so $a_n = 3 \cdot 2^{k-n} - 1$ since for $n = 1, 2, \dots, 2001$, $a_n \neq 2$ we may have $k = 2002$ and then $a_0 = 3 \cdot 2^{2002} - 1$.

Solution-2: lets define $b_n = \frac{1}{1+a_n}$ so we can rephrase the statement of the problem as :

$$b_{n+1} = \begin{cases} 2b_n & \text{if } 2b_n < l \\ 2b_n - 1 & \text{if } 2b_n > l \end{cases}$$

We can preclude the case with $b_n = 1$, which is fixed point of the sequence, so by induction we could find that $b_n = \{2^n b_0\}$, so if we set $1 + a_0 = k$ we have $b_n = \left\{\frac{2^n}{k}\right\}$, since $\left\{\frac{2^{2002}}{k}\right\} = \frac{1}{3}$, we must have $\frac{2^{2002}}{k} - \frac{1}{3} = l \in \mathbb{Z}$, so $3 \cdot 2^{2002} = k(3l + 1)$ thus $k = 3 \cdot 2^m$ for some positive integer $m \leq 2002$ and then $b_n = \left\{\frac{2^n}{k}\right\} = \left\{\frac{2^{n-m}}{3}\right\}$ since $b_m = \frac{1}{3}$ we find that $m = 2002$, so $k = 3 \cdot 2^{2002}$ and then $a_0 = 3 \cdot 2^{2002} - 1$.

Problem-3. We know about sequences $\{x_n\}, \{y_n\}$, such that $x_1 = \frac{1}{10}, y_1 = \frac{1}{8}, x_{n+1} = x_n + x_n^2, y_{n+1} = y_n + y_n^2$, prove that for all n, m , x_n, y_m couldn't be equal.

Saint Petersburg

First Solution: it is obvious that terms of both sequences are rational numbers assume that $x_n = \frac{p}{q}$ where

$\gcd(p, q) = 1$, whence $x_{n+1} = \frac{pq + p^2}{q^2}$ it is obvious that $\gcd(pq + p^2, q^2) = 1$, thus at each step the denominator of x_n is being square and analogously the denominator of y_n , thus denominator of x_n in its reduced form must be power of 10 and denominator of y_n in its reduced form must be power of 8, thus they couldn't being equal.

Second Solution: since $x_1 = 0.1, y_1 = 0.125$ we prove inductively that for all n , x_n ends with 1 and y_n ends with 5, let's assume the statement being true for all $k \leq n$ thus $x_n = \frac{x}{10^N}$, where x ends with 1 so $x_{n+1} = \frac{10^N x + x^2}{10^{2N}}$, since $10^N x$ ends with zero and x^2 ends with unit, $10^N x + x^2$ ends with unit and our hypothesis is proven, thus for all n , x_n ends with unit, by the same argument we find that y_n ends with 5, thus they couldn't being equal for all n, m .

Third Solution: since $x_2 = 0.11, x_3 = 0.1221, x_4 = 0.1221 + 0.1221^2 > 0.13$ so $x_3 < 0.125 = y_1 < x_4$ we prove by induction that: $x_{n+2} < y_n < x_{n+3}$, assume it holds true for all $k \leq n$, then $x_{n+2}^2 < y_n^2 < x_{n+3}^2$ adding the obtained inequality by the original inequality we find that: $x_{n+3} < y_{n+1} < x_{n+4}$. and our claim is proved. Now we receive an infinite sequence $x_3 < y_1 < x_4 < y_2 < x_5 < y_3 < \dots$ thus no terms could be equal at all.

Induction and Polynomials

Problem-1. Polynomials $F(x), G(x)$ has real coefficients and we know that the points

$$(F(1), G(1)), (F(2), G(2)), \dots, (F(2011), G(2011))$$

Are vertex of a 2011-gon in plane, prove that $\deg F$ or $\deg G \geq 2010$.

Solution: assume that the center of 2011-gon is located in (0,0). Thus one can say that :

$F(k) = \cos(\alpha + \frac{2k\pi}{2011})$, $G(k) = \sin(\alpha + \frac{2k\pi}{2011})$ (*) for $1 \leq k \leq 2011$. Now by induction on n , we show that if for $k = 1, 2, \dots, n$ the (*) relations holds ($n \leq 2011$) then at least one of F, G hasn't degree less than $n - 1$. Now take $n = 2011$ and we are done, for $n = 1$ the statement is true, now assume the statement is true for all $n \leq 2010$, let the polynomials F, G satisfies (*) for $k = 1, 2, \dots, n + 1$. define $F^*(x) = F(x + 1) - F(x)$, $G^*(x) = G(x + 1) - G(x)$, now for all $k = 1, 2, \dots, n$ we have

$$F^*(k) = -2 \sin \frac{\pi}{2011} \cdot \sin(\alpha + \frac{(2k+1)\pi}{2011}),$$

$$G^*(k) = 2 \sin \frac{\pi}{2011} \cdot \cos(\alpha + \frac{(2k+1)\pi}{2011})$$

We can divide F^*, G^* by $2 \sin \frac{\pi}{2011}$, thus they satisfies the problem criteria. We know that by induction hypothesis at least one of $\deg F^*$ or $\deg G^*(k) \geq n - 1$ since $-1 + \deg F = \deg F^*$, $-1 + \deg G = \deg G^*$ we have $\deg F$ or $\deg G \geq n$.

Problem-2. Prove there exist a quadratic polynomial $f(x)$, such that $f(f(x))$ has 4 non-positive real roots, and $\underbrace{f(f(\dots(x) \dots))}_n$ has 2^n real roots.

Bulgarian Olympiads

Solution: Without loss of generality assume that $f(x)$ has positive leading coefficient, (since change the sign of leading coefficient doesn't change locations of roots) it is obvious that $f(x)$ must have two real roots, say $x_1 < x_2$, if $x_2 > 0$, then there exist a positive real number s such that $f(s) = x_2$, thus $f(f(s)) = f(x_2) = 0$ and $f(f(x))$ find positive real root, a contradiction, thus we may assume that $x_1 < x_2 \leq 0$. since the roots of $f(f(x))$ are indeed, roots of the equations: $f(x) = x_1$, $f(x) = x_2$ and both of the equations must have 2 real roots, let's show the minimal value of the f , by m . So we find that $m < x_1 < x_2 \leq 0$. And then roots of $f(f(x))$ lies on the interval (x_1, x_2) . Now we complete our proof by induction, we prove that for all $n > 1$, the polynomial f_n has 2^n real roots lies on the interval (x_1, x_2) . For $n = 2$ it is clear, assume y_1, \dots, y_{2^n} being roots of $f_n = \underbrace{f(f(\dots(x) \dots))}_n$ lies on the interval (x_1, x_2) , we find that roots of f_{n+1} are roots of $f(x) = y_k$ where $1 \leq k \leq 2^n$. since $m < x_1$ all the equations of the form $f(x) = y_k$ has two real roots, lies in the interval (x_1, x_2) . whence the polynomial f_{n+1} , has 2^{n+1} real roots lies on the interval (x_1, x_2) . We are done.

Problem-3. We know about the polynomial $P(x, y)$ with real coefficients such that, there exist a function f , such that

$$P(x, y) = f(x + y) - f(x) - f(y)$$

Prove that, f can take polynomial values infinitely many times.

Saint-Petersburg Olympiads

Solution: we define polynomial $f_0(x)$ such that $(x, 1) = f_0(x + 1) - f_0(x) - f_0(1)$ and $f_0(1) = f(1)$ it is obvious that one can find all the coefficients of $f_0(x)$ uniquely implies that $f_0(x)$ determines uniquely, now one can find that $f_0(x + 1) - f_0(x) = f(x + 1) - f(x)$ thus we can inductively prove that $f(n) = f_0(n)$ for all positive integers n , thus for all positive integers x, y we have $P(x, y) = f_0(x + y) - f_0(x) - f_0(y)$ and we are done.

On the degree of Polynomials

Problem-1. Prove that there exist positive real numbers a_0, a_1, \dots, a_n such that the polynomial $\pm a_n x^n \pm a_{n-1} x^{n-1} \pm \dots \pm a_0$ for all choose of sign has n real roots. **IMC-2014**

Solution: we prove the problem statement, by use of induction on n , by induction hypothesis, the polynomial $\overline{P(x)} = \pm a_n x^{n+1} \pm a_{n-1} x^{n-1} \pm \dots \pm a_0 x$, where $a_0 \neq 0$, has $n + 1$ distinct real roots, thus it hasn't any double root, thus any of its local extreme wasn't its roots, so for all polynomial $\overline{P(x)}$, for all of its n extremes s_1, \dots, s_n there are $\varepsilon > 0$, such that $|\overline{P(s_i)}| > \varepsilon$, so we prove all of the polynomial $P(x) = \pm a_n x^{n+1} \pm a_{n-1} x^{n-1} \pm \dots \pm a_0 x \pm \varepsilon$, has exactly $n + 1$ real roots, since $\overline{P(x)}$ has local extremum $s_1 < \dots < s_n$, such that $P(s_i), P(s_{i+1})$ has different signs thus it has roots in any interval of (s_i, s_{i+1}) adding up $(-\infty, s_1), (s_n, +\infty)$ so by induction, our proof is complete.

Variant:

Is there exist a sequence of nonzero real numbers $a_0, a_1, \dots, a_n, \dots$ such that for all n the polynomial $P_n(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_0$ has n distinct real roots? **Putnam-1990**

Solution: besides previous solution, we can construct an infinite sequence $a_n = (-1)^n 10^{-n^2}$, thus $(-1)^k 10^{-k^2} P(10^{2k}) = \sum_{i=0}^n (-1)^{i-k} 10^{-(i-k)^2} = \sum_{j=-k}^{j=k} (-1)^j 10^{-j^2} > 1 - 2 \sum_{j=1}^{\infty} 10^{-j^2} > 0$

Henceforth, $P(1), \dots, P(10^{2n})$ changes their signs, alternatively....

Problem-2. We know about function $f(x)$ such that for all integer x , we have $f(x) \in \mathbb{Z}$, for all prime p there exist a polynomial $Q_p(x)$ of degree less than or equal 2013, with integer coefficients such that $f(n) - Q_p(n)$ is divisible by p , prove that there are polynomial $g(x)$ with real coefficients such that for all positive integers we have $f(n) = g(n)$. **Moscow-2013**

First Solution: we prove this problem, by induction on degree of polynomial $Q_p(x)$ such that for all polynomial $Q_p(x)$ of degree less than or equal to k , $f(x)$ takes polynomial values in positive integer points. For $= 0$, $Q_p(x)$ must be constant and $f(n) - Q_p(n) = f(n) - c$ is divisible by p thus for all $m, n \in \mathbb{N}$ $f(m) - f(n)$ is

divisible by p , so take p large enough, we conceive that $f(x)$ must be constant integer. Now we prove two following lemma:

Lemma-1: let $h(x)$ be a polynomial of degree d , then the polynomial $\Delta h = h(x+1) - h(x)$ is of degree $d-1$.

Proof: the proof is indeed obvious.

Lemma-2: if for all integer x , the function $\Delta h(x)$ takes a polynomial value of degree less than or equal to $d-1$, then for all integer x , we have $h(x) = h_1(x)$ for some polynomial h_1 of degree less than or equal to d .

Proof: we prove this lemma by induction on d , for $d=1$, we can say that $h(x) = h(0) + cx$ for all integer x . now assume $\Delta h(x) = ax^d + \dots$ define $h_0(x) = h(x) - \frac{a}{d+1}x(x-1)\dots(x-d)$ then

$$\Delta h_0 = \Delta h(x) - ax(x-1)\dots(x-d+2)$$

Then $\deg \Delta h_0 \leq d-1$, so by induction hypothesis, h_0 is equal to polynomial of degree less than or equal to d since $\deg h_0(x) - h(x) \leq d+1$, whence $h(x)$ in integral points take the value of a polynomial of degree less than or equal to $d+1$, ensure that our proof is complete.

Take back to our problem, since the function Δf satisfies the statement of the problem (indeed $\Delta f(x) - \Delta Q_p(x)$ is divisible by p in positive integer points, note that $\deg \Delta Q_p(x) \leq k-1$. so by induction hypothesis $\Delta f(x)$ take polynomial values. thus by lemma-2, the polynomial $f(x)$ takes polynomial value of degree less than or equal one unit added up to whatever polynomial $\Delta f(x)$ takes its values on integral points.

Second Solution: by use of lagrange interpolation formula one can find that we can find the polynomial $f_0(x)$, by the values of $f(1), \dots, f(2014)$ with rational coefficients of degree less than or equal to 2013, now if we take $c = 2013!^2$, then one can easily find that $cf_0(x)$ has integer coefficients, take a prime $p > c$, then $\deg cQ_p(x) - cf_0(x) \leq 2013$, and has 2014 incongruent roots modulo p , thus polynomial $cQ_p(x) - cf_0(x)$ is zero polynomial modulo all large prime p note that:

$$c(f(x) - Q_p(x)) + c(Q_p(x) - f_0(x)) = c(f(x) - f_0(x))$$

Is divisible by sufficiently large p , and thus one can find that for all integer x , we must have: $f(x) = f_0(x)$. (fix x and take p , sufficiently large).

Problem-3. Prove that If a rational function that is not a polynomial assumes rational values at all positive integral points, then it is the quotient of two relatively prime polynomials with integral coefficients.

Solution: Let the function in question be $R(x) = \frac{P(x)}{Q(x)}$ where $P(x), Q(x)$ are relatively prime polynomials, r the sum of the degrees of $P(x), Q(x)$. For $r = 0$ the theorem is obvious. Consider if necessary $\frac{1}{R(x)}$ instead of $R(x)$ and assume that the degree of $P(x)$ is not less than that of $Q(x)$ and furthermore that a is a positive integer such that $Q(a) \neq 0$. Then $\frac{P(a)}{Q(a)}$ is rational and the rational function $\frac{1}{x-a} \left(R(x) - \frac{P(a)}{Q(a)} \right) = \frac{P_1(x)}{Q(x)}$ is a rational function whose value is rational for integral $x, x > a$, and the degree of $P_1(x) = \frac{Q(x)Q(a) - Q(x)P(a)}{Q(a)(x-a)}$ is less than that of $P(x)$; thus the sum of the degrees of $P_1(x), Q(x)$ is less than that of $P(x)$ and $Q(x)$. so by induction we are done.

Problem-4. Prove that If a rational function has integral values for infinitely many integral values of the variable, then it is a polynomial.

Solution: The function $f(x) = \frac{P(x)}{Q(x)}$ is considered, where $P(x)$ and $Q(x)$ are polynomials with integral coefficients. We can find an integer q such that $qf(x) = G(x) + r(x)$, where $G(x)$ denotes a polynomial with integral coefficients and $r(x)$ a rational function whose numerator is of lower degree than its denominator. The value of $r(x)$ is integral for infinitely many integral values of x . Since $\lim_{x \rightarrow \infty} r(x) = 0$, from a sufficiently large points we must have $|r(x)| < 1$, thus $r(x) = 0$ and hence is zero for all x for a rational function. so we are done.

Miscellaneous

Problem-1. Let $P(x) = a_d x^d + \dots + a_0$ prove that the polynomial is divisible by $(x-1)^m$ if and only if for all $s = 0, 1, \dots, m-1$ we have $a_n(n+1)^s + a_{n-1}n^s + \dots + a_1 2^s + a_0 = 0$

Solution: let $(x) = (x-1)^m Q(x)$, where $Q(x) = \sum_{i=0}^{n-m} q_i x^{n-m-i}$ since $(x-1)^m = \sum_{j=0}^m \binom{m}{j} x^{m-j}$, then:

$$P(x) = \left(\sum_{i=0}^{n-m} q_i x^{n-m-i} \right) \left(\sum_{j=0}^m \binom{m}{j} x^{m-j} \right) = \sum_{i=0}^{n-m} q_i \sum_{j=0}^m (-1)^j \binom{m}{j} x^{n-i-j}.$$
 The problem statement reduces to : $\sum_{i=0}^{n-m} q_i \left(\sum_{j=0}^m (-1)^j \binom{m}{j} (n-i-j+1)^s \right) = 0$. the term $\sum_{j=0}^m (-1)^j \binom{m}{j} (n-i-j+1)^s$ is equal to m -th difference of the polynomial x^s , which is identically zero.

Proof of Sufficiency: we prove this by induction on m , the case $m = 1$ is clear, assume the statement is true for $m = k$ and divide $P(x)$ by $(x-1)^{k+1}$ now we have $P(x) = (x-1)^{k+1} q(x) + r(x)$ where $\deg r(x) \leq k$. by induction hypothesis, we know that $P(x)$ is divisible by $(x-1)^k$ so $r(x) = a(x-1)^k$ the necessity condition depicts that $(x-1)^{k+1} q(x)$ satisfies the conditions for $s = 0, 1, \dots, k$. now we only need to check the condition for $r(x)$, for $s = k$, we have $r(x) = a \sum_{j=0}^k (-1)^j \binom{k}{j} x^{k-j}$ thus we have :

$$a((k+1)^k - k.k^k + \dots) = 0$$

The expression in the bracket is equal to ! . thus $a = 0$.

Comment: Instead of $1, 2, \dots, n+1$ we can use the $n+1$ consecutive terms of Arithmetic progression.

Problem-2. let $P(x)$ has n -distinct real root larger 1 and smaller 0, prove that:

1. $P'(x)$ has a root greater than $1 - \frac{1}{n}$.

2. Absolute value of difference between maximum and minimum root of $P'(x)$ is greater than $\sqrt{1 - \frac{2}{n}}$.

Saint Petersburg

Solution-1: without loss of generality assume that leading coefficient of the polynomial $P(x)$ is positive, we know that all the roots of $P'(x)$ lies on the interval $(0,1)$ so, for $x \geq 1$ we must have $P'(x) > 0$, if we show roots of $P(x)$ by $x_0 = 0 < x_1 < \dots < x_{n-2} < x_{n-1} = 1$. if all the roots of $P'(x)$ say $x_0 = 0 < y_1 < x_1 < y_2 < \dots < x_{n-2} < y_{n-1} < x_{n-1} = 1$ being less than $1 - \frac{1}{n}$ then we must have $P'(1 - \frac{1}{n}) > 0$ and $x_{n-2} < 1 - \frac{1}{n}$. so $1 - \frac{1}{n}$ lies between two consecutive roots of the polynomial $P(x)$ so we must have $P(1 - \frac{1}{n}) < 0$ we know that $P'(x) = P(x) \cdot \sum_{i=0}^{n-1} \frac{1}{x-x_i}$, set $x = 1 - \frac{1}{n}$ in there, we find that:

$$P'(1 - \frac{1}{n}) = P(1 - \frac{1}{n}) \cdot \sum_{i=0}^{n-1} \frac{1}{1 - \frac{1}{n} - x_i}$$

We know that $\sum_{i=0}^{n-1} \frac{1}{1 - \frac{1}{n} - x_i} = -n + \sum_{i=0}^{n-2} \frac{1}{1 - \frac{1}{n} - x_i}$, we also know that for $0 \leq i \leq n-2$

$$0 < 1 - \frac{1}{n} - x_i \leq 1 - \frac{1}{n}$$

Thus $\sum_{i=0}^{n-1} \frac{1}{1 - \frac{1}{n} - x_i} > -n + \frac{n-1}{1 - \frac{1}{n}} = 0$, since $P(1 - \frac{1}{n}) < 0$, we must have $P'(1 - \frac{1}{n}) < 0$, a contradiction.

Solution-2: At first we prove the following interesting lemma:

Lemma: let $0 < x_2 \leq y_2 \leq \dots \leq x_{k+1} < 1$ and $A = \frac{x_2 x_3 \dots x_{k+1}}{y_2 y_3 \dots y_k}$, $B = \frac{(1-x_2) \dots (1-x_{k+1})}{(1-y_2) \dots (1-y_k)}$ then $A + B \leq 1$.

Proof: for $k = 1$, the statement is obvious. now we prove the truth of the statement for k , fix all x_k, y_k and notice that $x_{k+1} \in [y_k, 1]$. if $x_{k+1} = 1$ we are done, if not, note that the expression $A + B$ is linear respect to x_{k+1} so attains its maximum values at the one ends of the interval if $x_{k+1} = y_k$ by induction hypothesis we are done.

Back to our problem, since $x_1 = 0 < y_1 < x_2 < y_2 < \dots < x_{n-1} < y_{n-1} < x_n = 1$, and we can say that

$P(x) = x(x - x_2) \dots (x - x_n)$ and $P'(x) = n(x - y_1) \dots (x - y_{n-1})$, since the coefficient of x in $P(x)$ is $\pm x_2 \dots x_n$ and the coefficient of constant term in $P'(x)$ is $\pm n y_1 \dots y_{n-1}$, one can easily find that :

$$x_2 \dots x_{n-1} = n y_1 \dots y_{n-1} \quad (*)$$

Now change $P(x)$ by $P(1 - x)$ one can see that $(1 - y_1) \dots (1 - y_{n-1}) = (1 - x_2) \dots (1 - x_{n-1})$.
 (**) now since for $i = 2, \dots, n - 1$ we have $1 - x_i \leq 1 - y_{i-1}$ then $n(1 - y_{n-1}) \leq 1$ so $y_{n-1} \geq 1 - \frac{1}{n}$
 which is another proof for the first part, use the lemma for $k = n - 2$, then one can find that :

$$\frac{x_2 \dots x_{n-1}}{y_2 \dots y_{n-2}} + \frac{(1-x_2) \dots (1-x_{n-1})}{(1-y_2) \dots (1-y_{n-2})} \leq 1$$

By use of our (*), (**) we find that $n y_1 y_{n-1} + n(1 - y_1)(1 - y_{n-1}) \leq 1$.

So $(y_1 - y_{n-1})^2 = (1 - y_1 - y_{n-1})^2 - 2(y_1 y_{n-1} + (1 - y_1)(1 - y_{n-1})) + 1 \geq 1 - \frac{2}{n}$. and we are done.

Induction and functional equations

Problem-1. Find all function $f: \mathbb{N} \rightarrow \mathbb{N}$ such that

$$2f(mn) \geq f(m^2 + n^2) - f^2(m) - f^2(n) \geq 2f(m)f(n)$$

Silk Road Math Competitions

Solution: define $A = \{k \mid f(kn) = k^2 f(n)\}$, $A_1 = \{k \mid f(k) = k^2\}$, from the above inequalities we find that

1. $f(mn) \geq f(m) \cdot f(n)$
2. $f(n^2 + m^2) \geq (f(m) + f(n))^2$
3. $f^2(m) + f^2(n) + 2f(mn) \geq f(n^2 + m^2)$

Now we prove following lemmas:

Lemma-1: if $f(mn) = f(m) \cdot f(n)$ then $f(n^2 + m^2) = (f(m) + f(n))^2$.

Proof: by three inequalities, it is clear.

Now take $n = 1$, then $f(1) = 1$, set $m = 1$ in the above lemma we receive $f(n^2 + 1) = (1 + f(n))^2$ (4), set $n = 1$ we find that $f(2) = 4$, take $m = n$ in the inequalities 1, 3 we find that: $f^2(n) \geq f(n^2)$, $f(n^2) \geq f^2(n)$. thus $f(n^2) = f^2(n)$. we prove following lemma:

Lemm-2: if for all k , $f(kn^2) = k^2 f^2(n) = k^2 f(n^2)$ then $k \in A$.

Proof: $f(k) = f(k \cdot 1^2) = k^2 f^2(1) = k^2$, and $k^2 f^2(n) = f(kn^2) \geq f(kn) \cdot f(n)$

so $f(kn) \leq k^2 f(n)$, on the other hand $f(kn) \geq f(k) \cdot f(n) = k^2 f(n)$ so $f(kn) = k^2 f(n)$, whence $k \in A$.

We know $f(n^2) = f^2(n)$ so $1 \in A$, then by lemme-1, for $m = n$ we have $f(2n^2) = 4f(n^2)$ so $2 \in A$, conclude that $f(2n) = 4f(n)$. Now set $n = p_1 \dots p_t$ where p_i are primes, we know that

$$f(n) \geq f(p_1) \dots f(p_t)$$

We prove by induction that $f(n) \geq n^2$, it is clear for $n = 1, 2$. let $n = mk$ where $1 < m, k < n$ by induction hypothesis $f(n) \geq f(m) \cdot f(k) \geq m^2 k^2 = n^2$. if $n \equiv 1 \pmod{4}$ there are integers x, y such that $n = x^2 + y^2$ and $x, y < n$. thus by induction hypothesis:

$$f(n) = f(x^2 + y^2) \geq (f(x) + f(y))^2 \geq (x^2 + y^2)^2 = n^2$$

If $n \equiv 3 \pmod{4}$ then $n^2 + 1 = 2p_1 \dots p_t$ where $p_i \equiv 1 \pmod{4}$ which wasn't necessarily distinct, so there are positive integers x_i, y_i such that $x_i^2 + y_i^2 = p_i \leq \frac{n^2 + 1}{2} < n^2$ so $x_i, y_i < n$, then by induction hypothesis: $f(p_i) = f(x_i^2 + y_i^2) \geq (f(x_i) + f(y_i))^2 \geq (x_i^2 + y_i^2)^2 = p_i^2$. thus:

$$f(n^2 + 1) \geq f(2) \cdot f(p_1) \dots f(p_t) \geq 4p_1^2 \dots p_t^2 = (n^2 + 1)^2$$

Since $f(n) = \sqrt{f(n^2 + 1)} - 1$ we have $f(n) \geq n^2 + 1 - 1 = n^2$. We are done.

Now we complete our proof with two different methods:

Method-1: we use induction to prove $f(n) = n^2$ assume it holds true for all $1 \leq i \leq m$, now one can see that $f(2(m^2 + 1)) = 4f(m^2 + 1) = 4(m^2 + 1)^2$ now set $n = m - 1$, $m = m + 1$ in the inequality (2) we find that

$$4(m^2 + 1)^2 = f(2(2(m^2 + 1))) = f((m - 1)^2 + (m + 1)^2) \geq (f(m + 1) + f(m - 1))^2 = (f(m + 1) + (m - 1)^2)^2$$

Thus we can find $f(m + 1) \leq (m + 1)^2$ but we proved $(m + 1) \geq (m + 1)^2$. so the equality occurred. and our inductive proof is complete.

Method-2: we continue our proof, by proving thi lemma:

Lemma-3: if $a, \in A$ then so is $ab, a^2 + b^2$.

Proof: since $f(an) = a^2 f(n), f(bn) = b^2 f(n)$ we find that $(abn) = a^2 f(bn) = (ab)^2 f(n)$. we know that $x = an, y = bn$ satisfies the equality $f(xy) = f(x) \cdot f(y)$ hence by lemma-1 we find that

$$f((a^2 + b^2)n^2) = f(x^2 + y^2) = (f(x) + f(y))^2 = (a^2 + b^2)^2 f^2(n)$$

Then our proof is complete.

Lemma-4: if $k \in A$ and $d|k$ then $d \in A$.

Proof: $k^2 f(n) = f(kn) \geq f(dn) \cdot f\left(\frac{k}{d}\right) \geq \left(\frac{k}{d}\right)^2 f(dn)$ therefore $d^2 f(n) \geq f(dn) \geq f(d) \cdot f(n) \geq d^2 f(n)$. thus $f(dn) = d^2 f(n)$ and it means that $d \in A$.

We construct set of primes $p_1 = 2, p_{i+1} | (p_1 \dots p_i)^2 + 1$ by induction on i and use of lemma-3 and lemma-4, we can see that $p_i \in A$.

Lemma-5: let $r = a^2 + b^2 \in A_1$ then $a, b \in A_1$

Proof: since $r \in A_1$ then $f(r) = r^2 = f(a^2 + b^2) \geq (f(a) + f(b))^2 \geq (a^2 + b^2)^2 = r^2$ so the equality occurred and then $f(a) = a^2, f(b) = b^2$.

Lemma-6: if $r \in A_1$ and $d|r$ then $d \in A_1$

Proof: write $r = kd$, then $d^2 \leq f(d) \leq \frac{f(r)}{f(k)} \leq \frac{r^2}{k^2} = d^2$ so $(d) = d^2$. shows that $d \in A_1$.

Now since there are infinitely many $p_i \equiv 1 \pmod{4}$ in the set A , we can find that $f(p_i) = p_i^2$ we know that for each of which there are integers x_i, y_i such that $x_i^2 + y_i^2 = p_i$. take arbitrary positive integer n and consider $\gcd(n, y_i)$, whence there is a certain number d which occurred infinitely many times, define the set

$$B = \{i | \gcd(n, y_i) = d\}$$

So $y_i = dz_i$ and $\gcd\left(z_i, \frac{n}{d}\right) = 1$ whence there are $i < j \in B$ such that $\frac{x_i}{z_i} \equiv \frac{x_j}{z_j} \pmod{\frac{n}{d}}$ set $s = |x_i y_j - x_j y_i| = d |x_i z_j - x_j z_i| \equiv 0 \pmod{n}$ now set $= x_i x_j + y_i y_j$, so $p_i p_j = s^2 + t^2$ we know that $p_i p_j \in A$ and $f(p_i p_j) = (p_i p_j)^2$ so $p_i p_j \in A_1$ so $s, t \in A_1$. we now that $n|s$ whence by lemma-6, $n \in A_1$ then $f(n) = n^2$.

Finally Our long-run proof is complete!

Problem-2. Prove that for all positive integers $a, b > 1$, there are a function $f: \mathbb{N} \rightarrow \mathbb{N}$ such that :

$$f(af(n)) = bn$$

Brazilian Olympiad

Solution: define two increasing sequence $\{S_n\}_{n=1}^{\infty} = \mathbb{N} - \{a, 2a, 3a, \dots\}$, $\{R_n\}_{n=1}^{\infty} = \mathbb{N} - \{b, 2b, \dots\}$ in turn, of non multiples a, b . one can easily find that, if for example $R_l = R_k$, then $l = k$. we prove that the following function satisfies the problem conditions:

$$f(n) = \begin{cases} R_k & a \nmid n, S_k = n \\ bS_l & n = ak, b \nmid k, R_l = k \\ abf(j) & n = abj \end{cases}$$

Now we prove that the above function is indeed a solution, we divide our proof in 3 cases:

Case-1: $a \nmid n$. We can write $S_k = n$. Then $f(af(n)) = f(aR_k) = bS_l = bn$ (since $\{S_n\}_{n=1}^{\infty}$ is an increasing sequence, then $k = l$)

Case-2: $n = ak, b \nmid k$. Then $f(af(n)) = f(af(ak)) = f(abS_l), k = R_l$ now $f(abS_l) = abf(S_l) = abR_k, (S_k = S_l)$ thus $f(abS_l) = abR_l = abk = nb$

Case-3: $n = abj$, then $f(af(n)) = f(af(abj)) = f(ab(af(j))) = abf(af(j))$ now if $ab \nmid j$ then by above cases: $f(af(j)) = bj$ and we are done. if $ab \mid j$ set $j = abi$ and then $i < j$ now we use induction on j and by use of induction hypothesis on i we can find

$$f(af(j)) = f(af(abi)) = f(ab(af(i))) = abf(af(i)) = ab.bi = bj$$

So, if $n = abj$ then $f(f(af(n))) = ab.bj = b.n$

Problem-3. Find all function: $\mathbb{R}^+ \rightarrow \mathbb{R}$, such that $f(x) + f(y) \leq \frac{1}{2}f(x+y)$, $\frac{f(x)}{x} + \frac{f(y)}{y} \geq \frac{f(x+y)}{x+y}$

Japanese Olympiads-2007

Solution: set $x = y = t$, then $4f(t) \leq f(2t)$ and $2\frac{f(t)}{t} \geq \frac{f(2t)}{2t}$ then $f(2t) = 4f(t)$ by induction it could be find that $f(2^m t) = 2^{2m} f(t)$. for all positive integer m . Define $g(x) = \frac{f(x)}{x}$ we now that for $x = 2^m, m \in \mathbb{N}$,

we have $g(nt) = ng(t)$ since $g(x) + g(y) \geq g(x+y)$, so inductively $g(nt) \leq ng(t)$. now take $2^m > n$, then $g(2^m t) \leq g(nt) + g((2^m - n)t) \leq ng(t) + (2^m - n)g(t) = 2^m g(t) = g(2^m t)$ see the equality occurs, thus we can see that $g(2^m t) = g(nt) + g((2^m - n)t)$ thus for all integer n , we have $g(nt) = ng(t)$. We know that $f(2t) = 4tg(t)$, $f(3t) = 9tg(t)$ take $x = 2t$, $y = t$ in the first inequality then $f(2t) + f(t) \leq \frac{f(3t)}{2}$ thus $5tg(t) \leq \frac{9}{2}tg(t)$, then $g(t) \leq 0$ for all positive t .

Then take $0 < x \leq y$ so $g(x) \geq g(x) + g(y-x) \geq g(y)$ so $g(x)$ is a decreasing function. take $(1) = a \leq 0$. then for all positive integer n we find that $g(n) = a \cdot n$, if $\gcd(p, q) = 1$. then $q \cdot g\left(\frac{p}{q}\right) = g(p) = a \cdot p$ so $g\left(\frac{p}{q}\right) = \frac{a \cdot p}{q}$. we also know that for all $t > \frac{p}{q}$, we have $g(t) < \frac{a \cdot p}{q}$. if there are $t \in \mathbb{R} - \mathbb{Q}$ such that $g(t) < at$, there exist a rational r , such that $g(t) < r < at$, since g is decreasing we have

$$ar = g(r) > g(at) = ag(t)$$

Since a isn't positive we must have $g(t) < at$, a contradiction, analogously for the case of existence of t for which $g(t) > at$. Thus $g(t) = at$. For all real t , thus $f(t) = at^2$ clearly satisfies the conditions of the problem.

On sum of num and den

Problem-2. Find all function $f: \mathbb{Q}^+ \rightarrow \mathbb{Q}^+$ such that $f\left(\frac{x}{x+1}\right) = \frac{f(x)}{x+1}$, $f(x) = x^3 f\left(\frac{1}{x}\right)$

Turkish Olympiads

Solution: we prove that the only functions that satisfy the problem conditions are $f_a(x) = a \cdot \frac{n^2}{d}$ where d, n are relatively prime positive integers, it is easy to check, function $f_a(x)$ satisfies. now we made an induction on $\max(n, d)$ (or $n + d$ respectively) for $n = d = 1$, we have $f(1) = a = f_a(1)$. Now we prove inductively that for all ≥ 2 , we must have $f\left(\frac{n}{s}\right) = f_a\left(\frac{n}{s}\right)$ where $\gcd(n, s) = 1$. assume the statement was true for $k = \max(n, d) < k$, now if $\gcd(n, d) = 1$, $\max(n, d) = k$ we know that $n \neq d$, so we have two cases. First. If $d < n$, then $\gcd(n, d - n) = 1$ so $f\left(\frac{n}{d}\right) = f\left(\frac{\frac{n}{d-n}}{1 + \frac{n}{d-n}}\right) = \frac{f\left(\frac{n}{d-n}\right)}{\frac{d}{d-n}} = a \cdot \frac{n^2}{d}$. Second. If $d < n$ then $f\left(\frac{n}{d}\right) = f\left(\frac{1}{\frac{d}{n}}\right) = \frac{f\left(\frac{d}{n}\right)}{\frac{d^3}{n^3}} = a \cdot \frac{n^2}{d}$. now our proof is complete.

Problem-3. When we have number x in hand, two operation $x \rightarrow \frac{1+x}{x}$, $x \rightarrow \frac{1-x}{x}$, is permitted, Is it true that each of the nonzero rational number can be obtained with a finite number of such operations? **Moscow-Olympiads-2007**

Solution: lets $f(x) = \frac{1+x}{x}$, $g(x) = \frac{1-x}{x}$ one can see that $f(g(f(x))) = -x$, $f(g(f(g(f(x)))))) = \frac{1}{1+x}$
so $f(g(f(f(g(f(x)))))) = x$ ($x \neq -1$), $f(g(f(g(f(g(x)))))) = x$, ($x \neq 1$)

Now number 2 could be obtain from -2 and vice versa, since $f(-2) = \frac{1}{2}$, $g(\frac{1}{2}) = 1$ we can receive number 1 from 2, besides, since $g(-1) = -2$, and -2 can get 2 from -1 . since

$$g(2) = \frac{-1}{2}, f\left(\frac{-1}{2}\right) = -1$$

we can get -1 from -2 . the cycle is complete(i.e. $2 \leftrightarrow 1$, $\underbrace{-2 \rightarrow -1 \rightarrow 2}_{2 \leftrightarrow -2}$)

now by induction on sum of num and den of the positive irreducible fraction $\frac{m}{n}$, we construct all positive rational number then by the equality $f(g(f(x))) = -x$ we construct all negative ones. For $m + n = 2$ we are done, assume the statement holds true for all $m + n < k$. so if, $m > n$ then $\frac{m}{n} = 1 + \frac{1}{\frac{n}{m-n}}$ since $n + m - n = m < k$ we can construct $\frac{n}{m-n}$ and so forth $\frac{m}{n}$ (by f). if $m < n$ then $\frac{m}{n} = \frac{1}{1 + \frac{n-m}{m}}$ since $n - m + m = n < k$. We can construct $\frac{n-m}{m}$ and so forth $\frac{m}{n}$ (by $f(g(f(g(f(x))))))$. and we are done.

Induction and Number Theory

Problem-1. Prove that for all positive integer n there exist an integer number such that decimal expansion of its square starts with n ones and ends with a block of length n with combinations of ones and two s.

Moscow-2014

Solution: we prove this by induction on n , such that for all n there exist an integer m_n whose decimal expansions ends with units and decimal expansion of m_n^2 ends with a block of length n with combinations of ones and two s.

for $n = 1$ it is clear, assume the statement of the problem holds for $n = k$, and the desired number was m_k , lets define $p_a = m_k + a \cdot 10^k$ for some $a \in \{1, 2, \dots, 9\}$ ends with unit. thus $p_a^2 = m_k^2 + 2a \cdot m_k \cdot 10^k + a^2 \cdot 10^{2k}$, the number m_k^2 ends with k ones and twos, the number $2a \cdot m_k \cdot 10^k$ ends with k zeros and $a^2 \cdot 10^{2k}$

ends with $2k$ zeros, name the $k + 1$ - th digits of m_k^2 by b and $k + 1$ - th digits of $2a \cdot m_k \cdot 10^k$ is $2a \pmod{10}$ (since m_k ends with unit) thus $k + 1$ - th digits of p_a^2 is $b + 2a \pmod{10}$ if b odd take a such that last digits of $b + 2a \pmod{10}$ being 1, if b being even, take a such that last digits of $b + 2a \pmod{10}$ is 1, 0. thus we can take $p_a = m_{k+1}$.

Now set $c_n = \underbrace{11 \dots 1}_n \cdot 10^{4n}$ and $d_n = c_n + 10^{4n}$, now:

$$\sqrt{d_n} - \sqrt{c_n} = \frac{d_n - c_n}{\sqrt{d_n} + \sqrt{c_n}} = \frac{10^{4n}}{\sqrt{d_n} + \sqrt{c_n}} > \frac{10^{4n}}{2 \cdot 10^{3n}} > 1$$

Thus there exist an integer in the interval $(\sqrt{c_n}, \sqrt{d_n})$ such that its square starts with n units, show by p_n now construct the number $p_n \cdot 10^k + m_n$ take k large enough (i.e. larger than number of digits of $2p_n m_n, m_n^2$). now one can see that $(p_n \cdot 10^k + m_n)^2 = p_n^2 \cdot 10^{2k} + 2p_n m_n \cdot 10^k + m_n^2$, now the first n digits of this number characterize by p_n^2 (string of units) and the last n digits of that number is characterized by m_n^2 (strings of units and twos)

Problem-2. Prove that for all positive integer, there exist, integer k such that $S(k) = n, S(k^2) = n^2, S(k^3) = n^3$.

Moskow-Olympiads-2013.

Solution: we prove more general statement: for any positive integer n there exist a nonnegative integer k consist only of 0,1 and k^2 consist only of 0,1,2 in theirs decimal representations such that:

$$S(k) = n, S(k^2) = n^2, S(k^3) = n^3.$$

the statement by induction on n , for $n = 1$ it is obvious, assume the statement holds true for n , now we must prove that there exist an integer m such that $S(m) = n + 1, S(m^2) = (n + 1)^2, S(m^3) = (n + 1)^3$. Lets name the number which satisfies the statement for n by t and assume that t has k digits, now take $m = t + 10^{10k}$, then we can see that $S(m) = 1 + S(t) = n + 1$ and

$$S(m^2) = S(t^2 + 2t \cdot 10^{10k} + 10^{20k}) = S(t^2) + 2S(t) + 1 = (n + 1)^2$$

Since, $t^2 < 10^{2k}, 2t \cdot 10^{10k} < 10^{20k}$, number $t^2, 2t \cdot 10^{10k}, 10^{20k}$ hasn't any in-common digits which affects theirs addition. And m^2 consists only of 0,1,2. now we can see that

$$m^3 = t^3 + 3t^2 \cdot 10^{10k} + 3t \cdot 10^{20k} + 10^{30k}$$

By the same argument, one can find that $S(m^3) = (n + 1)^3$.

Problem-3. Let a, b, c, m, n being positive integers and $f(x) = ax^2 + bx + c$, prove that there exist n consecutive integers $\alpha_1, \alpha_2, \dots, \alpha_n$ such that $f(\alpha_1), f(\alpha_2), \dots, f(\alpha_n)$ has at least m distinct prime factors.

Solution: we make an inductive reasoning on m , for $m = 1$ the statement of the problem holds, now choose $\alpha_1, \alpha_2, \dots, \alpha_n$ such that $f(\alpha_1), f(\alpha_2), \dots, f(\alpha_n)$ has at least m distinct prime divisors, now take $A = f^2(\alpha_1) \cdot f^2(\alpha_2) \cdot \dots \cdot f^2(\alpha_n)$ and $\beta_j = A + \alpha_j$, then $f(\beta_j) = f(\alpha_j) \cdot \underbrace{(1 + B \cdot f(\alpha_j))}_*$, for some integer B thus the $*$ term is coprime to $f(\alpha_j)$ and then has at least one prime number other than $f(\alpha_j)$. So we are done.

Problem-4. Let $\sigma(m) = \sum_{d|m, 0 < d < m} m$ prove that for every integer $t \geq 1$ there exist a positive integer m such that for all n , we have:

$$m < \sigma(m) < \sigma(\sigma(m)) < \dots < \sigma^{(t)}(m) \quad \text{AMM}$$

Solution: we prove stronger statement here:

For all positive integer n , there are a positive integer n_t such that for all positive integer m such that $\gcd\left(n_t, \frac{m}{n_t}\right) = 1$ (this means implicitly that $\frac{m}{n_t}$ is an integer) we have :

$$m < \sigma(m) < \sigma(\sigma(m)) < \dots < \sigma^{(t)}(m)$$

We prove this by induction on n , for $t = 1$, take $n_t = 12$. now suppose n_t has desired properties, we construct n_{t+1} as follows : take prime p not divide n_t , choose $k \geq 1$ such that $p^k \equiv 1 \pmod{n_t^2(p-1)}$, it is clear that such k exists (for example take $k = \varphi(n_t^2(p-1))$), we claim that $n_{t+1} = p^{k-1} \cdot n_t$, has desired property. set an integer m , such that $\gcd\left(n_{t+1}, \frac{m}{n_{t+1}}\right) = 1$ and let $A(m) = m + \sigma(m)$ it is obvious that $A(m)$ is multiplicative function. since $v_p(m) = k - 1$ then $A(p^{k-1}) = \frac{p^k - 1}{p - 1} A(m)$ so $n_t^2 \mid \frac{p^k - 1}{p - 1} A(m)$

Hence $\sigma(m) = A(m) - m \equiv -m \pmod{n_t^2}$, so $\gcd\left(n_t, \frac{\sigma(m)}{n_t}\right) = 1$ thus by induction hypothesis $\sigma(m)$ satisfies the condition we have $\sigma(m) < \sigma(\sigma(m)) < \dots < \sigma^{(t)}(\sigma(m)) = \sigma^{(t+1)}(m)$, at the same time, since $\gcd\left(n_t, \frac{m}{n_t}\right) = 1$ we have $m < \sigma(m)$ and we are done.

Problem-5. Let p being odd prime and a_1, \dots, a_{2p-1} being distinct numbers in the interval $[1, p^2]$ which their sum is divisible by p . prove that there exist positive integers b_1, \dots, b_{2p-1} none of which is divisible by p and their decimal representations in base p has only 0, 1 such that $\sum a_i b_i$ being divisible by p^{2012} .

Alexander Ivanov, Bulgarian Olympiads

Solution: we prove the statement for all positive integer n , instead of 2012, at first we prove following lemma :

Lemma: if x_1, \dots, x_{p-1} be positive integers not divisible by p , then for all $1 \leq r \leq p - 1$ we can choose some of them whose sum, leads to residue r modulo p .

Proof: set $S_0 = 0$ and for $0 \leq k \leq p - 2$ assume we receive the k different residues by numbers S_0, \dots, S_k , which made by x_1, \dots, x_k . add x_{k+1} to them we receive $x_{k+1} + S_0, \dots, x_{k+1} + S_k$, none of them has the same residue modulo p , if their residues being permutations of S_0, \dots, S_k then we must have $(k + 1)x_{k+1} \equiv 0 \pmod{p}$, which is impossible, so we receive at least new residue modulo p , so inductively we receive all non-zero residues, our proof is complete.

Back to our problem. we prove the statement by induction, since sum of a_i is divisible by p , the base is true, suppose there are positive integers b_1, \dots, b_{2p-1} none of which is divisible by p and their decimal representations in base p has only 0, 1 such that $\sum a_i b_i = p^n A$, Without loss of generality assume that A is not divisible by p . one can find that between a_1, \dots, a_{2p-1} there are at least $p - 1$ numbers say a_1, \dots, a_{p-1} , which is not divisible by p , now by the lemma there are a_1, \dots, a_k such that $a_1 + \dots + a_k + A \equiv 0 \pmod{p}$ now set $b'_i = b_i + p^n$ for $i = 1, \dots, k$ which satisfies the base- p criterion, and $b'_i = b_i$ for $i = k + 1, \dots, 2n - 1$. Then $\sum a_i b'_i$ is divisible by p^{n+1} . this completes our proof.

Problem-6. Prove that 2010 positive integers whose every sum of the integers selected from them is perfect power greater than unit.

Polish Olympiads

Solution: We prove inductively that for all there is a set S_k with such properties. for arbitrary set T , call a subset X of T good if sum of elements of this subset be a power number, for $S_1 = \{4\}$, the statement holds, assume there is a set S_k with desired properties, define $U_1 = S_1 \cup \{b\}$ if some subsets of U_1 wasn't good, we start to dis-eviling them, we construct the set U_i by multiplying elements of U_{i-1} by c_i . assume that in the set U_i there are t good subset which their sum is m_1, \dots, m_t power of integers, multiply all the elements by $c^{\text{lcm}(m_1, \dots, m_t)}$ now if sum of one non-good subset is c in U_i , now it is $c^{1+\text{lcm}(m_1, \dots, m_t)}$ in U_{i+1} , now the set U_{i+1} defines as $U_{i+1} = \{a \cdot c^{\text{lcm}(m_1, \dots, m_t)} \mid a \in U_i\}$ thus in each step the number of good subsets, increased, after finite steps say m , the set U_m has all subsets good, and then set $S_{k+1} = U_m$. So we are done.

Problem-7. Let C being a positive integer and p_1 being prime number and for all $n > 1$, p_n is the prime divisor of $p_{n-1} + C$. Which not enlisted in the sequence before, prove that sequence p_1, p_2, \dots is bounded. **Tuymada-2004**

Solution: consider, consecutive composite numbers $A = N(C + 1)! + 2, N(C + 1)! + 3, \dots, N(C + 1)! + C + 1$, and take natural number N , sufficiently large such that $A > p_1, C$. we prove by induction that for all natural number n , we must have $p_n < A$. for $n = 1$, it holds true, assume the statement true for natural numbers less than n but assume contrary that $p_n \geq A$, then $p_{n-1} + C \geq p_n \geq A$. since $p_{n-1} < A$ we find that $p_{n-1} + C$ will fall in the above composite sequence, so must be composite whence:

$$p_n \leq \frac{p_{n-1} + C}{2} < \frac{A + A}{2} = A$$

A contradiction, so for all integer n , we have $p_n < A$.

Problem-8. Prove that there exist positive integers $a_1 < a_2 < \dots < a_{2011}$, such that for all $1 \leq i < j \leq 2011$, such that $\gcd(a_i, a_j) = a_j - a_i$

Brazilian Olympiads

Solution: At first we prove that the condition of the problem is equivalent to $a_j - a_i | a_i$ (*), if $\gcd(a_i, a_j) = a_j - a_i$ then $a_j - a_i | a_i$ now if $a_j - a_i | a_i$ then $a_j - a_i | a_j$ so $a_j - a_i | \gcd(a_i, a_j)$ since $\gcd(a_i, a_j) \leq a_j - a_i$ we are done. Now we construct these numbers inductively , the case $n = 2$ is obvious , now assume we construct $x_1 < x_2 < \dots < x_{k-1}$ such that $\gcd(x_i, x_j) = x_j - x_i$ now we put k numbers

$$x_0 < x_0 + x_1 < x_0 + x_2 < \dots < x_0 + x_{k-1}$$

Where x_0 will be determined later, since the problem condition is equivalent to (*) we must have $x_j - x_i | x_0 + x_i$, $x_i | x_0$ take $x_0 = \text{lcm}(x_1, x_2, \dots, x_{k-1})$ and we prove the statement for $n = k$.

Problem-9. Let $a, m \geq 2$, $\gcd(a, m) = 1$ and $\text{ord}_m^a = k$, t being an odd number and every prime dividing t , also divides m , but $\gcd\left(t, \frac{a^k - 1}{m}\right) = 1$, prove that $\text{ord}_{mt}^a = kt$.

Bulgarian TST

Solution: we prove this problem by induction on number of prime factors of t , with their multiplicity . if t being prime , $d = \frac{a^k - 1}{m}$, $t | m$, set $a^k = 1 + md$, thus $a^{kt} = (1 + md)^t \equiv 1 \pmod{mt}$ thus $s = \text{ord}_{mt}^a | kt$ we know that $mt | a^s - 1$ thus $m | a^s - 1$ since $k = \text{ord}_m^a$ then $k | s | kt$ so $s = k$ or kt . if $s = k$ then $1 + md = a^k \equiv 1 \pmod{mt}$ this implies that $t | d$, a contradiction. So $s = kt$.

Now assume that t has at least to prime factor(not necessarily distinct) write $t = rt_0$ where r is prime and $t_0 > 1$ since r is prime by the use of base of the induction we find that $\text{ord}_{mr}^a = kr$. at first we prove $\gcd\left(t_0, \frac{a^{kr} - 1}{mr}\right) = 1$, then we will use the induction hypothesis . if prime number r_0 divides t_0 it also divides t so $r_0 | m | mr$, now lets regard : $d_0 = \frac{a^{kr} - 1}{mr} = \frac{a^k - 1}{m} \cdot \frac{a^{k(r-1)} + a^{k(r-2)} + \dots + 1}{r} = d \cdot \underbrace{\frac{a^{k(r-1)} + a^{k(r-2)} + \dots + 1}{r}}_c$ since $a^k \equiv 1 \pmod{m}$ then $a^k \equiv 1 \pmod{r}$ thus c is an integer , if $r \neq r_0$ then $a^k \equiv 1 \pmod{r_0}$ thus $c \equiv 1 \pmod{r_0}$ so $\gcd\left(t_0, \frac{a^{kr} - 1}{mr}\right) = 1$ and we can use induction hypothesis . if $r = r_0$ then one can set $a^k \equiv 1 + br \pmod{r^2}$ for some integer b , thus for all $j = 0, 1, \dots, r-1$ by use of binomial theorem we can find that $a^{kj} \equiv 1 + jbr \pmod{r^2}$.so

$$a^{k(r-1)} + a^{k(r-2)} + \dots + 1 \equiv r + br(1 + 2 + \dots + r - 1) \equiv r + \frac{b}{2}r^2(r - 1) \pmod{r^2}$$

Thus $c \equiv 1 \pmod{r}$ and again $\gcd\left(t_0, \frac{a^{kr} - 1}{mr}\right) = 1$, so our inductive reasoning is complete.

Problem-10. Prove that for all positive integers m, n there exist an integer k , such that $2^k - m$ has at least n distinct prime divisors.

Chinese TST

Solution: we prove the statement by induction on n , for $n = 1$ the statement is clear, assume now that there is an integer k_n such that $A_n = 2^{k_n} - m$ has at least n distinct prime divisors, WLOG, assume A_n is odd (we can easily preclude the power of 2 in prime decompositions of m , thus may assume m is odd), so $2^{k_n + \varphi(A_n^2)} - m \equiv A_n \equiv 2^{k_n} - m \pmod{A_n^2}$, whence $\frac{2^{k_n + \varphi(A_n^2)} - m}{A_n} \equiv 1 \pmod{A_n}$ so there exist a prime $p \nmid A_n$ but divides $\frac{2^{k_n + \varphi(A_n^2)} - m}{A_n}$, so we find at least $n + 1$ distinct prime divisors.

Problem-11. Let $p \geq 3$, being prime and for positive integers a_1, a_2, \dots, a_{p-2} . we know that neither $a_k, a_k^k - 1$ is not divisible by p where $k = 1, 2, \dots, p-2$. Prove that there is a positive integer m , such that $a_1 a_2 \dots a_m \equiv 2 \pmod{p}$.

Bulgarian Olympiads

Solution: we prove inductively, that for any $i = 2, 3, \dots, p-1$ there exist a sequence b_i which are the product of some of a_1, a_2, \dots, a_{p-2} such that for all $m \neq n$ we have $b_m \not\equiv b_n \pmod{p}$. for $i = 2$, take $b_1 = 1, b_2 = a_1$. And we are done. assume there are b_1, \dots, b_i such that for all $m \neq n$ we have $b_m \not\equiv b_n \pmod{p}$. consider numbers $b_1 a_i, \dots, b_i a_i$ by induction step none of which are congruent modulo p now if there are an index l such that $b_j a_i \equiv b_l \pmod{p}$ for any j then numbers $b_1 a_i, \dots, b_i a_i$ and b_1, \dots, b_i are same modulo p , comparing theirs product we have $a_i^i \equiv 1 \pmod{p}$, a contradiction. Thus we can adjoin new element to the set of incongruencies. Then we can say that for all $t = 2, \dots, p-1$ there are some product of a_i 's such that is congruent to t modulo p .

Strong Induction and Number theory

Problem-1. Prove that for all positive integer m , there exist an integer n , such that $\varphi(n) = m!$

Solution: we know that $3! = \varphi(18), 4! = \varphi(72)$, now take an induction, set $p_n \geq 5$, n -th prime number we know that for $k < p_n$, $k!$ wasn't divisible by p_n . We know that $(p_n - 2)! = \varphi(p_1^{e_1} \cdot p_2^{e_2} \dots p_{n-1}^{e_{n-1}})$, now $p_n! = p_n(p_n - 1) \cdot (p_n - 2)! = \varphi(p_n^2) \cdot \varphi(p_1^{e_1} \cdot p_2^{e_2} \dots p_{n-1}^{e_{n-1}}) = \varphi(p_1^{e_1} \cdot p_2^{e_2} \dots p_{n-1}^{e_{n-1}} \cdot p_n^2)$. now for all $p_n < m < p_{n+1} < 2p_n$, if $m = \prod_{p_i < p_n} p_i^{\alpha_i}$, then as $(m-1)!$ could be constructed before we can say that $m! = m \cdot (m-1)!$, now as $1 + p_n = p_1^{\beta_1} \cdot p_2^{\beta_2} \dots p_{n-1}^{\beta_{n-1}}$, we can construct $(1 + p_n)!$ by $(1 + p_n) \cdot p_n! = p_1^{\beta_1} \cdot p_2^{\beta_2} \dots p_{n-1}^{\beta_{n-1}} \varphi(p_1^{e_1} \cdot p_2^{e_2} \dots p_{n-1}^{e_{n-1}} \cdot p_n^2) = \varphi(p_1^{e_1 + \beta_1} \cdot p_2^{e_2 + \beta_2} \dots p_{n-1}^{e_{n-1} + \beta_{n-1}} \cdot p_n^2)$ by the same method we can construct all the numbers m , between (p_n, p_{n+1}) so by induction on prime numbers our proof is complete.

Problem-2. Let a, b, m being positive integers such that $\gcd(b, m) = 1$ prove that the set $\{a^n + bn \mid n = 1, 2, \dots, m^2\}$ contains complete residue system modulo m . **Z.We.Sun**

Solution: we prove the statement of the problem by induction on m but at first we divide the problem in two cases:

Case-1: $\gcd(m, ab) = 1$, the statement is clear for $m = 1$, assume $m = p_1^{\alpha_1} \dots p_t^{\alpha_t}$ where $p_1 < \dots < p_t$ are distinct prime numbers, set $m_0 = \frac{m}{p_t}$ thus by induction hypothesis there exist a number $k \leq m_0^2$, such that for all integer r , we have $a^k + bk = r + m_0 q_0$ for some integer q_0 since $\gcd(b(p_1 - 1) \dots (p_t - 1), p_t) = 1$ there exist a nonnegative integer $q < p_t$ such that $qb(p_1 - 1) \dots (p_t - 1) \equiv -q_0 \pmod{p_t}$.

Set $n = k + m_0 q(p_1 - 1) \dots (p_t - 1)$ and since $\varphi(m) \mid m_0(p_1 - 1) \dots (p_t - 1)$ then

$$a^n + bn \equiv a^k + b(k + m_0 q(p_1 - 1) \dots (p_t - 1)) \equiv r + m_0(q_0 + qb(p_1 - 1) \dots (p_t - 1)) \equiv r \pmod{m}$$

Now we can see that :

$$0 < k \leq n \leq m_0^2 + m_0(p_1 - 1) \dots (p_t - 1) < m_0^2 + m_0(p_t - 1)m = m_0^2(1 + p_t^2 - p_t) < m_0^2 p_t^2 = m^2$$

Case-2: $\gcd(m, a) > 1$, set $m = uv$ where $u > 1$ such that a is divisible by any prime divisors of u and $\gcd(v, a) = 1$. set r as arbitrary integer then there exist an integer $s \in \{0, 1, \dots, u - 1\}$ such that $bs \equiv r \pmod{u}$, we also know that $\gcd(a^u, v) = 1$ henceforth $\gcd((a^{-1})^s bu, v) = 1$ whence by previous case, there are $k \leq v^2$ such that $(a^u)^k + (a^{-1})^s buk \equiv (a^{-1})^s(r - bs) \pmod{v}$, now set $n = uk + s$ then $n \leq uv^2 + u - 1 < u(v^2 + 1) \leq u(u^2 + v^2) \leq u^2 v^2 = m^2$, and then :

$$a^{uk+s} + buk \equiv r - bs \pmod{v}$$

Implies that $a^n + bn \equiv r \pmod{v}$, now for any prime p divides a we know that $v_p(u) < u$ since $p^u \geq 2^u \geq u + 1$ so $a^n + bn = a^{uk+s} + b(uk + s) \equiv 0 + bs \equiv r \pmod{u}$. and our proof is complete.

Problem-3. Let $a^n - 1$ being divisible by n , prove that $a + 1, a^2 + 2, \dots, a^n + n$ is distinct modulo n . **Komal**

Solution: we will proceed induction for this, the case $n = 1$ is obvious lets suppose the statement holds true for all integers less than n , now assume $\text{ord}_n^a = k$ then $k \mid n$ and thus $k < n$, thus the statement of the problem is true for k , hence $a + 1, a^2 + 2, \dots, a^k + k$ are all distinct modulo k , now we prove that for $1 \leq x, y \leq n$ $a^x + x \not\equiv a^y + y$, let $x = kz + t, y = ku + v$ where $0 \leq t, v \leq k - 1, 0 \leq z, u \leq \frac{n}{k}$ then $a^x \equiv a^t, a^y \equiv a^v \pmod{n}$ now we have two cases:

Case-1: $t \neq v$. then $a^x + x \equiv a^t + t \not\equiv a^v + v \equiv a^y + y \pmod{k}$ the central incongruence is derived from induction hypothesis.

Case-2: $= v$, then we essentially have $z \neq u$, so

$$a^x + x \equiv a^t + kz + t = a^v + ku + v + k(z - u) \equiv a^y + y + k(z - u) \not\equiv a^y + y \pmod{n}$$

Our proof is complete.

Problem-4. Prove that for all positive integer n , there exist an integer k , such that $3^{k+1} - 2^k - k$ is divisible by n .

Serbia

Solution: define the sequence $x_0 = 2, x_{n+1} = 3^{x_n} - 2^{x_n}$ we prove the following lemma:

Lemma: for all d the congruence $x_{n+1} \equiv x_n \pmod{d}$ holds for all sufficiently large n .

Proof: we prove this lemma by induction on d , for $d = 1$ it is obvious, assume that the statement holds for all integers less than d , since $\varphi(d) < d$ the statement holds for $\varphi(d)$, ensure that $x_{n+1} \equiv x_n \pmod{\varphi(d)}$ for all sufficiently large n , thus $3^{x_{n+1}} \equiv 3^{x_n}, 2^{x_{n+1}} \equiv 2^{x_n} \pmod{d}$. so our lemma is proven.

Back to our problem, we set $k = x_n, d = m$ and we are done.

$n, \frac{n}{p}$ -Technique

Problem-1. We know about the sequence a_n such that $a_1 = 1, a_n = a_{\lfloor \frac{n}{2} \rfloor} + a_{\lfloor \frac{n}{3} \rfloor} + \dots + a_{\lfloor \frac{n}{n} \rfloor} + 1$, prove that there exist infinitely many n such that $a_n \equiv n \pmod{2^{2010}}$ **USA-TST-2010.**

Solution: first we prove the following lemma:

Lemma: if $v_p(n) \geq s$, then $2^{s-1} | a_n - a_{n-1}$.

Proof: assume $a_0 = 0$, the statement is obvious for $s = 1$. assume that the statement holds true for all integers less than n , we will prove the truth of the proposition for n , assume contrary so n is the least integer such that $v_p(n) \geq s$, but $2^{s-1} \nmid a_n - a_{n-1}$, take $b_n = a_n - a_{n-1}$, then one can find that:

$$b_n = \sum_{i>1, i|n} \frac{b_n}{i} = \sum_{i<n, i|n} b_i$$

Thus $\frac{b_n}{p} = \sum_{i<\frac{n}{p}, i|\frac{n}{p}} b_i$ so $b_n = \frac{b_n}{p} + \sum_{i<\frac{n}{p}, i|\frac{n}{p}} b_i + \sum_{i<n, i|\frac{n}{p}, i|n} b_i = 2\frac{b_n}{p} + \sum_{i<n, i|\frac{n}{p}, i|n} b_i$

Since $\frac{n}{p} < n$ and $p^{s-1} \mid \frac{n}{p}$ thus by induction hypothesis, we must have $2^{s-2} \mid \frac{n}{p}$ so $2^{s-1} \mid 2\frac{n}{p}$. now those $i < n$ such that $i \nmid \frac{n}{p}$ but $i \mid n$ are divisible by p^s so by induction hypothesis we must have $2^{s-1} \mid b_i$, then we must have $2^{s-1} \mid b_n$. a contradiction, so Our proof is complete.

Let p_1, \dots, p_m be different primes, by Chinese remainder theorem we can find k such that for all $1 \leq i \leq m$ we have $p_i^s \mid k + i$ so by our lemma $2^{s-1} \mid a_{k+i} - a_{k+i-1}$ so $a_k \equiv a_{k+1} \equiv \dots \equiv a_{k+m} \equiv N$ (say!) (mod 2^{s-1}) take $> N + 2^{s-1} + 1$. Then since the set $\{k + N, \dots, k + N + 2^{s-1} - 1\}$ covers all residues modulo 2^{s-1} , there exist n belongs to the above set, such that $a_n \equiv n \equiv N$ (mod 2^{s-1}), $n > N$.

Problem-2. For all, prove that there exist an integer n with at least m prime factors (no necessarily distinct) such that $2^{kn^2} + 3^{kn^2}$ being divisible by n^3 . Where k is a positive integer.

Solution: at first we present the following lemma:

Lemma: let $\gcd(a, b) = 1$ being integers and p be an odd prime such that $v_p(a + b) = s \geq 1$, then $v_p(a^p + b^p) = s + 1$.

Proof: set $a + b = x$ is divisible by p , now

$$\frac{a^p + b^p}{a + b} = \frac{(x-b)^p + b^p}{x} = x^{p-1} - bp x^{p-2} + \dots - \binom{p}{2} b^{p-2} x + p b^{p-1} \equiv p b^{p-1} \pmod{p^2}$$

Since $v_p(a^p + b^p) = v_p(a + b) + v_p\left(\frac{a^p + b^p}{a + b}\right)$ we are done.

we prove the statement by induction on m , for $m = 0$ take $n_1 = 1$, assume that for $m - 1$ there are an integer n_i with at least $m - 1$ prime factors such that n_i^3 divides $2^{kn_i^2} + 3^{kn_i^2}$, now we divide the problem in two case:

Case-1: there exist a prime p such that $p \nmid n_i$ but $2^{kn_i^2} + 3^{kn_i^2}$ being divisible by p . since $p \mid 2^{kn_i^2} + 3^{kn_i^2}$ by our lemma $p^3 \mid 2^{kp^2 n_i^2} + 3^{kp^2 n_i^2}$ now, take $n_{i+1} = p n_i$, we are done.

Case-2: n_i and $2^{kn_i^2} + 3^{kn_i^2}$ has same prime factors. we know that $n_i^3 \neq 2^{kn_i^2} + 3^{kn_i^2}$ otherwise $2^{kn_i^2} + 3^{kn_i^2} > 3^{n_i^2} > n_i^3$ thus there exist a prime q such that $v_q(n_i^3) = \alpha$, $v_q(2^{kn_i^2} + 3^{kn_i^2}) = \beta$ and $\beta \geq 1 + \alpha$, take $n_{i+1} = q n_i$ so $q^{\alpha+3} \mid q^{\beta+2} = v_q(2^{kn_{i+1}^2} + 3^{kn_{i+1}^2})$

Problem-3. Prove that for all positive integer k , there exist an integer n Which has exactly k prime divisors and $\frac{2^{n^2}+1}{n^3}$ being an integer.

Solution: we first prove two important lemma:

Lemma-1: Let a be an integer, two following statements are equivalent:

- a. $v_p(a+1) = s \geq 1$
- b. $v_p(a^p+1) = s+1$

Proof: if $v_p(a+1) = s \geq 1$, then $\frac{a^p+1}{a+1} \equiv p \pmod{p^2}$, since $v_p(a^p+1) = v_p(a+1) + v_p\left(\frac{a^p+1}{a+1}\right)$ we are done, on the other hand, since $p|a^p+1$, by fermat's little theorem we find that $a+1$ is divisible by p and by the above proof, we are done.

Lemma-2: Let a be a positive integer, then there exist prime q such that $q|\frac{a^p+1}{a+1}$ but $q \nmid a+1$ except the case $a=2, p=3$.

Proof: assume contrary then all of theirs common primes will appear in theirs greatest prime divisors, since $\gcd\left(\frac{a^p+1}{a+1}, a+1\right) = 1$ or p , we find that $\frac{a^p+1}{a+1}$ must be power of p and at the simultaneously with $a+1$, since by the previous lemma, we find that if $p|a+1$, then $v_p\left(\frac{a^p+1}{a+1}\right) = 1$, we find that $\frac{a^p+1}{a+1} = p$ but $\frac{a^p+1}{a+1} > a^2 - a + 1 > a+1 \geq p$ except the case $a=2, p=3$. Our proof is complete.

Back to our problem, by lemma-1, we find that, if $p|2^m+1$ then $p^3|2^{mp^2}+1$. now take $n_1 = p_1 = 3$ since $2^9+1 = 513 = 27 \cdot 19$, take $p_2 = 19$, $n_2 = p_1 p_2$, we construct n_i inductively, assume we construct $n_k = p_1 p_2 \dots p_k$ by lemma-2, there are prime p_{k+1} such that $p_{k+1} \nmid 2^{p_1 p_2 \dots p_{k-1}} + 1 = a+1$ but $p_{k+1} | 2^{p_1 p_2 \dots p_k} + 1 = a^{p_k} + 1$, now set $n_{k+1} = p_{k+1} n_k$. since $n_k^3 | 2^{n_k^2} + 1$ and by lemma-1, $p_{k+1}^3 | 2^{n_k^2 \cdot p_{k+1}^2} + 1$ we find that $n_k^3 p_{k+1}^3 = n_{k+1}^3 | 2^{n_k^2 \cdot p_{k+1}^2} + 1 = 2^{n_{k+1}^2} + 1$. so we are done.

Problem-4. We know about the sequence a_n , that $\sum_{d|n} da_d = k^n$ where k is a positive integer , prove that terms of this sequence are all positive integers a_n is an integer.

Polish training camps.

Solution: we prove the statement by induction on n , for $n = 1$ it is obvious , assume it was true for all integers less than n , write the equality as $n \cdot a_n + \sum_{d|n, d < n} da_d = k^n$ we will prove that $n|k^n - \sum_{d|n, d < n} da_d$ which leads to the desired conclusion. Assume that $v_p(n) = r$, $n = p^r x$ for some integer x which is not divisible by p we prove that above summation is divisible by p^r , thus by use of induction hypothesis we can write:

$$k^n - \sum_{d|n, d < n} da_d \equiv k^n - \sum_{d|p^{r-1}x} da_d \equiv k^{p^r x} - k^{p^{r-1}x} \equiv k^{p^{r-1}x}(k^{p^{r-1}(p-1)x} - 1) \pmod{p^r}$$

Now if $p|k$ we are done , otherwise by Euler's theorem the term in the bracket is divisible by p^r . Since the above proof is indeed for all primes dividing n , we are done.

Miscellaneous

Problem-I. For $n = 1, 2, 3$ n -type-number include the following numbers:

- a. Zero
- b. Geometric progression $1, n + 2, (n + 2)^2, \dots$
- c. Sum of its various numbers.

Prove that every natural number could be represented as the sum of first type , second type and the third type number.

Moscow-2012

Solution: first we prove the following lemma

Lemma: let $a_1, a_2, \dots, a_n, \dots$ be a non-decreasing sequence of positive integers and define $S_1 = 0$, for ≥ 2 . $S_k = a_1 + \dots + a_{k-1}$, then any positive integers could be written as sum of different terms of the sequence $a_1, a_2, \dots, a_n, \dots$ if and only if: $a_k \leq 1 + S_k$.

Proof: if for some k , we have $a_k > 1 + S_k$ then the number $1 + S_k$ can't be represented by that form. For the necessity proof , we use induction assume that the statement of the problem , holds true for all the integers in the interval $[1, S_k]$ using only the numbers included in S_k , i.e. a_1, \dots, a_{k-1} . for the base of induction since $a_1 \leq 1 + S_1 = 1$, we have $S_2 = a_1 = 1$. we are done for $k = 1, 2$. Assume we can write $1, 2, \dots, S_k$ without a_k , add a_k to $1, 2, \dots, S_k$, we receive $1 + a_k, 2 + a_k, \dots, S_k + a_k = S_{k+1}$, whence We can construct all of them . since $a_k \leq 1 + S_k$. By induction hypothesis we can construct all the numbers in the interval $[1, S_{k+1}]$, so we are done.

Now we reorder the three type numbers , inclusively as :

$$1,1,1,3,4,5,9,16,25,27, \dots$$

So $a_1 = a_2 = a_3 = 1$,we may prove that the inequality $a_k \leq 1 + S_k$ holds for $k > 3$. assume that in the first k terms of the sequence we have $(3^0, \dots, 3^{n-1}), (4^0, \dots, 4^{m-1}), (5^0, \dots, 5^{l-1})$ where $l = k - 1 - n - m$ so we have

$$\begin{aligned} S_k &= \sum_{j=0}^{n-1} 3^j + \sum_{j=0}^{m-1} 4^j + \sum_{j=0}^{l-1} 5^j = \frac{3^n-1}{2} + \frac{4^m-1}{3} + \frac{5^l-1}{4} \geq (\min(3^n, 4^m, 5^l) - 1) \left(\frac{1}{2} + \frac{1}{3} + \frac{1}{4} \right) \\ &= \frac{13}{12} (\min(3^n, 4^m, 5^l) - 1) > (\min(3^n, 4^m, 5^l) - 1) = a_k - 1 \end{aligned}$$

And we are done.

Problem-2. Let $0 < x_k \leq \frac{1}{2}$ prove that $\left(\frac{n}{x_1+x_2+\dots+x_n} - 1 \right)^n \leq \left(\frac{1}{x_1} - 1 \right) \dots \left(\frac{1}{x_n} - 1 \right)$ **Tuymada-2000**

First Solution: we prove the following lemmaL

Lemma: that if x, y being two positive reals such that $x+y \leq 1$, then we have:

$$\left(\frac{1}{x} - 1 \right) \left(\frac{1}{y} - 1 \right) \geq \left(\frac{2}{x+y} - 1 \right)^2$$

Proof: rewrite the inequality in the form $\frac{1-x-y}{xy} + 1 \geq \frac{1-x-y}{(\frac{x+y}{2})^2} + 1$ which is true by the AM-GM Inequality $\leq \left(\frac{x+y}{2} \right)^2$.

now we can prove inductively (on m) that for all $N = 2^m$ real numbers x_1, \dots, x_N such that $0 < x_k \leq \frac{1}{2}$, for $1 \leq k \leq N$ we have $\left(\frac{N}{x_1+x_2+\dots+x_N} - 1 \right)^N \leq \left(\frac{1}{x_1} - 1 \right) \dots \left(\frac{1}{x_N} - 1 \right)$, now by back-ward induction we prove the inequality for all n , assume ,we want to prove the inequality for $n < N = 2^m$, numbers x_1, \dots, x_n $0 < x_k \leq \frac{1}{2}$, and note that $x_1 + \dots + x_n = nd$,for some positive real d , since the inequality is true for any N , numbers , take specific(Particular) N , numbers as : $x_1, \dots, x_n, x_{n+1} = \dots = x_N = d$, notice that $x_1 + \dots + x_N = nd + (N - n)d = Nd$.

so we write the statement as :

$$\left(\frac{N}{x_1 + x_2 + \dots + x_N} - 1\right)^N \leq \left(\frac{1}{x_1} - 1\right) \dots \left(\frac{1}{x_N} - 1\right)$$

Whence the inequality leads to the following statement:

$$\begin{aligned} \left(\frac{N}{x_1 + x_2 + \dots + x_N} - 1\right)^N &= \left(\frac{N}{Nd} - 1\right)^N = \left(\frac{1}{d} - 1\right)^N \leq \left(\frac{1}{x_1} - 1\right) \dots \left(\frac{1}{x_N} - 1\right) = \\ &\left(\frac{1}{x_1} - 1\right) \dots \left(\frac{1}{x_n} - 1\right) \cdot \left(\frac{1}{d} - 1\right)^{N-n} \end{aligned}$$

So one can find that $\left(\frac{1}{d} - 1\right)^n = \left(\frac{n}{x_1 + x_2 + \dots + x_n} - 1\right)^n \leq \left(\frac{1}{x_1} - 1\right) \dots \left(\frac{1}{x_n} - 1\right)$.

We are done.

Second Solution: take $x_1 + \dots + x_n = nd$ as past, assume there are to index, say: x_1, x_2 such that $x_1 < d, x_2 > d$. now we prove the following lemma:

Lemma: let $x_1 < d, x_2 > d$ then: $\left(\frac{1}{x_1} - 1\right)\left(\frac{1}{x_2} - 1\right) > \left(\frac{1}{d} - 1\right)\left(\frac{1}{x_1 + x_2 - d} - 1\right)$

Proof: by simplifying both sides we get $(1 - x_1 - x_2)(d - x_1)(x_2 - d) > 0$. and our proof is complete.

Now, if we change the set $\{x_1, \dots, x_n\}$ by $\{d, x_1 + x_2 - d, x_3, \dots, x_n\}$, the numbers which isn't equal to d , decreased by one, we continue these procedure (which takes finitely) and final all the numbers are equal to d and thus their products is greater than $\left(\frac{1}{d} - 1\right)^n$.

Comment: the following hard inequality, is indeed the particular case of the above inequality:

Let a_1, a_2, \dots, a_5 being positive real numbers, prove that:

$$\begin{aligned} \frac{a_1 + a_2}{2} \cdot \frac{a_2 + a_3}{2} \cdot \frac{a_3 + a_4}{2} \cdot \frac{a_4 + a_5}{2} \cdot \frac{a_5 + a_1}{2} \\ \leq \frac{a_1 + a_2 + a_3}{3} \cdot \frac{a_2 + a_3 + a_4}{2} \cdot \frac{a_3 + a_4 + a_5}{3} \cdot \frac{a_4 + a_5 + a_1}{3} \end{aligned}$$

Just take $S = a_1 + a_2 + a_3 + a_4 + a_5$, and take $a_k = a_{k-5}$, then $\frac{a_{k+2} + a_{k+3} + a_{k+4}}{a_{k+1} + a_{k+2}} = \frac{S}{a_{k+1} + a_{k+2}} - 1$

,hence the problem leads to proving $\left(\frac{S}{a_1 + a_2} - 1\right) \dots \left(\frac{S}{a_1 + a_5} - 1\right) \geq \left(\frac{3}{2}\right)^5$, we know $\frac{a_k + a_{k+1}}{S} + \frac{a_{k+2} + a_{k+3}}{S} \leq 1$ then:

$$\left(\frac{S}{a_k + a_{k+1}} - 1\right) \left(\frac{S}{a_{k+2} + a_{k+3}} - 1\right) \geq \left(\frac{2S}{a_k + a_{k+1} + a_{k+2} + a_{k+3}} - 1\right)^2 = \left(\frac{2S}{S - a_{k+4}} - 1\right)^2 = \left(\frac{2S}{S - a_{k-1}} - 1\right)^2$$

Now we must prove $\prod_{i=1}^5 \left(\frac{2S}{S - a_i} - 1\right) \geq \left(\frac{3}{2}\right)^5$, we set $x_k = \frac{S - a_k}{2S} \leq \frac{1}{2}$. And $x_1 + \dots + x_5 = 2$ implies that:

$$\left(\frac{5}{x_1+x_2+\dots+x_5} - 1\right)^5 = \left(\frac{3}{2}\right)^5 \leq \left(\frac{1}{x_1} - 1\right) \dots \left(\frac{1}{x_5} - 1\right)$$

Variant:

Let $p = \sqrt[n]{x_1 \dots x_n}$, prove that $\left(1 + \frac{1}{x_1(x_1+1)}\right) \dots \left(1 + \frac{1}{x_n(x_n+1)}\right) \geq \left(1 + \frac{1}{p(1+p)}\right)^n$.

Solution: first we prove the following lemma:

Lemma: Let $p = \sqrt{xy}$ and x, y being positive real numbers then:

$$\left(1 + \frac{1}{x(1+x)}\right) \left(1 + \frac{1}{y(1+y)}\right) \geq \left(1 + \frac{1}{p(1+p)}\right)^2$$

Solution: the left hand side of the inequality is equal to: $1 + \frac{x^2+x+y^2+y}{xy(1+x)(1+y)} = 1 + \frac{1}{xy} + \frac{x^2-xy+y^2}{xy(1+x)(1+y)}$ now we must prove $\frac{x^2-xy+y^2}{p^2(1+x)(1+y)} \geq \frac{1}{(1+p)^2}$, set $x + y = s$, we must prove $(s^2 - 3p^2)(p+1)^2 \geq (p^2 + s + 1)p^2$, since $s \geq 2p$ the inequality is proven. Now by induction the statement holds for all powers of 2, now by back-ward induction we prove the inequality for all n , assume we want to prove the inequality for $n < N = 2^m$, numbers x_1, \dots, x_n such that $x_1 \dots x_n = p^n$, since the inequality is true for any N , numbers, take specific (Particular) N , numbers as: $x_1, \dots, x_n, x_{n+1} = \dots = x_N = p$, whence $x_1 \dots x_N = p^N$, thus:

$$\begin{aligned} & \left(1 + \frac{1}{x_1(x_1+1)}\right) \dots \left(1 + \frac{1}{x_N(x_N+1)}\right) \\ &= \left(1 + \frac{1}{x_1(x_1+1)}\right) \dots \left(1 + \frac{1}{x_n(x_n+1)}\right) \cdot \left(1 + \frac{1}{p(1+p)}\right)^{N-n} \\ &\geq \left(1 + \frac{1}{p(1+p)}\right)^N \end{aligned}$$

So our proof is complete.

Problem-3. Let a_1, \dots, a_n being real numbers, prove that two following statements are equivalent:

- $a_i + a_j \geq 0$, for all $i \neq j$.
- If x_1, x_2, \dots, x_n being non-negative real numbers adding up to one, then:

$$a_1 x_1 + \dots + a_n x_n \geq a_1 x_1^2 + \dots + a_n x_n^2$$

Chinese Olympiads

Solution: if b, being true, set for $1 \leq i \leq j \leq n$ $x_i = x_j = \frac{1}{2}$, and otherwise, $x_k = 0$ then, $\frac{a_i+a_j}{2} \geq \frac{a_i+a_j}{4}$ and we are done. Now assume a is true, we prove that a implies b, by induction on n , if $n = 2$, then $a_1 x_1 + a_2 x_2 - a_1 x_1^2 - a_2 x_2^2 = (a_1 + a_2)x_1 x_2 \geq 0$. and we are done. assume the statements holds for all $3, \dots, n$. and let $x_1, x_2, + \dots, x_{n+1}$ being

non-negative real numbers adding up to one , if $x_{n+1} = 1$ we are done , if $x_{n+1} < 1$, then $\sum_{i=1}^n \frac{x_i}{1-x_{n+1}} = 1$ then by induction hypothesis we have $\sum_{k=1}^n \frac{a_k x_k}{1-x_{n+1}} \geq \sum_{k=1}^n a_k \left(\frac{x_k}{1-x_{n+1}}\right)^2$ or

$$(1 - x_{n+1}) \sum_{k=1}^n a_k x_k \geq \sum_{k=1}^n a_k x_k^2 .$$

We know that

$$\sum_{k=1}^{n+1} a_k x_k = (1 - x_{n+1}) \sum_{k=1}^n a_k x_k + x_{n+1} \sum_{k=1}^n a_k x_k + (1 - x_{n+1}) a_{n+1} x_{n+1} + a_{n+1} x_{n+1}^2 \geq \sum_{k=1}^{n+1} a_k x_k^2 + x_{n+1} \sum_{k=1}^n a_k x_k + a_{n+1} x_{n+1} \sum_{k=1}^n x_k \geq \sum_{k=1}^{n+1} a_k x_k^2$$

Since $(a_k + a_{n+1})x_k x_{n+1} \geq 0$ we are done.