

Branching processes with immigration in a random environment

Péter Kevei

University of Szeged

ISCPS 2025

Outline

Introduction

- GWI in deterministic environment
- Random environment

Results

- Tail asymptotic
- Stationary Markov chain

Stochastic recurrence equation

- Perpetuity equation
- Goldie's implicit renewal theory

Outline

Introduction

GWI in deterministic environment

Random environment

Results

Tail asymptotic

Stationary Markov chain

Stochastic recurrence equation

Perpetuity equation

Goldie's implicit renewal theory

GWI

Let $X_0 = 0$,

$$X_{n+1} = \sum_{i=1}^{X_n} A_i^{(n+1)} + B_{n+1} =: \theta_{n+1} \circ X_n + B_{n+1}, \quad n \geq 0,$$

offsprings $\{A_i^{(n)} : i = 1, 2, \dots, n = 1, 2, \dots\}$ iid, immigrants

$\{B_n : n = 1, 2, \dots\}$ iid.

Subcritical, critical, supercritical: $\mathbf{EA} < 1, = 1, > 1$.

Stationary distribution – existence

Theorem (Quine (1970), Foster & Williamson (1971))

Unique stationary distribution exists iff

$$\int_0^1 \frac{1 - \mathbf{E}s^B}{\mathbf{E}s^A - s} ds < \infty.$$

$$X_\infty = B_1 + \theta_1 \circ B_2 + \theta_1 \circ \theta_2 \circ B_3 + \dots = \sum_{i=0}^{\infty} \Pi_i \circ B_{i+1}.$$

If $m = \mathbf{E}A < 1$, then $\mathbf{E} \log B < \infty$ is necessary and sufficient.

If $m = 1$, then the condition holds if $\mathbf{P}(A > n) \sim \ell_A(n)n^{-1-\alpha}$,

$\mathbf{E}B < \infty$, $\alpha \in (0, 1)$, or $\mathbf{P}(B > n) \sim \ell_B(n)n^{-\beta}$, $\beta > \alpha$.

Stationary distribution – tail

Theorem (Basrak & Kulik & Palmowski (2013))

- (i) If $m = \mathbf{E}A < 1$, $\mathbf{E}A^2 < \infty$, and $\mathbf{P}(B > x)$ is regularly varying with index $-\alpha \in (-2, 0)$, then

$$\mathbf{P}(X_\infty > x) \sim c \mathbf{P}(B > x), \quad c > 0.$$

- (ii) If $m = \mathbf{E}A < 1$, $\mathbf{P}(A > x)$ is regularly varying with index $-\alpha \in (-2, -1)$, and $\mathbf{P}(B > x) \sim c' \mathbf{P}(A > x)$, $c' \geq 0$ then

$$\mathbf{P}(X_\infty > x) \sim c \mathbf{P}(A > x), \quad c > 0.$$

More general tail behavior: Foss & Miyazawa (2020)

Second order GWI: Barczy & Bősze & Pap (2020)

Critical case, $m = 1$

Theorem (Guo & Hong (2024))

Assume $\mathbf{P}(A > n) \sim \ell_A(n)n^{-1-\alpha}$, $\alpha \in (0, 1)$,
 $\mathbf{P}(B > n) \sim \ell_B(n)n^{-\beta}$, $\beta > \alpha$, additional assumption on ℓ_A , ℓ_B .
Then

$$\mathbf{P}(X_\infty > x) \sim \ell(x)x^{-(\beta-\alpha)}.$$

Without immigration: Fleischmann & Vatutin & Wachtel (2008):

$\mathbf{P}(W > n) \sim \ell(n)n^{-1/(1+\alpha)}$, W : total progeny

Borovkov & Vatutin (1996): $\mathbf{P}(M > x) \sim \frac{\alpha}{x}$

Critical case, $m = 1$

Assume $\mathbf{P}(A > n) \sim \ell_A(n)n^{-1-\alpha}$, $\alpha \in (0, 1)$, $\mathbf{E}B < \infty$.

K & Kubatovics (2025+, work in progress):

$$\mathbf{P}(X_\infty > x) \sim \ell(x)x^{-(1-\alpha)}.$$

Stationary chain $(X_n)_{n \geq 0}$ is regularly varying, tail process:

$$\mathcal{L}((X_n/x)|X_0 > x) \longrightarrow Y(1, 1, \dots),$$

$\mathbf{P}(Y > y) = y^{-(1-\alpha)}$. Extremal index = 0. The anticlustering condition does not hold.

Explicit calculations are possible: Alsmeyer & Hoang (2025):

Power fractional distributions, Lindo & Sagitov (2016):

θ -branching

Outline

Introduction

GWl in deterministic environment

Random environment

Results

Tail asymptotic

Stationary Markov chain

Stochastic recurrence equation

Perpetuity equation

Goldie's implicit renewal theory

GWRE with immigration (GWIRE)

- ▶ Δ probability measures on $\mathbb{N} = \{0, 1, \dots\}$
- ▶ ξ, ξ_1, \dots iid on Δ^2 (environment), $\xi = (\nu_\xi, \nu_\xi^\circ)$

GWRE with immigration (GWIRE)

- ▶ Δ probability measures on $\mathbb{N} = \{0, 1, \dots\}$
- ▶ ξ, ξ_1, \dots iid on Δ^2 (environment), $\xi = (\nu_\xi, \nu_\xi^\circ)$
- ▶ $X_0 = 0$,

$$X_{n+1} = \sum_{i=1}^{X_n} A_i^{(n+1)} + B_{n+1} =: \theta_{n+1} \circ X_n + B_{n+1}, \quad n \geq 0,$$

conditioned on \mathcal{E} , $\{A_i^{(n)}, B_n : i = 1, 2, \dots, n = 1, 2, \dots\}$ are independent and for n fix $(A_i^{(n)})_{i=1,2,\dots}$ are iid with distribution ν_{ξ_n} , and B_n has distribution $\nu_{\xi_n}^\circ$.

Subcritical / critical / supercritical: $\mathbf{E} \log m(\xi) < / = / > 0$.

Kersting, Vatutin: Discrete Time Branching Processes in Random Environment, 2017, Wiley.

Stationary distribution – existence

Theorem (Key (1987))

If $\mathbf{E} \log m(\xi) < 0$ (offspring) and $\mathbf{E} \log^+ m^\circ(\xi) < \infty$ (immigration) then there exists a unique stationary distribution

$$X_\infty = B_1 + \theta_1 \circ B_2 + \theta_1 \circ \theta_2 \circ B_3 + \dots = \sum_{i=0}^{\infty} \Pi_i \circ B_{i+1}.$$

Outline

Introduction

GWI in deterministic environment

Random environment

Results

Tail asymptotic

Stationary Markov chain

Stochastic recurrence equation

Perpetuity equation

Goldie's implicit renewal theory

Kesten–Grincevičius–Goldie setup

$$X_\infty = B_1 + \theta_1 \circ B_2 + \theta_1 \circ \theta_2 \circ B_3 + \dots = \sum_{i=0}^{\infty} \Pi_i \circ B_{i+1}.$$

Theorem (Basrak & K 2022)

*Assume: $\mathbf{E}m(\xi)^\kappa = 1$, $\mathbf{E}A^\kappa < \infty$, $\mathbf{E}B^\kappa < \infty$,
 $\mathbf{E}m(\xi)^\kappa \log m(\xi) < \infty$, $\log m(\xi)$ is non-arithmetic. Then*

$$\mathbf{P}(X_\infty > x) \sim Cx^{-\kappa} \quad x \rightarrow \infty,$$

with $C > 0$.

Related papers

- ▶ Afanasyev (2001): $\mathbf{P}(\sup_n X_n > x) \sim cx^{-\kappa}$, $c > 0$.
- ▶ Large deviation results: Buraczewski & Dyszewski (2022), Guo & Hong & Sun (2025)
- ▶ Arithmetic case: Jelenković and Olvera-Cravioto (2012), K (2017): implicit renewal theory in the arithmetic case

Grincevičius – Grey setup

Theorem (K 2024)

Assume: $\mathbf{E}(m(\xi)^{1 \vee \kappa}) < 1$, $\mathbf{E}(A^{(1 \vee \kappa) + \delta}) < \infty$.

Let ℓ be a slowly varying function. Then

$$\mathbf{P}(B > x) \sim \frac{\ell(x)}{x^\kappa}, \quad \text{as } x \rightarrow \infty,$$

if and only if

$$\mathbf{P}(X_\infty > x) \sim \frac{\ell(x)}{x^\kappa} \frac{1}{1 - \mathbf{E}(m(\xi)^\kappa)}, \quad \text{as } x \rightarrow \infty.$$

Valid in the deterministic environment setup – Basrak, Kulik, Palmowski (2013); Foss, Miyazawa (2020)

Outline

Introduction

GWI in deterministic environment

Random environment

Results

Tail asymptotic

Stationary Markov chain

Stochastic recurrence equation

Perpetuity equation

Goldie's implicit renewal theory

Setup

- ▶ $X_{n+1} = \sum_{i=1}^{X_n} A_i^{(n+1)} + B_{n+1} =: \theta_{n+1} \circ X_n + B_{n+1}$, $n \in \mathbb{Z}$, strictly stationary
- ▶ $\mathbf{P}(X_0 > x) \sim c\ell(x)x^{-\kappa}$
- ▶ a_n is defined by $n\mathbf{P}(X_0 > a_n) \sim 1$.

Kulik, Soulier (2020): Heavy-tailed time series

Mikosch, Wintenberger (2024): Extreme value theory for time series

Asymptotic properties of Markov chain

- ▶ tail process (Basrak & Segers (2009))
- ▶ point process convergence (ergodicity, anticlustering)
- ▶ convergence of partial sums (vanishing small values)

Stable limit for partial sums

Theorem (Basrak & K 2022)

Let $b_n = 0$, $\kappa < 1$, $b_n = n\mathbf{E}(X_\infty/a_n I(X_\infty \leq a_n))$, $\kappa \in [1, 2)$. Then

$$V_n = \sum_{k=1}^n \frac{X_k}{a_n} - b_n \xrightarrow{\mathcal{D}} V, \quad n \rightarrow \infty,$$

with V κ -stable. If $\kappa > 2$,

$$\frac{1}{\sqrt{n}\sigma} \sum_{j=1}^n (X_j - \mathbf{E}X_\infty) \xrightarrow{\mathcal{D}} Z \sim N(0, 1).$$


Random walk in random environment

Kozlov and Solomon:

- ▶ $\{\alpha_i\}_{i \in \mathbb{Z}}$ iid in $[0, 1]$
- ▶ $\mathcal{A} = \sigma(\alpha_i : i \in \mathbb{Z})$, σ -algebra generated by the environment,
- ▶ $X_0 = 0$,

$$\mathbf{P}(W_{n+1} = W_n + 1 | \mathcal{A}, W_0, \dots, W_n) = \alpha_i \text{ on } \{W_n = i\}$$

$$\mathbf{P}(W_{n+1} = W_n - 1 | \mathcal{A}, W_0, \dots, W_n) = 1 - \alpha_i \text{ on } \{W_n = i\}$$



KKS result

Let T_n be the first hitting time of n . $T_n \approx 2 \sum_{k=1}^n X_k - n$.

KKS result

Let T_n be the first hitting time of n . $T_n \approx 2 \sum_{k=1}^n X_k - n$.

Theorem (Kesten & Kozlov & Spitzer 1975)

For $\kappa \in (0, 2)$,

$$n^{-1/\kappa}(T_n - A_n) \xrightarrow{\mathcal{D}} \kappa - \text{stable r.v.}$$

where $A_n \equiv 0$ if $\kappa < 1$, $A_n = nc_1$ if $\kappa > 1$.

For $\kappa > 2$

$$n^{-1/2}(T_n - nc) \xrightarrow{\mathcal{D}} N(0, 1).$$

Moreover, $n^{-\kappa}(W_n - B_n)$ also converges.

Outline

Introduction

GWI in deterministic environment

Random environment

Results

Tail asymptotic

Stationary Markov chain

Stochastic recurrence equation

Perpetuity equation

Goldie's implicit renewal theory

$(A_n, B_n)_n$ iid random vectors, and X_0 a random variable independent of them. The stochastic recurrence equation is

$$X_{n+1} = A_{n+1}X_n + B_{n+1}.$$

The stationary solution should be

$$X_\infty = B_1 + A_1B_2 + \dots + A_1A_2\dots A_nB_{n+1} + \dots =: \sum_{n=0}^{\infty} \Pi_n B_{n+1}.$$

Satisfies the fixed point equation

$$X \stackrel{\mathcal{D}}{=} AX + B,$$

where (A, B) and X on the RHS are independent.

Tail of the stationary distribution

Theorem (Grincevičius – Kesten – Goldie)

If $\mathbf{E}A^\kappa = 1$, $\mathbf{E}A^\kappa \log_+ A < \infty$, $\log A$ is nonarithmetic, $\mathbf{E}B^\kappa < \infty$
then for the solution to the equation $X \stackrel{\mathcal{D}}{=} AX + B$ we have

$$\mathbf{P}(X > x) \sim cx^{-\kappa},$$

with $c > 0$.

Tail of the stationary distribution

Theorem (Grincevičius – Grey)

If $A \geq 0$, $\mathbf{E}A^\kappa < 1$, $\mathbf{E}A^{\kappa+\epsilon} < \infty$ then the tail of X is regularly varying with parameter $-\kappa$ if and only if the tail of B is.

Damek & Kołodziejek 2020: Between Kesten and Grincevičius – Grey

Outline

Introduction

GWI in deterministic environment

Random environment

Results

Tail asymptotic

Stationary Markov chain

Stochastic recurrence equation

Perpetuity equation

Goldie's implicit renewal theory

Goldie's setup - stochastic fixed point equations

$$X_{n+1} = \sum_{i=1}^{X_n} A_i^{(n+1)} + B_{n+1} =: \theta_{n+1} \circ X_n + B_{n+1}, \quad n \geq 0,$$

X stationary law:

$$X \stackrel{\mathcal{D}}{=} \sum_{i=1}^X A_i + B = \theta \circ X + B$$

(θ, B) and X are independent.

Examples

- ▶ Perpetuity: $X \stackrel{\mathcal{D}}{=} AX + B$, (A, B) and X are independent.

Examples

- ▶ Perpetuity: $X \stackrel{\mathcal{D}}{=} AX + B$, (A, B) and X are independent.
- ▶ Supremum of RW with negative drift: $X \stackrel{\mathcal{D}}{=} AX \vee B$.

Examples

- ▶ Perpetuity: $X \stackrel{\mathcal{D}}{=} AX + B$, (A, B) and X are independent.
- ▶ Supremum of RW with negative drift: $X \stackrel{\mathcal{D}}{=} AX \vee B$.
- ▶ $X \stackrel{\mathcal{D}}{=} \sum_{i=1}^X A_i + B$

Stationary distributions of a Markov chain.

Examples

- ▶ Perpetuity: $X \stackrel{\mathcal{D}}{=} AX + B$, (A, B) and X are independent.
- ▶ Supremum of RW with negative drift: $X \stackrel{\mathcal{D}}{=} AX \vee B$.
- ▶ $X \stackrel{\mathcal{D}}{=} \sum_{i=1}^X A_i + B$

Stationary distributions of a Markov chain.

Buraczewski, Damek, Mikosch: Stochastic models with power law tails. The equation $X = AX + B$. (2016)

Iksanov: Renewal theory for perturbed random walks and similar processes. (2016)

General: $X \stackrel{\mathcal{D}}{=} \Psi(X)$, where $\Psi : \mathbb{R} \times \Omega \rightarrow \mathbb{R}$ random operator,
independent of X .

General: $X \stackrel{\mathcal{D}}{=} \Psi(X)$, where $\Psi : \mathbb{R} \times \Omega \rightarrow \mathbb{R}$ random operator, independent of X .

Assume: $A \geq 0$, $\mathbf{E}A^\kappa = 1$ for some $\kappa > 0$, $\mathbf{E}A^\kappa \log^+ A < \infty$, $\log A$ is not arithmetic.

General: $X \stackrel{\mathcal{D}}{=} \Psi(X)$, where $\Psi : \mathbb{R} \times \Omega \rightarrow \mathbb{R}$ random operator, independent of X .

Assume: $A \geq 0$, $\mathbf{E}A^\kappa = 1$ for some $\kappa > 0$, $\mathbf{E}A^\kappa \log^+ A < \infty$, $\log A$ is not arithmetic.

Theorem (Goldie (1991), Grincevicius (1975))

X is the solution to $X \stackrel{\mathcal{D}}{=} \Psi(X)$, assume

$\mathbf{E}|(\Psi(X))^\kappa - (AX)^\kappa| < \infty$. Then $\mathbf{P}(X > x) \sim cx^{-\kappa}$, where $c = \mathbf{E}(\Psi(X)^\kappa - (AX)^\kappa) / \mathbf{E}(A^\kappa \log A) \geq 0$.

$$\mathbf{P}(X > x) \sim cx^{-\kappa}$$

- ▶ Problem: $c = 0$ is possible!
- ▶ If $\mathbf{E}X^\kappa < \infty$, then $c = 0$.
- ▶ Idea: $\Psi(x) \sim Ax$, $x \rightarrow \infty$. ($x \rightarrow \pm\infty$)
- ▶ Alsmeyer, Brofferio, Buraczewski: Asymptotically linear iterated function systems on the real line (2023)
- ▶ K (2016): additional slowly varying factor, or $\mathbf{E}A^\kappa < 1$ is possible